

Contents

Chapter 1 Basic knowledge	1
1.1 Calaulation of sets	1
1.1.1 Intersecton and union	1
1.1.2 Limit of Set Sequences	3
1.1.3 Direct product of sets	4
1.2 Mappings and cardinality	4
Chapter 2 Measure theory and probability spaces	7
2.1 Algebras and σ -algebras	7
2.2 Measurable spaces	9
2.3 Extension Theorem	13
2.4 Inclusion and exclusion principle	13
2.5 Conditional probability	15
2.5.1 Definitons	15
2.5.2 Multiplication formula	16
2.5.3 Total probability formula	17
2.5.4 Bayes formula	18
Chapter 3 Random variables and their distribution	25
3.1 Random variables; Distribution function	25
3.2 Properties of Borel functions	28
3.3 σ -algebra generated by a random variable	29
3.4 Independence	30
Chapter 4 Discrete and continuous random variables	33
4.1 Definitions	33
4.2 Expectation and variance	33
4.3 Lebesgue integration	38
4.4 Absoulte continuity and Radon-Nikodym derivative	41
4.5 Common discrete distribution	43
4.6 Inequalities	53
4.6.1 Chebychev	53
4.6.2 Cauchy-Schwarz	54
Chapter 5 Multi-dimensional random variables	56
5.1 Joint distribution function	56
5.2 Conditional distribution	62
5.2.1 Discrete cases	62
5.2.2 Continuous cases	63
5.2.3 Total probability formula; Bayes formula	65
5.3 Conditional expectation	65

5.3.1	Discrete cases	65
5.3.2	Continuous cases	67
Chapter 6	Random Walk	71
Chapter 7	Functions of random variables	78
7.1	Functions of random variables	78
7.2	Generating functions	83
7.3	Galton-Walter process	90
7.4	Characteristic functions	92
7.5	Common characteristic functions	96
7.6	Inversion and continuity	97
Chapter 8	Convergence of random variables	102
8.1	Different modes of convergence	102
8.2	Tail events	108

Chapter 1 Basic knowledge

1.1 Calculation of sets

1.1.1 Intersecton and union


Let Ω be the universe.

Definition 1.1

If a set contains all elements of A, B, \dots , it is called the **union** of A, B, \dots , denoted as $A \cup B \cup \dots$. 

The operation of union is commutative and associative.

Definition 1.2

Let A, B, \dots be a set sequence. The set, which consists of elements common to A, B, \dots is called the **intersection** of A, B, \dots , denoted as $A \cap B \dots$ (or $AB \dots$). 

Definition 1.3

The set which consists of elements in Ω but not in A , is called the **complement** of A , denoted as A^c . 

For every element in Ω , either it belongs to A or it belongs to A^c , i.e. $x \notin A \iff x \in A^c$. Of course, $(A^c)^c = A$.


One can verify that

$$(A \cup B)^c = A^c \cap B^c, \quad (A \cap B)^c = A^c \cup B^c.$$


Similarly,

$$\left(\bigcup A_n\right)^c = \bigcap A_n^c, \quad \left(\bigcap A_n\right)^c = \bigcup A_n^c.$$

In fact, suppose that $x \in \bigcup A_n$, then $x \in A_n$ for some n . $x \in \left(\bigcup A_n\right)^c$ implies that for all n , $x \notin A_n$, i.e., $x \in A_n^c$. Thus $x \in \bigcap A_n^c$. Hence $\bigcup A_n$ is the complement of $\bigcap A_n^c$. One can obtain the second equality by substituting A_n^c with A_n .



Definition 1.4

The set consists of elements that belong to A but not belong to B is denoted as $A - B$. 

One can see that

$$A - B = A - AB = AB^c = B^c - A^c, \tag{1.1}$$

$$B - A = B - AB = BA^c = A^c - B^c,$$

$$A \cup B = (A - B) + (B - A) + AB.$$

Note that the three sets in the right hand side of the last equality are disjoint.

Theorem 1.1

Let A, B and C be three sets. Then one has the commutative law

$$A \cup B = B \cup A, \quad A \cap B = B \cap A; \quad (1.2)$$

the associative law

$$\begin{aligned} A \cup (B \cup C) &= (A \cup B) \cup C, \\ A \cap (B \cap C) &= (A \cap B) \cap C; \end{aligned} \quad (1.3)$$

and the distributive law

$$\begin{aligned} (A \cup B) \cap C &= (A \cap C) \cup (B \cap C), \\ (A \cap B) \cup C &= (A \cup C) \cap (B \cup C). \end{aligned} \quad (1.4)$$



Similarly, for set sequence, one has

$$\begin{aligned} \bigcap_i \left(\bigcup_j A_j^{(i)} \right) &= \bigcup \left(A_{j_1}^{(1)} \cap A_{j_2}^{(2)} \cap A_{j_3}^{(3)} \cap \dots \right), \\ \bigcup_i \left(\bigcap_j A_j^{(i)} \right) &= \bigcap \left(A_{j_1}^{(1)} \cup A_{j_2}^{(2)} \cup A_{j_3}^{(3)} \cup \dots \right), \end{aligned}$$

where j_1, j_2, \dots is a permutation of $1, 2, \dots$.

Theorem 1.2

One can use intersection and union to obtain the product of sets, i.e.,

$$\bigcap_{n \geq 1} A_n = A_1 - \bigcup_{n \geq 2} (A_1 - A_n). \quad (1.5)$$



Exercise 1.1

- Let A, B be two subsets of the whole set X . If for any $E \subset X$, one has $E \cap A = E \cup B$, prove that $A = X, B = \emptyset$.
- Let A, B be two sets. Prove that $A = B \iff$ there is set C such that

$$A \cap C = B \cap C, \quad A \cup C = B \cup C.$$

Proof. 1. Take $E = X$, we get $A = X$. Take $E = A^c$, we get $B = \emptyset$.

2. \implies . Take $C = A$.

\impliedby . From

$$A = A \cap (C \cup C^c) = (A \cap C) \cup (A \cap C^c),$$

we know $A \cup C = (A \cap C^c) \cup C$. Similarly, $B \cup C = (B \cap C^c) \cup C$. Note that $C \cap (A \cap C^c) = \emptyset = C \cap (B \cap C^c)$, hence $A \cap C^c = B \cap C^c$, implying

$$A = (A \cap C) \cup (A \cap C^c) = (B \cap C) \cup (B \cap C^c) = B.$$

□

1.1.2 Limit of Set Sequences

Definition 1.5

For a set sequence $\{A_n\}$, let

$$B_j = \bigcup_{k=j}^{\infty} A_k (j = 1, 2, \dots), \quad C_j = \bigcap_{k=j}^{\infty} A_k (j = 1, 2, \dots),$$

then $B_j \supset B_{j+1}, C_j \subset C_{j+1}, (j = 1, 2, \dots)$. We define its *lime superior* $\overline{\lim} A_n$ and its *lime inferior* $\underline{\lim} A_n$ as

$$\begin{aligned} \overline{\lim} A_n &= \lim_{k \rightarrow \infty} B_k = \bigcap_{j=1}^{\infty} B_j = \bigcap_{n \geq 1} \left(\bigcup_{i \geq n} A_i \right), \\ \underline{\lim} A_n &= \lim_{k \rightarrow \infty} C_k = \bigcup_{j=1}^{\infty} C_j = \bigcup_{n \geq 1} \left(\bigcap_{i \geq n} A_i \right). \end{aligned} \quad (1.6)$$

When $\overline{\lim} A_n = \underline{\lim} A_n$, this result is defined as $\lim A_n$.



In convention, one usually uses “ A_n occurs infinitely often” to read $\overline{\lim} A_n$ and “ A_n occurs almost always” to read $\underline{\lim} A_n$. Consider the following statements:

$$\begin{aligned} x \in \text{infinitely } A_n\text{'s} &\iff \forall p \in \mathbb{N}, \exists n \geq p, \text{ such that } x \in A_n \\ &\iff \forall p \in \mathbb{N}, x \in \bigcup_{n \geq p} A_n \\ &\iff x \in \bigcap_{p \in \mathbb{N}} \left(\bigcup_{n \geq p} A_n \right) \end{aligned}$$

and

$$\begin{aligned} x \text{ is in all but a finite number of the } A_n\text{'s} &\iff \exists m_x \in \mathbb{N}, \forall n > m_x, x \in A_n \\ &\iff x \in \bigcup_{m_x \geq 1} \left(\bigcap_{n=m_x+1}^{\infty} A_n \right). \end{aligned}$$

One can see that $\overline{\lim}, \underline{\lim}$ have nothing to do with the arrangement of A_n . Actually, $\overline{\lim} A_n$ is the collection of elements that are common to **infinitely** A_n and $\underline{\lim} A_n$ is the collection of elements that are common to a ‘deleted’ A_n (i.e. **finite** A_k ’s are deleted from the original $\{A_n\}$). Hence, we have $\liminf A_n \subset \limsup A_n$.

When $\{A_n\}$ is increasing, i.e., $A_1 \subset A_2 \subset \dots \subset A_n \subset \dots$,

$$\lim A_n = \bigcup_{n \geq 1} A_n. \quad (1.7)$$

When $\{A_n\}$ is decreasing, i.e., $A_1 \supset A_2 \supset \dots \supset A_n \supset \dots$,

$$\lim A_n = \bigcap_{n \geq 1} A_n. \quad (1.8)$$

Generally, we have

$$\overline{\lim} A_n = \lim_{n \rightarrow \infty} \bigcup_{i \geq n} A_i, \quad \underline{\lim} A_n = \lim_{n \rightarrow \infty} \bigcap_{i \geq n} A_i. \quad (1.9)$$

For the complement, we have

$$(\overline{\lim} A_n)^c = \underline{\lim} A_n^c, \quad (\underline{\lim} A_n)^c = \overline{\lim} A_n^c. \quad (1.10)$$

Example 1.1 Let E, F be two sets. Consider the set sequence

$$A_F = \begin{cases} E, & k \text{ is odd,} \\ F, & k \text{ is even} \end{cases} \quad k = (1, 2, \dots),$$

then we have

$$\overline{\lim}_{k \rightarrow \infty} A_k = E \cup F, \quad \underline{\lim}_{k \rightarrow \infty} A_k = E \cap F.$$

Example 1.2 Let $\{f_n(x)\}$ and $f(x)$ be real-valued functions defined on \mathbb{R} . Then D , the set of points where $f_n(x)$ does not converge to $f(x)$ can be described as

$$D = \bigcup_{k=1}^{\infty} \bigcap_{N=1}^{\infty} \bigcup_{n=N}^{\infty} \left\{x : |f_n(x) - f(x)| \geq \frac{1}{k}\right\}.$$

Indeed, if $f_n(x)$ does not converge to $f(x)$ on x_0 , for a fixed $\varepsilon_0 > 0$, for any natural number k , there is $n \geq k$ such that

$$|f_n(x) - f(x)| \geq \varepsilon_0,$$

i.e., x_n is in infinitely many E_n 's, where

$$E_n(\varepsilon_0) := \{x : |f_n(x) - f(x)| \geq \varepsilon_0\},$$

namely, x_n is in the lime superior of $\{E_n(\varepsilon_0)\}$. Moreover, taking union of ε , the discontinuous points are given by the union of these lime superiors. Finally, we use a decreasing sequence $\{\varepsilon_k = 1/k\}$ to supersede ε and reach our desired D .

Example 1.3 Let $\{f_n(x)\}$ and $f(x)$ be real-valued functions defined on \mathbb{R} with

$$\lim_{n \rightarrow \infty} f_n(x) = f(x), \quad x \in \mathbb{R}.$$

Then for $t \in \mathbb{R}$,

$$\begin{aligned} & \{x \in \mathbb{R} : f(x) \leq t\} \\ &= \bigcap_{k=1}^{\infty} \bigcup_{m=1}^{\infty} \bigcap_{n=m}^{\infty} \left\{x \in \mathbb{R} : f_n(x) < t + \frac{1}{k}\right\}. \end{aligned}$$

1.1.3 Direct product of sets

Definition 1.6

Let X, Y be two sets. The collection of all ordered "element pair" (x, y) (where $x \in X, y \in Y$) is called the **direct product** of X and Y , denoted by $X \times Y$, i.e.,

$$X \times Y = \{(x, y) : x \in X, y \in Y\},$$

where $(x, y) = (x', y')$ refers to $x = x', y = y'$. $X \times X$ is also denoted by X^2 .




1.2 Mappings and cardinality

Definition 1.7

Let X, Y be two non-empty sets. If for every $x \in X$, there is a unique $y \in Y$ corresponding to it, we call this corresponding relation a **mapping**. If we use f to denote this relationship, we write

$$f : X \rightarrow Y,$$

and f is called a mapping from X to Y . Now the unique $y \in Y$ corresponding to a certain $x \in X$ is called the **image** of x under f whereas x is called a **pre-image** of y , denoted by $f(x) = y$. If for any $y \in Y$, there is $x \in X$ such that $f(x) = y$, f is called a **surjection** of f from X to Y . 

For $f : X \rightarrow Y$ and $A \subset X$, we write

$$f(A) = \{y \in Y : x \in A, y = f(x)\}$$

with stipulation $f(\emptyset) = \emptyset$. Clearly,

$$(i) f\left(\bigcup_{\alpha \in I} A_\alpha\right) = \bigcup_{\alpha \in I} f(A_\alpha);$$

$$(ii) f\left(\bigcap_{\alpha \in I} A_\alpha\right) \subset \bigcup_{\alpha \in I} f(A_\alpha).$$

For $f : X \rightarrow Y$ and $B \subset Y$, we write

$$f^{-1}(B) = \{x \in X : f(x) \in B\}.$$

Similarly,

$$(i) \text{ for } B_1 \subset B_2, \text{ we have } f^{-1}(B_1) \subset f^{-1}(B_2);$$

$$(ii) f^{-1}\left(\bigcup_{\alpha \in I} B_\alpha\right) = \bigcup_{\alpha \in I} f^{-1}(B_\alpha) (B_\alpha \subset Y, \alpha \in I);$$

$$(iii) f^{-1}\left(\bigcap_{\alpha \in I} B_\alpha\right) = \bigcap_{\alpha \in I} f^{-1}(B_\alpha) (B_\alpha \subset Y, \alpha \in I);$$

$$(iv) f^{-1}(B^c) = (f^{-1}(B))^c (B \subset Y).$$


Definition 1.8

Let $f : X \rightarrow Y$. If f satisfies that for $x_1, x_2 \in X$ and $x_1 \neq x_2$,

$$f(x_1) \neq f(x_2),$$

f is called an **injection** from X to Y . If f is both injective and surjective, f is called a **bijection** from X to Y . Take f bijective from X to Y , then for every $y \in Y$, there is a unique $x \in X$ such that $f(x) = y$, i.e., there is a mapping

$$g : Y \rightarrow X, \quad g(y) = x,$$

where x is determined by $y = f(x)$. g is then called the **inverse** of f , denoted as f^{-1} . 

Definition 1.9

The **indicator mapping** of set A , denoted as $\mathbb{1}_A$, is defined as

$$\mathbb{1}_A(x) := \begin{cases} 1, & \text{if } x \in A \\ 0, & \text{if } x \notin A. \end{cases} \quad (1.11) \quad \text{img alt="clover icon" data-bbox="888 773 903 786}$$

Example 1.4 Show that

$$\mathbb{1}_{\limsup A_n} = \limsup \mathbb{1}_{A_n}, \text{ and } \mathbb{1}_{\liminf A_n} = \liminf \mathbb{1}_{A_n}.$$

Proof. Observe that both mappings $\mathbb{1}_{\limsup A_n}$ and $\limsup \mathbb{1}_{A_n}$ only take values in $\{0, 1\}$.

For an arbitrary $x \in \Omega$, consider the following statements:

$$\mathbb{1}_{\limsup A_n} = 1 \iff x \in \overline{\lim A_n}$$

$$\begin{aligned} &\iff \forall p \in \mathbb{N}, \exists n \geq p, \text{ such that } x \in A_n \\ &\iff \forall p \in \mathbb{N}, \exists n \geq p, \text{ such that } \mathbb{1}_{A_n}(x) = 1 \\ &\iff \limsup \mathbb{1}_{A_n}(x) = 1. \end{aligned}$$

□

Definition 1.10

Let X be a non-empty set. The collection of all subsets of X (including \emptyset and X) is called the **power set** of X , denoted as $\mathcal{P}(X)$.



For example, there are 2^n elements in the power set of an n -element set E .

Example 1.5(Fixed points of monotonous mappings) Let X be a non-empty set. Let $f : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ satisfying $f(A) \subset f(B)$ for any $A \subset B \in \mathcal{P}(X)$, then there is $T \subset \mathcal{P}(X)$ such that $f(T) = T$.

Proof. Consider sets S and T , where

$$\begin{aligned} S &= \{A : A \in \mathcal{P}(X) \text{ and } A \subset f(A)\}, \\ T &= \bigcup_{A \in S} A (\in \mathcal{P}(X)). \end{aligned}$$

I claim that $f(T) = T$.

Indeed, from $A \in S$ we know that $A \subset f(A)$. Moreover, $A \subset T$, implying that $f(A) \subset f(T)$, hence we deduce $f(A) \subset f(T)$ from $A \in S$, following

$$\bigcup_{A \in S} A \subset f(T), \quad T \subset f(T).$$

On the other hand, based on $T \subset f(T)$, we know $f(T) \subset f(f(T))$, implying $f(T) \in S$ thus $f(T) \subset T$. □

Chapter 2 Measure theory and probability spaces

2.1 Algebras and σ -algebras

Definition 2.1

Let E be a set and let \mathcal{A} be a **set of subsets** of E . We say that \mathcal{A} is an algebra if for all $B, A \in \mathcal{A}$,

$$\emptyset \in \mathcal{A}, A^c \in \mathcal{A}, \text{ and } A \cup B \in \mathcal{A}.$$

We say that \mathcal{A} is a π -system if for all $B, A \in \mathcal{A}$,

$$\emptyset \in \mathcal{A}, \text{ and } A \cap B \in \mathcal{A}.$$

We say that \mathcal{A} is a λ -system if

$$\emptyset \in \mathcal{A}, \text{ and } B \setminus A \in \mathcal{A} \text{ given } A, B \in \mathcal{A}, A \subseteq B, \text{ and}$$

$$A \in \mathcal{A} \text{ given } A_n \uparrow A, A_n \in \mathcal{A}.$$

We say that \mathcal{A} is a σ -algebra if for all $A \in \mathcal{A}$, and all sequences $\{A_n\}_{n \in \mathbb{N}}$,

$$\emptyset \in \mathcal{A}, A^c \in \mathcal{A}, \text{ and } \bigcup_{n \in \mathbb{N}} A_n \in \mathcal{A}.$$



Immediately, we have the following lemmas:

Lemma 2.1

Every σ -algebra is an algebra.



Lemma 2.2

Every algebra is a π -system.



Example 2.1 $\mathcal{A} = \{\emptyset, E\}$ is a(n) (σ -)algebra, usually called the trivial (σ -)algebra.

Example 2.2 The power set $\mathcal{P}(E) := \{A : A \subset E\}$ is a(n) (σ -)algebra.

Lemma 2.3

The intersection of any collection of (σ -)algebras is a(n) (σ -)algebra, i.e., if \mathcal{A}_1 and \mathcal{A}_2 are both (σ -)algebras, then their intersection $\mathcal{A}_1 \cap \mathcal{A}_2$ is also a(n) (σ -)algebra.



Definition 2.2

Note that the intersection of any collection of σ -algebras is a σ -algebra. Thus for any set of subsets \mathcal{C} the intersection of all the σ -algebras (there is at least one) containing \mathcal{C} is itself a σ -algebra. We call this the σ -algebra generated by \mathcal{C} , and we denote it by $\sigma(\mathcal{C})$. Similarly, the intersection of all the algebras (there is at least one) containing \mathcal{C} is itself an algebra. We call this the algebra generated by \mathcal{C} , and we denote it by $a(\mathcal{C})$. To be more precise,


$$\sigma(\mathcal{C}) = \bigcap_{\substack{\mathcal{A} \text{ is a } \sigma\text{-algebra} \\ \mathcal{C} \subseteq \mathcal{A}}} \mathcal{A};$$

$$a(\mathcal{C}) = \bigcap_{\substack{\mathcal{A} \text{ is an algebra} \\ \mathcal{C} \subseteq \mathcal{A}}} \mathcal{A}.$$




Based on definition 2.2, we know

Proposition 2.1

$\sigma(\mathcal{C})$ is the smallest σ -algebra containing \mathcal{C} whereas $a(\mathcal{C})$ is the smallest algebra containing \mathcal{C} . 

Proposition 2.2

Let \mathcal{C} be a set of subsets. Then

$$a(\mathcal{C}) \subseteq \sigma(\mathcal{C}), \quad \sigma(a(\mathcal{C})) = \sigma(\mathcal{C}).$$


Proof. The first assertion is trivial since every σ -algebra is an algebra. For the second one, first note that $\mathcal{C} \subseteq a(\mathcal{C})$ thus $\sigma(a(\mathcal{C}))$ is actually a σ -algebra containing \mathcal{C} , implying $\sigma(\mathcal{C}) \subseteq \sigma(a(\mathcal{C}))$. Conversely, since $a(\mathcal{C}) \subseteq \sigma(\mathcal{C})$, we know that $\sigma(\mathcal{C})$ is a σ -algebra containing $a(\mathcal{C})$. Hence $\sigma(a(\mathcal{C})) \subseteq \sigma(\mathcal{C})$. \square

Example 2.3 For $A \subseteq E$, let $\mathcal{C} = \{\emptyset, A\}$. Then $a(\mathcal{C}) = \{\emptyset, A, A^c, E\}$.

Proof. We use the shorthand $\mathcal{F} = \{\emptyset, A, A^c, E\}$. It is easy to observe that \mathcal{F} is an algebra containing \mathcal{C} . Then $a(\mathcal{C}) \subseteq \mathcal{F}$ according to the “smallest”, implying $a(\mathcal{C}) \subseteq \mathcal{F}$.

On the other hand, for every member in \mathcal{F} , they are in $a(\mathcal{C})$, implying $\mathcal{F} \subseteq a(\mathcal{C})$. \square

Example 2.4 Let $\pi = \{A_1, A_2, \dots, A_n\}$ be a partition of E , i.e., $A_i \cap A_j = \emptyset$ and $\bigcup_{i=1}^m A_i = E$. Then

$$\begin{aligned} a(\pi) &= \{\text{finite disjoint unions of } \{A_i\}_{i=1}^m\} \\ &= \{\bigcup_{i \in I} A_i \text{ for some } I \subseteq \{1, 2, \dots, m\}\}. \end{aligned}$$

Example 2.5 Let \mathcal{A} be an algebra of \mathbb{R} and $X : E \rightarrow \mathbb{R}$ be a set function. Then $\{X^{-1}(A) : A \in \mathcal{A}\}$ is an algebra where $X^{-1}(A) := \{\omega \in E : X(\omega) \in A\}$.

Example 2.6 Let \mathcal{E} , a set of subset of \mathbb{R} be $\mathcal{E} = \{(a, b) : -\infty \leq a < b \leq +\infty, (a, +\infty) : a \in \mathbb{R}\}$. Then $a(\mathcal{E}) = \{I_1 \cup I_2 \cup \dots \cup I_n, I_k \in \mathcal{E}, I_i \cap I_j = \emptyset, 1 \leq i, j, k \leq n\}$.

Example 2.7(π -systems generating $\mathcal{B}(\mathbb{R})$) The followings are four methods to generate $\mathcal{B}(\mathbb{R})$:

1. $\mathcal{E}_1 := \{(a, b) : -\infty \leq a \leq b < +\infty\}$ is a π -system.
2. $\mathcal{E}_2 := \{(a, b) : -\infty \leq a \leq b \leq +\infty\}$ is a π -system.
3. $\mathcal{E}_3 := \{A \subset \mathbb{R} : A \text{ is open}\}$ is a π -system.
4. $\mathcal{E}_4 := \{A \subset \mathbb{R} : A \text{ is closed}\}$ is a π -system.


Proof. Only for 3. It suffices to show $\sigma(\mathcal{E}_2) = \sigma(\mathcal{E}_3)$. $\sigma(\mathcal{E}_2) \subset \sigma(\mathcal{E}_3)$ is trivial since $\mathcal{E}_2 \subset \mathcal{E}_3$. Conversely, let $A \subseteq \mathbb{R}$ be open, and then

$$A = \bigcup_{i=1}^{\infty} (x_i - \varepsilon_i, x_i + \varepsilon_i)$$


implying $\mathcal{E}_3 \subset \sigma(\mathcal{E}_2)$ and thus $\sigma(\mathcal{E}_3) \subset \sigma(\mathcal{E}_2)$ according to the “smallest”. \square

 **Exercise 2.1** Let \mathcal{A} be an algebra on E . If $A \notin \mathcal{A}$ and $B \notin \mathcal{A}$, is it the same for $A \cup B$ and $A \cap B$?

Solution. Let $E = \{0, 1\}$, $\mathcal{A} = \{\emptyset, E\}$, $A = \{1\}$, $B = \{0\}$. Then $A \cap B = \emptyset \in \mathcal{A}$ and $A \cup B = E \in \mathcal{A}$. \square

 **Exercise 2.2** Provide a counterexample in which $\mathcal{A} \cup \mathcal{B}$ is not an algebra whereas \mathcal{A} and \mathcal{B} are both algebras, separately.


Solution. Set $\Omega = \{0, 1, 2\}, A = \{0\}, B = \{0, 1\}$. Then $a(A) = \{\emptyset, \Omega, A, A^c\}, a(B) = \{\emptyset, \Omega, B, B^c\}$. Note that $a(A) \cup a(B) = \{\emptyset, \{0\}, \{0, 1\}, \{2\}, \{1, 2\}, \Omega\}$ whereas $\{0, 1\} \cap \{1, 2\} = \{1\} \notin a(A) \cup a(B)$. \square

 **Exercise 2.3** Let E be an infinite set (countable or not). Let \mathcal{A} be the set of subsets of E that are either finite or with finite complement in E . Prove that \mathcal{A} is an algebra but not a σ -algebra.

Proof. Take a subset A of E where $A \in \mathcal{A}$. If A is finite, its complement must also be in \mathcal{A} since $(A^c)^c = A$. If A^c is finite, it is the same. This has already implied $\emptyset, E \in \mathcal{A}$.

Now take another $B \subseteq E$ with $B \in \mathcal{A}$. If A and B are both finite, $A \cup B$ is also finite thus in \mathcal{A} . We finished the proof. Otherwise, assume A is not finite. Note that $(A \cup B)^c = A^c \cap B^c$ is finite since A^c is finite. Hence $A \cup B$ is indeed in \mathcal{A} .

Now assume that $\mathbb{N} \subset E$. Consider the even number set $2\mathbb{N} = \{2n, n \in \mathbb{N}\}$. If \mathcal{A} is a σ -algebra, since each $\{n\}$ is in \mathcal{A} , their countable union $2\mathbb{N}$ must also be in \mathcal{A} . However, neither $2\mathbb{N}$ nor $E \setminus 2\mathbb{N}$ is finite. \square

 **Exercise 2.4** Let $\Omega = \mathbb{N}$. For $n \geq 0$, let $\mathcal{F}_n = \sigma(\{\{0\}, \dots, \{n\}\})$, the smallest σ -algebra containing all of $\{0\}, \dots, \{n\}$. Show that $\bigcup_{n \geq 0} \mathcal{F}_n$ is not a σ -algebra.


Proof. Let us write down several \mathcal{F}_n 's.

$$\begin{aligned} \mathcal{F}_0 &= \{\emptyset, \mathbb{N}, \{0\}, \mathbb{N} \setminus \{0\}\}; \\ \mathcal{F}_1 &= \{\emptyset, \mathbb{N}, \{0\}, \{1\}, \mathbb{N} \setminus \{0\}, \mathbb{N} \setminus \{1\}, \mathbb{N} \setminus \{0, 1\}\}; \\ &\vdots \end{aligned}$$

We can observe that for each natural even number $2k, \{2k\} \in \bigcup_{n \geq 0} \mathcal{F}_n$, whereas $2\mathbb{N}$ does not belong to $\bigcup_{n \geq 0} \mathcal{F}_n$. \square

2.2 Measurable spaces


Definition 2.3

A pair (E, \mathcal{E}) , where E is a set and \mathcal{E} is a σ -algebra on E , is called a measurable space. Given a measurable space (E, \mathcal{E}) each $A \in \mathcal{E}$ is called a measurable set (or an E -measurable set). 

Recall the set \mathcal{E} in Example 2.6. We have had an understanding of $a(\mathcal{E})$. Now let's take a look at $\sigma(\mathcal{E})$. First, one can observe that any single point $\{a\}, a \in \mathbb{R}$ is in $\sigma(\mathcal{E})$ since $\{a\} = \bigcup_{n \geq 1} (a - \frac{1}{n}, a]$. Thus every countable set (for example, \mathbb{Q} and transcendental numbers) is also in $\sigma(\mathcal{E})$. Moreover, every open interval (a, b) is in $\sigma(\mathcal{E})$ since $(a, b) = \bigcup_{n \geq 1} (a, b - \frac{1}{n})$. Finally, $[a, b)$ is in $\sigma(\mathcal{E})$ since $[a, b) = \bigcup_{n \geq 1} [a, b - \frac{1}{n})$. Thus, $\sigma(\mathcal{E})$ actually basically contains the "nice subsets of \mathbb{R} ", usually it is called the **Borel set of \mathbb{R}** , denoted as $\mathcal{B}(\mathbb{R})$.

We will keep using the notation \mathcal{E} to represent the set mentioned in Example 2.6 in this section if not specifically stated.

Definition 2.4

Let E be a set, and let \mathcal{A} be an algebra on E . A set function is any function $\mu : \mathcal{A} \rightarrow [0, \infty]$ with $\mu(\emptyset) = 0$. 

Let μ be a set function. We say that μ is increasing if, for all $A, B \in \mathcal{A}$ with $A \subseteq B$,

$$\mu(A) \leq \mu(B).$$

We say that μ is additive if, for all disjoint sets $A, B \in \mathcal{A}$,

$$\mu(A \cup B) = \mu(A) + \mu(B).$$

We say that μ is countably additive if, for all sequences of disjoint sets $\{A_n\}_{n \in \mathbb{N}} \subseteq \mathcal{A}$, with $\bigcup_{n \in \mathbb{N}} A_n \in \mathcal{A}$, we have

$$\mu\left(\bigcup_{n \in \mathbb{N}} A_n\right) = \sum_{n \in \mathbb{N}} \mu(A_n).$$

Definition 2.5

Let \mathcal{A} be an algebra (of subsets of E). A set function $\mu : \mathcal{A} \rightarrow [0, +\infty)$ is called a **content** if

1. $\mu(\emptyset) = 0$;
2. μ is finitely additive.



Lemma 2.4

Let $\mu : \mathcal{A} \rightarrow [0, +\infty)$ be a content. Then for any $A, B \in \mathcal{A}$,

1. $\mu(A \cup B) + \mu(A \cap B) = \mu(A) + \mu(B)$;
2. If $A \subseteq B$ and $\mu(A) < +\infty$ then $\mu(B \setminus A) = \mu(B) - \mu(A)$;
3. If $A \subseteq B$, then $\mu(A) \leq \mu(B)$.
4. (Subadditivity) If $A_1, A_2, \dots, A_n \in \mathcal{A}$, then

$$\mu\left(\bigcup_{i=1}^n A_i\right) \leq \sum_{i=1}^n \mu(A_i).$$

5. (σ -superadditivity) If $\{A_i\}_{i=1}^{+\infty}$ is a sequence of disjoint elements and $\bigcup_{i \geq 1} A_i \in \mathcal{A}$, then

$$\mu\left(\bigcup_{i \geq 1} A_i\right) \geq \sum_{i \geq 1} \mu(A_i).$$



Proof. Only prove 5. By 4, we have for any $N \in \mathbb{N}$

$$\mu\left(\bigcup_{i=1}^N A_i\right) \leq \sum_{i=1}^N \mu(A_i).$$

Thus the superadditivity holds by nature since by 3,

$$\mu\left(\bigcup_{i=1}^{\infty} A_i\right) \geq \mu\left(\bigcup_{i=1}^N A_i\right) \geq \sum_{i=1}^N \mu(A_i).$$

□

Example 2.8 (Density of \mathbb{R}) Let $E = \mathbb{R}$ and $\mathcal{A} = a(\mathcal{E})$. For any $A \in a(\mathcal{E})$, define $\mu(A) = \lim_{L \rightarrow \infty} \frac{|A \cap (0, L)|}{L}$. Then μ is finitely additive but not countably additive. In fact, note that

$$1 = \mu\left(\bigcup_{i \geq 0} [i, i+1)\right) > \sum_{i \geq 0} \mu([i, i+1)) = 0.$$

Definition 2.6

Let (E, \mathcal{E}) be a measurable space. A countably additive set function $\mu : \mathcal{E} \rightarrow [0, \infty]$ is called a **measure**; the triple (E, \mathcal{E}, μ) is called a **measure space**.



Example 2.9 Let $E = \mathbb{R}$. Define a set function $m : a(\mathcal{E}) \rightarrow [0, +\infty]$. Let $m((a, b]) = b - a$, $m((a, +\infty)) = +\infty$ and extend for every $A \in a(\mathcal{E})$, i.e., $m(A) = \sum_{j=1}^n m(I_j)$ where $A = \bigcup_{j=1}^n I_j$ and I_j 's are two by two disjoint.

Then m is actually a content.

Lemma 2.5

The set function m defined in Example 2.9 is countably additive, i.e., for $\{A_k\}_{k=1}^{\infty} \subset a(\mathcal{E})$ such that A_k 's are two by two disjoint and $A = \bigcup_{k \geq 1} A_k \subset a(\mathcal{E})$, one has $m(A) = \sum_{k \geq 1} m(A_k)$.



Proof. We introduce a new partition of A , i.e., set $A = \bigcup_{i=1}^n I_i$, where every $I_i \in \mathcal{E}$ is disjoint with each other.

Moreover, let $A_k = \bigcup_{j=1}^{m_k} J_{j,k}$, where every $J_{j,k} \in \mathcal{E}$ is disjoint with respect to j with each other. Then

$$\begin{aligned} m(A) &= \sum_{i=1}^n m(I_i) = \sum_{i=1}^n \sum_{j=1}^{m_k} \sum_{k \geq 1} m(I_i \cap J_{j,k}) \\ &= \sum_{j=1}^{m_k} \sum_{k \geq 1} m(I_i \cap J_{j,k}) = \sum_{k=1}^{+\infty} m(A_k). \end{aligned}$$

□

Definition 2.7

If $\mu(E) = 1$ then μ is called a probability measure and (E, \mathcal{E}, μ) is called a probability space. The notation (Ω, \mathcal{F}, P) is often used to denote a probability space.



Lemma 2.6

Let (E, \mathcal{F}, μ) be a measure space. Then if $A_1, A_2, \dots \in \mathcal{F}$, one has

$$\mu\left(\bigcup_{i \geq 1} A_i\right) \leq \sum_{i \geq 1} \mu(A_i).$$



Proof. Define $B_1 = A_1, B_2 = A_2 \setminus A_1, \dots, B_n = A_n \setminus \bigcup_{i=1}^{n-1} A_i, \dots$. Then the sequence $\{B_i\}$ satisfies:

- each $B_i (i = 1, 2, \dots, n, \dots)$ is disjoint with each other;
- $\bigcup_{i \geq 1} B_i = A$;
- $\mu(B_i) \leq \mu(A_i)$

Thus

$$\mu(A) = \mu\left(\bigcup_{i \geq 1} B_i\right) = \sum_{i \geq 1} \mu(B_i) \leq \sum_{i \geq 1} \mu(A_i).$$

□

Theorem 2.1 (Continuity of measures)

Let μ be a measure and A_1, A_2, \dots be an increasing sequence of events, so that $A_1 \subseteq A_2 \subseteq A_3 \subseteq \dots$, and write A for their limit:

$$A = \bigcup_{i=1}^{\infty} A_i = \lim_{i \rightarrow \infty} A_i.$$

Then $\mu(A) = \lim_{i \rightarrow \infty} \mu(A_i)$. Similarly, if B_1, B_2, \dots is a decreasing sequence of events, so that $B_1 \supseteq B_2 \supseteq B_3 \supseteq \dots$, then

$$B = \bigcap_{i=1}^{\infty} B_i = \lim_{i \rightarrow \infty} B_i$$

satisfies $\mu(B) = \lim_{i \rightarrow \infty} \mu(B_i)$.



Proof. $A = A_1 \cup (A_2 \setminus A_1) \cup (A_3 \setminus A_2) \cup \dots$ is the union of a disjoint family of events. Thus, by definition,

$$\begin{aligned} \mu(A) &= \mu(A_1) + \sum_{i=1}^{\infty} \mu(A_{i+1} \setminus A_i) \\ &= \mu(A_1) + \lim_{n \rightarrow \infty} \sum_{i=1}^{n-1} [\mu(A_{i+1}) - \mu(A_i)] \\ &= \lim_{n \rightarrow \infty} \mu(A_n). \end{aligned}$$

To show the result for decreasing families of events, take complements and use the first part. \square

Definition 2.8

Given a measure space (E, \mathcal{E}, μ) , μ is said to be finite if

$$\mu(E) < \infty$$

and μ is said to be σ -finite if there exist sets $\{E_n\}_{n \in \mathbb{N}} \subseteq E$ such that $\mu(E_n) < \infty$ and $\bigcup_{n \in \mathbb{N}} E_n = E$.



Definition 2.9

An element $F \in \mathcal{E}$ is called μ -null if $\mu(F) = 0$. A statement φ about points $s \in E$ is said to hold almost everywhere if

$$F := \{s \in E : \varphi(s) \text{ is false}\} \in \mathcal{E}, \text{ and } \mu(F) = 0.$$



Exercise 2.5 Recall the definition of limit of a set sequence $\{A_n\}$. Prove that $\mu(A_n) \rightarrow \mu(A)$, where $A = \limsup A_n = \liminf A_n$.

Proof. We use the shorthand $B_n = \bigcup_{i=n}^{\infty} A_i$ and $C_n = \bigcap_{i=n}^{\infty} A_i$. Then $\{B_n\}$ and $\{C_n\}$ are set sequences decreasing and increasing, respectively. Moreover,

$$\begin{aligned} B &:= \limsup A_n = \bigcap_{n \geq 1} \bigcup_{m \geq n} A_m = \bigcap_{n \geq 1} B_n = \lim_{n \rightarrow \infty} B_n \\ C &:= \liminf A_n = \bigcup_{n \geq 1} \bigcap_{m \geq n} A_m = \bigcup_{n \geq 1} C_n = \lim_{n \rightarrow \infty} C_n. \end{aligned}$$


We have that

$$C_n = \bigcap_{i=n}^{\infty} A_i \subseteq A_n \subseteq \bigcup_{i=n}^{\infty} A_i = B_n,$$

and therefore $\mu(C_n) \leq \mu(A_n) \leq \mu(B_n)$. By the continuity of measures, if $C_n \uparrow C$ then $\mu(C_n) \uparrow \mu(C)$, and if $B_n \downarrow B$ then $\mu(B_n) \downarrow \mu(B)$. If $B = C = A$ then

$$\mu(A) = \mu(C) \leq \lim_{n \rightarrow \infty} \mu(A_n) \leq \mu(B) = \mu(A).$$

\square

 **Exercise 2.6** Let m be Lebesgue measure on $[0, 1]$, and $0 \leq a \leq b \leq c \leq d \leq 1$ such that $a + d \geq b + c$. Give an example of a sequence of sets A_1, A_2, \dots in $[0, 1]$, such that $m(\liminf_n A_n) = a$, $\liminf_n m(A_n) = b$, $\limsup_n m(A_n) = c$ and $m(\limsup_n A_n) = d$.

2.3 Extension Theorem

Theorem 2.2 (Caratheodory Extension Theorem)

Let \mathcal{F} be an algebra on Ω and μ be a countably additive content on (Ω, \mathcal{F}) . If μ is σ -finite, i.e., there is $\{A_i\}_{i=1}^{\infty} A_i \subseteq \mathcal{F}$ such that $\mu(A_i) < +\infty$ and $\bigcup_{i=1}^{\infty} A_i = \Omega$, then μ extend to a measure on $(\Omega, \sigma(\mathcal{F}))$.



Example 2.10 (Lebesgue-Stieltjes measure) Let $m_F : a(\mathcal{E}) \rightarrow \mathbb{R}$ be a set function such that $m_F((a, b]) = F(b) - F(a)$, where $F : \mathbb{R} \rightarrow [0, 1]$ is a right-continuous increasing function with convention $F(+\infty) = \sup\{F(x), x \in \mathbb{R}\}$, $F(-\infty) = \inf\{F(x), x \in \mathbb{R}\}$. Extend m_F to $a(\mathcal{E})$ by finite subadditivity, m_F is countable additive. Hence by the Extension Theorem, m_F extends to a measure on $(\mathbb{R}, \sigma(\mathcal{E}))$. That is the Lebesgue-Stieltjes measure.

2.4 Inclusion and exclusion principle

Proposition 2.3 (Inclusion and Exclusion)

$$\mathbb{P}\left(\bigcup_{1 \leq j \leq n} A_j\right) = \sum_j \mathbb{P}(A_j) - \sum_{j_1 < j_2} \mathbb{P}(A_{j_1} \cap A_{j_2}) + \dots + (-1)^{n-1} \sum_{j_1 < \dots < j_n} \mathbb{P}(A_{j_1} \cap \dots \cap A_{j_n}). \quad (2.1)$$



Proof. Apply induction on n . For the case $n = 2$, it is trivial that the formula is true. Let $m \geq 2$ and suppose that the result is true for $n \leq m$. Then it is true for pair of events, so that

$$\begin{aligned} \mathbb{P}\left(\bigcup_{i=1}^{m+1} A_i\right) &= \mathbb{P}\left(\bigcup_{i=1}^m A_i\right) + \mathbb{P}(A_{m+1}) - \mathbb{P}\left\{\left(\bigcup_{i=1}^m A_i\right) \cap A_{m+1}\right\} \\ &= \mathbb{P}\left(\bigcup_{i=1}^m A_i\right) + \mathbb{P}(A_{m+1}) - \mathbb{P}\left\{\bigcup_{i=1}^m (A_i \cap A_{m+1})\right\}. \end{aligned}$$

Using the induction hypothesis, we may expand the two relevant terms on the right-hand side to obtain the result. \square

Remark If we truncate (2.1), we may obtain several inequalities:

1. $\mathbb{P}(E_1 \cup E_2 \cup \dots \cup E_n) \geq \sum_{i=1}^n \mathbb{P}(E_i) - \sum_{i < j} \mathbb{P}(E_i \cap E_j)$;
2. $\mathbb{P}(E_1 \cup E_2 \cup \dots \cup E_n) \leq \sum_{i=1}^n \mathbb{P}(E_i) - \sum_{i < j} \mathbb{P}(E_i \cap E_j) + \sum_{i < j < k} \mathbb{P}(E_i \cap E_j \cap E_k)$.

Example 2.11 (Birthday problem) Consider a group of n people. What is the probability that at least two of them have the same birthday?

Solution. We start by considering the complement, i.e., working with the condition where no one has the same

birthday. Indeed,

$$\begin{aligned} \mathbb{P}(\text{no one has the same birthday}) &= \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{(365)^n} \\ &= 1 \cdot \left(1 - \frac{1}{365}\right) \cdot \left(1 - \frac{1}{365}\right) \cdots \left(1 - \frac{n-1}{365}\right). \end{aligned}$$

If we consider the approximation $e^{-x} \approx 1 - x$, the final result becomes

$$e^{-1/365} \cdot e^{-1/365} \cdots e^{-(n-1)/365} = e^{-n(n-1)/730}.$$

□

Example 2.12(Matching problem) n people are to pick n hats. What is the probability that no one picks his/her own hat?

Solution. Still consider complement. Define E_i as $E_i = \{\text{the } i\text{-th person gets his/her own hat}\}$. We see that

$$\begin{aligned} \mathbb{P}(E_i) &= \frac{(n-1)!}{n!} = \frac{1}{n}, \quad \mathbb{P}(E_1 \cap E_2 \cap \cdots \cap E_r) = \frac{(n-r)!}{n!}, \\ (-1)^{r-1} \sum_{j_1 < j_2 < \cdots < j_r} \mathbb{P}(E_{j_1} \cap E_{j_2} \cap \cdots \cap E_{j_r}) \\ &= (-1)^{r-1} \binom{n}{r} \frac{(n-r)!}{n!} = (-1)^{r-1} \frac{1}{r!}, \quad 1 \leq r \leq n. \end{aligned}$$

From (2.1),

$$\mathbb{P}(E_1 \cup E_2 \cup \cdots \cup E_n) = \sum_{r=1}^n (-1)^{r-1} \frac{1}{r!}.$$

If we send $n \rightarrow \infty$, the result will converge to e^{-1} and what we want is $1 - e^{-1}$.

□

Example 2.13(Texas Holder) In a poker game, what is the probability that you have

1. a straight?
2. a full house?

The number you can use are from Ace to 10.


Solution. 1. To get a straight, there will be 10 choices of your starting number. For the five card position, there will be $4^5 - 4$ choices in total for separate unit. Thus the probability is

$$\frac{10 \cdot (4^5 - 4)}{C_{52}^5}.$$

2. The probability is

$$\frac{13 \cdot 12 \cdot C_4^3 \cdot C_4^2}{C_{52}^5}.$$

□

 **Exercise 2.7** Six cups and saucers come in pairs: there are two cups and saucers which are red, two white, and two with stars. If the cups are placed randomly onto the saucers (one each), find the probability that no cup is upon a saucer of the same pattern.

Solution. We first place the saucers in a certain order, for example, put it as “RRWWSS”. Note that there are $\frac{6!}{2! \cdot 2! \cdot 2!} = 90$ methods to put on the cups. Now to fulfill the requirement of the problem, a basic idea is to first consider three different cups. That is, for the cup tuple (S, R, W) , its position must be either $S()R()W()$

or $W()S()R()$, where $()$ denotes an empty seat. By enumeration or a little calculation, there are 10 ways in total fulfilling the requirement. Hence the probability is $10/90 = 1/9$. \square

Exercise 2.8 You are given that at least one of the events $A_r, 1 \leq r \leq n$, is certain to occur, but certainly no more than two occur. If $\mathbb{P}(A_r) = p$ and $\mathbb{P}(A_r \cap A_s) = q, r \neq s$, show that $p \geq 1/n$ and $q \leq 2/n$.

Proof. Since at least one of the A_r occurs,

$$\begin{aligned} 1 &= \mathbb{P}\left(\bigcup_{r=1}^n A_r\right) = \sum_r \mathbb{P}(A_r) - \sum_{r < s} \mathbb{P}(A_r \cap A_s) \\ &= np - \frac{n(n-1)}{2}q. \end{aligned}$$

Hence $p \geq 1/n$ and $\frac{1}{2}n(n-1)q = np - 1 \leq n - 1$. \square

Exercise 2.9 You are given that at least one, but no more than three, of the events $A_r, 1 \leq r \leq n$, is certain to occur, where $n \geq 3$. The probability of at least two occurring is $1/2$. If $\mathbb{P}(A_r) = p$ and $\mathbb{P}(A_r \cap A_s) = q, r \neq s$, and $\mathbb{P}(A_r \cap A_s \cap A_t) = x, r < s < t$. Show that $p \geq 3/(2n)$ and $q \leq 4/n$.

Proof. Similarly first we have

$$\begin{aligned} 1 &= \mathbb{P}\left(\bigcup_{r=1}^n A_r\right) = \sum_r \mathbb{P}(A_r) - \sum_{r < s} \mathbb{P}(A_r \cap A_s) + \sum_{r < s < t} \mathbb{P}(A_r \cap A_s \cap A_t) \\ &= np - \binom{n}{2}q + \binom{n}{3}x. \end{aligned}$$

Since at least two of the event occur with probability $\frac{1}{2}$,

$$\frac{1}{2} = \mathbb{P}\left(\bigcup_{r < s} (A_r \cap A_s)\right) = \sum_{r < s} \mathbb{P}(A_r \cap A_s) - \frac{1}{2} \sum_{\substack{r < s \\ t < u \\ (r,s) \neq (t,u)}} \mathbb{P}(A_r \cap A_s \cap A_t \cap A_u).$$

Since no more than three events can occur, in the item $A_r \cap A_s \cap A_t \cap A_u$, there must be a repetition in index and there are four choices for the repeated one. Hence one deduces that

$$\mathbb{P}(A_r \cap A_s \cap A_t \cap A_u) = 4\mathbb{P}(A_r \cap A_s \cap A_t).$$

Substitute back,

$$\frac{1}{2} = \binom{n}{2}q - 2\binom{n}{3}x.$$

Hence $\frac{3}{2} = np - C_n^3 x$ so that $p \geq 3/(2n)$. Also $C_n^2 q = 2np - \frac{5}{2}$ whence $q \leq 4/n$. \square

2.5 Conditional probability

2.5.1 Definitions

Definition 2.10

Consider an experiment carried out many times. For two event A and B , we use $N(\cdot)$ to represent the number of times that event \cdot occurs. If the event of interest is A and the event B is known or assumed to have occurred, “the conditional probability of A given B ”, or “the probability of A under the condition

B ”, is usually written as $\mathbb{P}(A | B)$ and

$$\mathbb{P}(A | B) = \frac{N(A \cap B)}{N(B)} = \frac{N(A \cap B)/N}{NP(B)/N} = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)}.$$



Example 2.14(Two kids) Assume that a family has two kids. What is the probability that

1. both kids are boys given at least one of them is a boy?
2. both kids are boys given at least one of them is a boy born on Wednesday?

Solution. We use the notation

$$A := \{\text{both kids are boys}\}, C := \{\text{at least one of them is a boy}\},$$

$$D := \{\text{at least one of them is a boy born on Wednesday}\}$$

and letter B and G to represent boy and girl, respectively.

1. The full sample space is $\Omega = \{BB, GB, BG, GG\}$ together with $A = \{BB\}, C = \{BB, BG, GB\}$. Thus the result is $1/3$.
2. The full sample now becomes $\Omega = \{B_i B_j, G_i B_j, B_i G_j, G_i G_j : i, j \in \{1, 2, \dots, 7\}\}$, where i and j denote the birth date (for example, $B_1 B_1$ denote the event that both boys are born on Monday). Now

$$\begin{aligned} |A \cap D| &= |\{\text{two boys at least one is born on Wednesday}\}| \\ &= |\{B_i B_3, B_3 B_j\}| = 13 \end{aligned}$$

and

$$\begin{aligned} |D| &= |\{\text{at least one of the two boys is born on Wednesday}\}| \\ &= |\{B_3 B_j, B_j B_3, B_3 G_j, G_j B_3\}| = 27. \end{aligned}$$

Thus the final is $13/27$.

□

2.5.2 Multiplication formula

Theorem 2.3 (Multiplication formula)

For events A_1, A_2, \dots, A_n satisfying $\mathbb{P}(A_1 \cap A_2 \cap \dots \cap A_{n-1}) > 0$, prove that

$$\mathbb{P}(A_1 \cap A_2 \cap \dots \cap A_n) = \mathbb{P}(A_1) \mathbb{P}(A_2 | A_1) \mathbb{P}(A_3 | A_1 \cap A_2) \cdots \mathbb{P}(A_n | A_1 \cap A_2 \cap \dots \cap A_{n-1}).$$



Proof. Check by direct calculation:

$$\begin{aligned} \text{RHS} &= \mathbb{P}(A_1) \cdot \frac{\mathbb{P}(A_1 \cap A_2)}{\mathbb{P}(A_1)} \cdot \frac{\mathbb{P}(A_1 \cap A_2 \cap A_3)}{\mathbb{P}(A_1 \cap A_2)} \cdots \frac{\mathbb{P}(A_1 \cap A_2 \cap \dots \cap A_n)}{\mathbb{P}(A_1 \cap \dots \cap A_{n-1})} \\ &= \mathbb{P}(A_1 \cap A_2 \cap \dots \cap A_n). \end{aligned}$$

□

Example 2.15(Can model) There are b black balls and r red balls in a can. Every time a man randomly picks one ball from the can and then put it back, together with c balls with the same color and d balls with the other color. We use B_i to denote the event “the i -th picked ball is black” whereas R_j denote the event “the j -th picked ball is red”.

Consider the scenario where two red and one black appear in three consecutive selected balls. Then by Multiplication formula,

$$\begin{aligned}\mathbb{P}(B_1R_2R_3) &= \mathbb{P}(B_1)\mathbb{P}(R_2 | B_1)\mathbb{P}(R_3 | B_1 \cap R_2) \\ &= \frac{b}{b+r} \cdot \frac{r+d}{b+r+c+d} \cdot \frac{r+d+c}{b+r+2c+2d}, \\ \mathbb{P}(R_1B_2R_3) &= \mathbb{P}(R_1)\mathbb{P}(B_2 | R_1)\mathbb{P}(R_3 | R_1 \cap B_2) \\ &= \frac{r}{b+r} \cdot \frac{b+d}{b+r+c+d} \cdot \frac{r+d+c}{b+r+2c+2d}, \\ \mathbb{P}(R_1R_2B_3) &= \mathbb{P}(R_1)\mathbb{P}(R_2 | R_1)\mathbb{P}(B_3 | R_1 \cap R_2) \\ &= \frac{r}{b+r} \cdot \frac{r+c}{b+r+c+d} \cdot \frac{b+2d}{b+r+2c+2d}.\end{aligned}$$

When $c > 0, d = 0$, the can model becomes the **pandemic model**, i.e., every time you pick a red (black) ball, the probability that the next time you still pick a red (black) will increase. Under this restriction,

$$\mathbb{P}(B_1R_2R_3) = \mathbb{P}(R_1B_2R_3) = \mathbb{P}(R_1R_2B_3) = \frac{br(r+c)}{(b+r)(b+r+c)(b+r+2c)}.$$

2.5.3 Total probability formula

Theorem 2.4 (Total probability formula)

For any events A and B such that $0 < \mathbb{P} < 1$,

$$\mathbb{P}(A) = \mathbb{P}(A | B)\mathbb{P}(B) + \mathbb{P}(A | B^c)\mathbb{P}(B^c).$$

More generally, let B_1, B_2, \dots, B_n be a partition of Ω such that $\mathbb{P}(B_i) > 0$ for all i . Then

$$\mathbb{P}(A) = \sum_{i=1}^n \mathbb{P}(A | B_i)\mathbb{P}(B_i).$$



Proof. $A = (A \cap B) \cup (A \cap B^c)$. This is a disjoint union and so

$$\begin{aligned}\mathbb{P}(A) &= \mathbb{P}(A \cap B) + \mathbb{P}(A \cap B^c) \\ &= \mathbb{P}(A | B)\mathbb{P}(B) + \mathbb{P}(A | B^c)\mathbb{P}(B^c).\end{aligned}$$

For the second part, recall the definition of partition,

$$\begin{aligned}A &= (A \cap B_1) \cup (A \cap B_1^c) \\ &= (A \cap B_1) \cup \{A \cap (B_2 \cup B_3 \cup \dots \cup B_n)\} \\ &= (A \cap B_1) \cup (A \cap B_2) \cup \dots \cup (A \cap B_n).\end{aligned}$$

Note that $A \cap B_i (1 \leq i \leq n)$ is disjoint two by two hence we have the desired. \square

Example 2.16(Lottery) Assume that there will be one “win” in n lotteries. What is the probability that the second person wins the lottery?

Solution. Let A_i denote the event that “the i -th person wins the lottery”, $i = 1, 2, \dots, n$. The desired is $\mathbb{P}(A_2)$. Note that A_1 occurs or not will influence A_2 , i.e.,

$$\mathbb{P}(A_2 | A_1) = 0, \quad \mathbb{P}(A_2 | A_1^c) = \frac{1}{n-1}.$$

Moreover, note that

$$\mathbb{P}(A_1) = \frac{1}{n}, \quad \mathbb{P}(A_1^c) = \frac{n-1}{n}$$

and hence by the total probability formula,

$$\mathbb{P}(A_2) = \mathbb{P}(A_1)\mathbb{P}(A_2 | A_1) + \mathbb{P}(A_1^c)\mathbb{P}(A_2 | A_1^c) = \frac{1}{n} \cdot 0 + \frac{n-1}{n} \frac{1}{n-1} = \frac{1}{n}.$$

□

Similarly, one deduces

$$\mathbb{P}(A_3) = \mathbb{P}(A_4) = \dots = \mathbb{P}(A_n) = \frac{1}{n},$$

implying that the lottery is fair.

If $k(\leq n)$ among n lotteries have the reward,

$$\mathbb{P}(A_1) = \mathbb{P}(A_2) = \dots = \mathbb{P}(A_n) = \frac{k}{n}.$$

2.5.4 Bayes formula

Theorem 2.5 (Bayes formula)

Let B_1, B_2, \dots, B_n be a partition of the sample space Ω . Assume that $\mathbb{P}(A) > 0, \mathbb{P}(B_i) > 0, i = 1, 2, \dots, n$, then

$$\mathbb{P}(B_i | A) = \frac{\mathbb{P}(B_i)\mathbb{P}(A | B_i)}{\sum_{j=1}^n \mathbb{P}(B_j)\mathbb{P}(A | B_j)}, \quad i = 1, 2, \dots, n. \quad (2.2)$$

♥

Proof. Note that

$$\begin{aligned} \mathbb{P}(B_i | A) &= \frac{\mathbb{P}(A \cap B_i)}{\mathbb{P}(A)}, \\ \mathbb{P}(A \cap B_i) &= \mathbb{P}(B_i)\mathbb{P}(A | B_i), \\ \mathbb{P}(A) &= \sum_{j=1}^n \mathbb{P}(B_j)\mathbb{P}(A | B_j), \end{aligned}$$

and hence

$$\mathbb{P}(B_i | A) = \frac{\mathbb{P}(B_i)\mathbb{P}(A | B_i)}{\sum_{j=1}^n \mathbb{P}(B_j)\mathbb{P}(A | B_j)}.$$

□

🔗 **Exercise 2.10** Prove that $\mathbb{P}(A | B) = \mathbb{P}(B | A)\mathbb{P}(A)/\mathbb{P}(B)$ whenever $\mathbb{P}(A)\mathbb{P}(B) \neq 0$. Show that, if $\mathbb{P}(A | B) > \mathbb{P}(A)$, then $\mathbb{P}(B | A) > \mathbb{P}(B)$.

Proof. Check by definition. □

🔗 **Exercise 2.11** A man possesses five coins, two of which are double-headed, one is double-tailed, and two are normal. He shuts his eyes, picks a coin at random, and tosses it.

1. What is the probability that the lower face of the coin is a head?
2. He opens his eyes and sees that the coin is showing heads; what is the probability that the lower face is a head?
3. He shuts his eyes again, and tosses the coin again. What is the probability that the lower face is a head?
4. He opens his eyes and sees that the coin is showing heads; what is the probability that the lower face is a head?
5. He discards this coin, picks another at random, and tosses it. What is the probability that it shows heads?

Solution. 1. Let HH be the event the selected coin is double headed, TT be the event that it is double-tailed and N that it is normal. Let H_l denote the event that the lower face of the coin is a head. Then,

$$\begin{aligned}\mathbb{P}(H_l) &= \mathbb{P}(H_l | HH) \mathbb{P}(HH) + \mathbb{P}(H_l | TT) \mathbb{P}(TT) + \mathbb{P}(H_l | N) \mathbb{P}(N) \\ &= 1 \cdot \frac{2}{5} + 0 \cdot \frac{1}{5} + \frac{1}{2} \cdot \frac{2}{5} = \frac{3}{5}.\end{aligned}$$

2. Let H_u be the event that the upper face of the coin is a head. Then,

$$\mathbb{P}(H_l | H_u) = \frac{\mathbb{P}(HH)}{\mathbb{P}(H_u)}.$$

Since $\mathbb{P}(H_u) = \frac{3}{5}$ (the calculation is the same as for $\mathbb{P}(H_l)$), therefore

$$\mathbb{P}(H_l | H_u) = \frac{2/5}{3/5} = \frac{2}{3}.$$

3. Let H_l^2 be the event that the coin's lower in the second toss is a head. Then,

$$\begin{aligned}\mathbb{P}(H_l^2 | H_u) &= \mathbb{P}(H_l^2 | HH, H_u) \mathbb{P}(HH | H_u) + \mathbb{P}(H_l^2 | N, H_u) \mathbb{P}(N | H_u) \\ &= 1 \cdot \mathbb{P}(HH | H_u) + \frac{1}{2} \cdot (1 - \mathbb{P}(HH | H_u)).\end{aligned}$$

Since $HH = H_l \cap H_u$, we can write this probability as

$$\mathbb{P}(H_l^2 | H_u) = 1 \cdot \mathbb{P}(H_l | H_u) + \frac{1}{2} \cdot (1 - \mathbb{P}(H_l | H_u)) = \frac{2}{3} + \frac{1}{2} \cdot \frac{1}{3} = \frac{5}{6}.$$

4.


$$\mathbb{P}(H_l^2 | H_u^2, H_u) = \frac{\mathbb{P}(HH)}{\mathbb{P}(H_u^2 \cap H_u)} = \frac{\mathbb{P}(HH)}{\mathbb{P}(H_u^2 | H_u) \mathbb{P}(H_u)} = \frac{\frac{2}{5}}{\frac{5}{6} \cdot \frac{3}{5}} = \frac{4}{5},$$

where we used results of (1) and (3). (The probabilities $\mathbb{P}(H_u^2 | H_u)$ and $\mathbb{P}(H_u)$ are the same as $\mathbb{P}(H_l^2 | H_u)$ and $\mathbb{P}(H_l)$, respectively).

5. The probability that he discards a double-headed coin is $\frac{4}{5}$ by the previous part, the probability that he discards a normal coin is $\frac{1}{5}$. In the first case we have 1 double-headed coin, 1 double-tailed, and 2 normal coins. In the second case, we have 2 double-headed coins, 1 double-tailed and 1 normal. Hence, by conditioning on the discard we have:

$$\mathbb{P}(H_u^3) = \frac{4}{5} \left(1 \cdot \frac{1}{4} + \frac{1}{2} \cdot \frac{2}{4} \right) + \frac{1}{5} \left(1 \cdot \frac{2}{4} + \frac{1}{2} \cdot \frac{1}{4} \right) = \frac{21}{40}$$

□

 **Exercise 2.12** There are n urns of which the r th contains $r - 1$ red balls and $n - r$ magenta balls. You pick an urn at random and remove two balls without replacement. Find the probability that:

1. the second ball is magenta;
2. the probability that the second ball is magenta given the first is red.

Solution. 1. Note that we have $n - 1$ balls for each urn. First we pick an random urn with equal probability $\frac{1}{n}$. We need to sum up all the possibility that the second removed ball is magenta. For the r th urn, there are $(r - 1)(n - r) + (n - r)(n - r - 1)$ ways of removal (the first is red and the second is magenta or


both are magenta). Hence what we want is

$$\begin{aligned}
 \mathbb{P}(B_2 = M) &= \frac{1}{n} \sum_{r=1}^n \left(\frac{n-r}{n-2} \frac{r-1}{n-1} + \frac{n-r-1}{n-2} \frac{n-r}{n-1} \right) \\
 &= \frac{1}{n(n-1)} \sum_{r=1}^n (n-r) \cdot \frac{r-1+n-r-1}{n-2} \\
 &= \frac{n^2 - \sum_{r=1}^n r}{n^2 - n} \\
 &= \frac{n^2 - \frac{1}{2}n(n+1)}{n^2 - n} \\
 &= \frac{1}{2}
 \end{aligned}$$

2. For this question, we first calculate the probability that both removals are magenta:

$$\begin{aligned}
 \mathbb{P}(B_1 = M, B_2 = M) &= \frac{1}{n} \sum_{r=1}^n \frac{(n-r)(n-r-1)}{(n-1)(n-2)} \\
 &= \frac{1}{n(n-1)(n-2)} \sum_{r=1}^n (n-r)^2 - (n-r) \\
 &= \frac{1}{n(n-1)(n-2)} \left[\frac{(n-1)n(2n-1)}{6} - n^2 + \frac{n(n-1)}{2} \right] \\
 &= \frac{1}{3}.
 \end{aligned}$$


The probability that the first removal is magenta is $\frac{1}{2}$ since there are $n(n-1)$ balls in total and half of them are magenta. Hence the result is $\frac{2}{3}$. □

 **Exercise 2.13** There is one ball with unknown color (either black or white) in a bag. Now put another white ball into the bag and pick one of the two balls randomly, obtaining a white one. What is the probability that the original ball is white?

Solution. We use A to denote the event that the original one is white, B to denote the event that the original one is black and C to denote the event that the picked ball is white. By the condition,

$$\begin{aligned}
 \mathbb{P}(A) &= \mathbb{P}(B) = \frac{1}{2}, \\
 \mathbb{P}(C | A) &= 1, \quad \mathbb{P}(C | B) = \frac{1}{2}.
 \end{aligned}$$

Solving this, we get $\mathbb{P}(A | C) = \frac{2}{3}$. □

 **Exercise 2.14** There are $2n$ vertices among n ropes. Now connecting one vertex with exactly one another vertex, what is the probability that n circles occur?


Solution. We treat the head and tail of a certain rope as different vertices. We first put all $2n$ vertices in a line. Then one connection corresponds to one permutation of $2n$ (for example, 1234 can be regarded as we connect vertices 1 and 2, 3 and 4). Thus the total number of connection method is $(2n)!$.

The only way to form n circles is that we connect each rope's head with their tail. To be more specific, we must place every rope's head and tail in position $2k-1$ and $2k$ for $1 \leq k \leq n$. Since we can switch the head

and tail, there are $2^n \cdot n!$ ways in total. Thus the result is

$$\frac{2^n \cdot n!}{(2n)!} = \frac{(2n)!!}{(2n)!} = \frac{1}{(2n-1)!!},$$

where $(2n)!! = 2n \cdot (2n-1) \cdot \dots \cdot 4 \cdot 2$, $(2n-1)!! = (2n-1) \cdot (2n-3) \cdot \dots \cdot 3 \cdot 1$. \square


 **Exercise 2.15** m students are passing ball to each other starting from Tsubaki. Every time each student (except the one who is passing) has a equal probability to receive the ball. What is the probability that the n -th pass is still finished by Tsubaki?

Solution. We use p_k to denote the probability of the event that the k -th pass is finished by Tsubaki. Then $p_1 = 1$ by condition. Note that if p_{k+1} happens, Tsubaki can not finish the k -th pass and thus

$$p_{k+1} = \frac{1}{m-1}(1-p_k),$$

$$p_{k+1} - \frac{1}{m} = \left(\frac{1}{1-m}\right)\left(p_k - \frac{1}{m}\right).$$

\square

 **Exercise 2.16** There are b black balls and r red balls in a can at first. Every time we pick a ball from it, adding c ($c > 0$) balls with the same color as the picked one and finally put back the picked one. Show that the probability that we pick a black ball at k -th time is $b/(b+r)$, $k = 1, 2, \dots$

Proof. We use $A_i(b, r)$ to denote the event that “there are b black balls and r red balls in the can and the i -th ball taken out is black”. Consider applying induction on i . By condition, $\mathbb{P}(A_1(b, r)) = \frac{b}{b+r}$. Now assume

$\mathbb{P}(A_{k-1}(b, r)) = \frac{b}{b+r}$. By total probability formula, one has

$$\mathbb{P}(A_k(b, r)) = \mathbb{P}(A_1(b, r)) \cdot \mathbb{P}(A_k(b, r) \mid A_1(b, r)) + \mathbb{P}(A_1^c(b, r)) \cdot \mathbb{P}(A_k(b, r) \mid A_1^c(b, r)).$$

Note that


$$\mathbb{P}(A_k(b, r) \mid A_1(b, r)) = \mathbb{P}(A_{k-1}(b+c, r))$$

$$\mathbb{P}(A_k(b, r) \mid A_1^c(b, r)) = \mathbb{P}(A_{k-1}(b, r+c)).$$

To see this, for the first equation, given the event $A_1(b, r)$, c black balls are added. Now $(k-1)$ picks are required in the new can $(b+c, r)$. The second equation is the same. Substituting back and from the induction assumption,

$$\begin{aligned} \mathbb{P}(A_k(b, r)) &= \frac{b}{b+r} \cdot \frac{b+c}{b+c+r} + \frac{r}{b+r} \cdot \frac{b}{b+c+r} \\ &= \frac{b}{b+r}. \end{aligned}$$

\square

 **Exercise 2.17** A bag contains a white balls, b black balls and n red balls. The balls are taken out one by one without returning. Show that the probability that white balls appear earlier than black balls is always $\frac{a}{a+b}$ and has nothing to do with n .

Proof. Let A denote the event that “the first ball taken out is white”, B denote the event that “the first ball taken out is black” and C denote the event that “the first ball taken out is red”. It is easy to see that A, B and C contradicts with each other whereas $A \cup B \cup C = \Omega$. Let E_k denote the event that “when there are k red balls, the white balls appear earlier than the black balls”. Apply induction on k . If $k = 0$, E_0 means the first ball taken out must be white, i.e., $\mathbb{P}(E_0) = a/(a+b)$.

Now assume $\mathbb{P}(E_{k-1}) = \frac{a}{a+b}$. By the total probability formula,

$$\begin{aligned}\mathbb{P}(E_k) &= \mathbb{P}(A)\mathbb{P}(E_k | A) + \mathbb{P}(B)\mathbb{P}(E_k | B) + \mathbb{P}(C)\mathbb{P}(E_k | C) \\ &= \frac{a}{a+b+n} \cdot 1 + \frac{b}{a+b+n} \cdot 0 + \frac{n}{a+b+n} \cdot \mathbb{P}(E_{k-1}) \\ &= \frac{a}{a+b}.\end{aligned}$$

□

Exercise 2.18 Given $\mathbb{P}(A | B) > \mathbb{P}(A | B^c)$, prove that $\mathbb{P}(B | A) > \mathbb{P}(B | A^c)$.

Proof. Since

$$\frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} > \frac{\mathbb{P}(A \cap B^c)}{\mathbb{P}(B^c)} = \frac{\mathbb{P}(A) - \mathbb{P}(B \cap A)}{1 - \mathbb{P}(B)},$$

we know $\mathbb{P}(A \cap B) > \mathbb{P}(A)\mathbb{P}(B)$. Thus

$$\begin{aligned}\mathbb{P}(B | A^c) &= \frac{\mathbb{P}(B \cap A^c)}{\mathbb{P}(A^c)} = \frac{\mathbb{P}(B) - \mathbb{P}(B \cap A)}{1 - \mathbb{P}(A)} \\ &< \frac{\mathbb{P}(B) - \mathbb{P}(B)\mathbb{P}(A)}{1 - \mathbb{P}(A)} = \frac{\mathbb{P}(B)(1 - \mathbb{P}(A))}{1 - \mathbb{P}(A)} \\ &= \mathbb{P}(B)\end{aligned}$$

whereas

$$\mathbb{P}(B | A) = \frac{\mathbb{P}(B \cap A)}{\mathbb{P}(A)} > \frac{\mathbb{P}(A)\mathbb{P}(B)}{\mathbb{P}(A)} = \mathbb{P}(B).$$

□

Exercise 2.19 Let $\mathbb{P}(A) = p$, $\mathbb{P}(B) = 1 - \varepsilon$, prove that:

$$\frac{p - \varepsilon}{1 - \varepsilon} \leq \mathbb{P}(A | B) \leq \frac{p}{1 - \varepsilon}.$$

Proof. On one hand,

$$\mathbb{P}(A | B) = \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} \leq \frac{\mathbb{P}(A)}{\mathbb{P}(B)} = \frac{p}{1 - \varepsilon}.$$

On the other hand,

$$\begin{aligned}\mathbb{P}(A | B) &= \frac{\mathbb{P}(A \cap B)}{\mathbb{P}(B)} = \frac{\mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)}{\mathbb{P}(B)} \\ &\geq \frac{\mathbb{P}(A) + \mathbb{P}(B) - 1}{\mathbb{P}(B)} = \frac{p - \varepsilon}{1 - \varepsilon}.\end{aligned}$$

□

Exercise 2.20 Prove $\mathbb{P}(B^c | A^c) = 1$ provided $\mathbb{P}(A | B) = 1$.

Proof. Apply the formula $\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B) - \mathbb{P}(A \cap B)$.

□

Exercise 2.21 The 'ménages' problem poses the following question. Some consider it to be desirable that men and women alternate when seated at a circular table. If n heterosexual couples are seated randomly according to this rule, show that the probability that nobody sits next to his or her partner is

$$\frac{1}{n!} \sum_{k=0}^n (-1)^k \frac{2n}{2n-k} \binom{2n-k}{k} (n-k)!$$

Proof. Label the seats $1, 2, \dots, 2n$ clockwise. For the sake of definiteness, we dictate that seat 1 be occupied by a woman; this determines the sex of the occupant of every other seat. For $1 \leq k \leq 2n$, let A_k be the event that seats $k, k + 1$ are occupied by one of the couples (we identify seat $2n + 1$ with seat 1). The required probability is

$$\mathbb{P} \left(\bigcap_1^{2n} A_i^c \right) = 1 - \mathbb{P} \left(\bigcup_1^{2n} A_i \right) = 1 - \sum_i \mathbb{P}(A_i) + \sum_{i < j} \mathbb{P}(A_i \cap A_j) - \dots$$

Now, $\mathbb{P}(A_i) = n(n - 1)!^2/n!^2$, since there are n couples who may occupy seats i and $i + 1$, $(n - 1)!$ ways of distributing the remaining $n - 1$ women, and $(n - 1)!$ ways of distributing the remaining $n - 1$ men. Similarly, if $1 \leq i < j \leq 2n$, then

$$\mathbb{P}(A_i \cap A_j) = \begin{cases} n(n - 1) \frac{(n - 2)!^2}{n!^2} & \text{if } |i - j| \neq 1 \\ 0 & \text{if } |i - j| = 1 \end{cases}$$

subject to $\mathbb{P}(A_1 \cap A_{2n}) = 0$. In general,

$$\mathbb{P}(A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}) = \frac{n!}{(n - k)!} \frac{(n - k)!^2}{n!^2} = \frac{(n - k)!}{n!}$$

if $i_1 < i_2 < \dots < i_k$ and $i_{j+1} - i_j \geq 2$ for $1 \leq j < k$, and $2n + i_1 - i_k \geq 2$; otherwise this probability is 0. Hence

$$\mathbb{P} \left(\bigcap_1^{2n} A_i^c \right) = \sum_{k=0}^n (-1)^k \frac{(n - k)!}{n!} S_{k,n}$$

where $S_{k,n}$ is the number of ways of choosing k non-overlapping pairs of adjacent seats. Finally, we calculate $S_{k,n}$. Consider first the number $N_{k,m}$ of ways of picking k non-overlapping pairs of adjacent seats from a line (rather than a circle) of m seats labelled $1, 2, \dots, m$. There is a one-to-one correspondence between the set of such arrangements and the set of $(m - k)$ -vectors containing k 1's and $(m - 2k)$ 0's. To see this, take such an arrangement of seats, and count 0 for an unchosen seat and 1 for a chosen pair of seats; the result is such a vector. Conversely take such a vector, read its elements in order, and construct the arrangement of seats in which each 0 corresponds to an unchosen seat and each 1 corresponds to a chosen pair. It follows that $N_{k,m} = \binom{m - k}{k}$.

Turning to $S_{k,n}$, either the pair $2n, 1$ is chosen or it is not. If it is chosen, we require another $k - 1$ pairs out of a line of $2n - 2$ seats. If it is not chosen, we require k pairs out of a line of $2n$ seats. Therefore

$$S_{k,n} = N_{k-1,2n-2} + N_{k,2n} = \binom{2n - k - 1}{k - 1} + \binom{2n - k}{k} = \binom{2n - k}{k} \frac{2n}{2n - k}$$

□

Exercise 2.22 Each member of a group of n players rolls a die.

1. For any pair of players who throw the same number, the group scores 1 point. Find the mean and variance of the total score of the group.
2. Find the mean and variance of the total score if any pair of players who throw the same number scores that number.

Solution. 1. Let I_{ij} be the indicator function of the event that players i and j throw the same number. Then

$$\mathbb{E}(I_{ij}) = \mathbb{P}(I_{ij} = 1) = \sum_{i=1}^6 \left(\frac{1}{6} \right)^2 = \frac{1}{6}, \quad i \neq j.$$

The total score of the group is $S = \sum_{i < j} I_{ij}$, so

$$\mathbb{E}(S) = \sum_{i < j} \mathbb{E}(I_{ij}) = \frac{1}{6} \binom{n}{2}$$

We claim that the family $\{I_{ij} : i < j\}$ is pairwise independent. The crucial calculation for this is as follows: if $i < j < k$ then

$$\mathbb{E}(I_{ij}I_{jk}) = \mathbb{P}(i, j, \text{ and } k \text{ throw same number}) = \sum_{r=1}^6 \left(\frac{1}{6}\right)^3 = \frac{1}{36} = \mathbb{E}(I_{ij})\mathbb{E}(I_{jk}).$$

Hence

$$\text{var}(S) = \text{var}\left(\sum_{i < j} I_{ij}\right) = \sum_{i < j} \text{var}(I_{ij}) = \binom{n}{2} \text{var}(I_{12})$$

by symmetry. But $\text{var}(I_{12}) = \frac{1}{6} \left(1 - \frac{1}{6}\right)$.

2. Let X_{ij} be the common score of players i and j , so that $X_{ij} = 0$ if their scores are different. This time the total score is $S = \sum_{i < j} X_{ij}$, and

$$\mathbb{E}(S) = \binom{n}{2} \mathbb{E}(X_{12}) = \binom{n}{2} \frac{1}{6} \cdot \frac{7}{2} = \frac{7}{12} \binom{n}{2}.$$

The X_{ij} are not pairwise independent, and you have to slog it out thus:

$$\begin{aligned} \text{var}(S) &= \mathbb{E}\left\{\left(\sum_{i < j} X_{ij}\right)^2\right\} - \mathbb{E}(S)^2 \\ &= \binom{n}{2} \mathbb{E}(X_{12}^2) + \binom{n}{3} \mathbb{E}(X_{12}X_{23}) \\ &\quad + \left\{\binom{n}{2}^2 - \binom{n}{2} - \binom{n}{3}\right\} \mathbb{E}(X_{12})^2 - \left(\frac{7}{12}\right)^2 \binom{n}{2}^2 \\ &= \frac{315}{144} \binom{n}{2} + \frac{35}{432} \binom{n}{3} \end{aligned}$$

□

Chapter 3 Random variables and their distribution

3.1 Random variables; Distribution function

Definition 3.1

Let (E, \mathcal{E}) and (G, \mathcal{G}) be measurable spaces and let μ be a measure on (E, \mathcal{E}) . A function $f : E \rightarrow G$ is said to be **measurable** if

$$f^{-1}(A) := \{x \in E : f(x) \in A\} \in \mathcal{E} \text{ whenever } A \in \mathcal{G}.$$



Note that f^{-1} maps subsets to subsets.

Definition 3.2

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, let (G, \mathcal{G}) be a measure space. Then a measurable function $X : \Omega \rightarrow G$ is called a **random variable**.



Definition 3.3

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, together with the special measure space $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$. Then a measurable function $X : \Omega \rightarrow \mathbb{R}$ is called a **real random variable** and

$$F_X(x) := \mathbb{P}_X((-\infty, x]) = \mathbb{P}(\{X \leq x\})$$

(where $\{X \leq x\} = \{\omega \in \Omega : X(\omega) \leq x\}$) is called the **distribution function of X**.



Proposition 3.1

The map f^{-1} preserves all set operations, i.e. $f^{-1}\left(\bigcup_i A_i\right) = \bigcup_i f^{-1}(A_i)$ and $f^{-1}(A^c) = [f^{-1}(A)]^c$.



Indeed, to verify measurability, we only need to check the preimage on a smaller set:

Lemma 3.1

Let $(\Omega_1, \mathcal{F}_1), (\Omega_2, \mathcal{F}_2)$ be measurable spaces. Suppose there is $\mathcal{E} \subseteq \mathcal{F}_2$ such that $\sigma(\mathcal{E}) = \mathcal{F}_2$. Then $X : \Omega_1 \rightarrow \Omega_2$ is measurable if and only if for any $A \in \mathcal{E}$, $X^{-1}(A) \in \mathcal{F}_1$.



Proof. \implies . Trivial.

\impliedby . Define $G := \{B \in \Omega_2 : X^{-1}(B) \in \mathcal{F}_1\}$. Then follow Proposition 3.1 one can deduce that G is a σ -algebra. By condition, $\mathcal{E} \subseteq G$. Therefore $\mathcal{F}_2 = \sigma(\mathcal{E}) \subseteq G$, implying that X is measurable. \square

Example 3.1 Consider (Ω, \mathcal{F}) and $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$. For any $A \in \mathcal{F}$, the function $X := \mathbb{1}_A$ is measurable.

Indeed, take any $B \in \mathcal{B}(\mathbb{R})$,

$$X^{-1}(B) = \{x \in \Omega : \mathbb{1}_A(x) \in B\} = \begin{cases} \Omega, & \text{if } 0 \in B \text{ and } 1 \in B \\ A^c, & \text{if } 0 \in B \text{ and } 1 \notin B \\ A, & \text{if } 0 \notin B \text{ and } 1 \in B \\ \emptyset, & \text{if } 0 \notin B \text{ and } 1 \notin B \end{cases}.$$

Since \mathcal{F} is a σ -algebra, all possibilities of $X^{-1}(B)$ must lie in \mathcal{F} .

Definition 3.4

Consider measure spaces (Ω, \mathcal{F}) and $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$. A measurable function $f : \Omega \rightarrow \mathbb{R}$ is called a **Borel function**.

**Corollary 3.1**

Let (Ω, \mathcal{F}) be a measurable space. The followings are equivalent:

1. $X : \Omega \rightarrow \mathbb{R}$ is a Borel function;
2. $\{X < a\} \in \mathcal{F}, \forall a \in \mathbb{R}$;
3. $\{X \geq a\} \in \mathcal{F}, \forall a \in \mathbb{R}$;
4. $\{X \leq a\} \in \mathcal{F}, \forall a \in \mathbb{R}$;
5. $\{X > a\} \in \mathcal{F}, \forall a \in \mathbb{R}$.



Proof. We prove $2 \implies 1$ and the remainings are similar. The statement in 2 is equivalent to $\forall a \in \mathbb{R}, X^{-1}((-\infty, a)) \in \mathcal{F}$. By Lemma 3.1, it suffices to show that $\mathcal{E}_2 = \{(-\infty, a) : a \in \mathbb{R}\}$ satisfies $\sigma(\mathcal{E}_2) = \mathcal{B}(\mathbb{R})$. We have know that $\mathcal{B}(\mathbb{R}) = \sigma(\mathcal{E}_0)$ where $\mathcal{E}_0 = \{(a, b] : -\infty \leq a < b < +\infty\}$. Note $\mathcal{E}_0 \subseteq \sigma(\mathcal{E}_2)$ since $(-\infty, b] = \bigcap_{n \geq 1} (-\infty, b + \frac{1}{n})$, $(a, b] = (-\infty, b] \setminus (-\infty, a]$. Conversely, $\mathcal{E}_2 \subseteq \sigma(\mathcal{E}_0)$ since $(-\infty, a) = \bigcup_{n \geq 1} (-\infty, a - \frac{1}{n}]$. □

Lemma 3.2

Let X be a random variable on some probability space. Then the distribution function of X , i.e. F_X , has the following properties:

1. $F_X : \mathbb{R} \rightarrow [0, 1]$, and F_X is monotonically increasing;
2. $\lim_{x \rightarrow +\infty} F_X(x) = 1, \lim_{x \rightarrow -\infty} F_X(x) = 0$;
3. F_X is right-continuous (i.e., $F_X(y) \downarrow F_X(x)$ as $y \downarrow x$).



Proof. 1. Note that $\{X \leq x\} \subseteq \{X \leq y\}$ and hence $F_X(x) \leq F_X(y)$ by the monotonicity of probability.

2. First we show $\lim_{n \rightarrow -\infty} F(n) = 0$. By “continuity from above”,

$$\begin{aligned} \lim_{n \rightarrow -\infty} F(n) &= \lim_{n \rightarrow -\infty} \mathbb{P}(\{X \leq n\}) = \mathbb{P}(\lim_{n \rightarrow -\infty} \{X \leq n\}) \\ &= \mathbb{P}\left(\bigcap_{n \in \mathbb{Z}} \{X \leq n\}\right) = \mathbb{P}(\emptyset) \\ &= 0. \end{aligned}$$

Symmetrically, by continuity from below,

$$\begin{aligned} \lim_{n \rightarrow +\infty} F(n) &= \lim_{n \rightarrow +\infty} \mathbb{P}(\{X \geq n\}) = \mathbb{P}(\lim_{n \rightarrow +\infty} \{X \geq n\}) \\ &= \mathbb{P}\left(\bigcup_{n \in \mathbb{Z}} \{X \geq n\}\right) = \mathbb{P}(\mathbb{R}) \\ &= 1. \end{aligned}$$

3. First we show $\lim_{n \rightarrow \infty} F(x + \frac{1}{n}) = F(x)$. By continuity from above,

$$\lim_{n \rightarrow \infty} F(x + \frac{1}{n}) = \lim_{n \rightarrow \infty} \mathbb{P}(\{X \leq x + \frac{1}{n}\}) = \mathbb{P}(\lim_{n \rightarrow \infty} \{X \leq x + \frac{1}{n}\})$$

$$\begin{aligned}
 &= \mathbb{P} \left(\bigcap_{n \geq 1} \left\{ X \leq x + \frac{1}{n} \right\} \right) = \mathbb{P}(\{X \leq x\}) \\
 &= F(x).
 \end{aligned}$$

The by monotonicity $F_X(x+h) \downarrow F_X(x)$ as $h \downarrow 0$. □

Indeed, we can also prove that functions satisfying the three properties in Lemma 3.2 are distribution functions of certain random variables. Hence **these three properties are necessary and sufficient judgements to determine whether a function is a distribution function (of a certain random variable) or not:**

Theorem 3.1

If F satisfies 1, 2, and 3 in Lemma 3.2, then it is the distribution function of some random variable. ♥

Proof. Let $\Omega = (0, 1)$, \mathcal{F} = the Borel sets, and P = probability (or, Lebesgue) measure. If $\omega \in (0, 1)$, let

$$X(\omega) = \sup\{y : F(y) < \omega\}$$

Once we show that

$$\{\omega : X(\omega) \leq x\} = \{\omega : \omega \leq F(x)\} \quad (\star)$$

the desired result follows immediately since $P(\omega : \omega \leq F(x)) = F(x)$. (Recall P is probability measure.) To check (\star) , we observe that if $\omega \leq F(x)$ then $X(\omega) \leq x$, since $x \notin \{y : F(y) < \omega\}$. On the other hand if $\omega > F(x)$, then since F is right continuous, there is an $\epsilon > 0$ so that $F(x+\epsilon) < \omega$ and $X(\omega) \geq x+\epsilon > x$. □

Lemma 3.3

Let X be a random variable and F be its distribution function. Then for $x, y \in \mathbb{R}$,

1. $\mathbb{P}(x < X \leq y) = F(y) - F(x)$;
 2. $\mathbb{P}(X = x) = F(x) - \lim_{y \rightarrow x^-} F(y)$.
- ♥

Proof. Statement 1 is trivial. To see 2, let $B_n = \{x - \frac{1}{n} < X \leq x\}$. By 1, $\mathbb{P}(B_n) = F(x) - F(x - \frac{1}{n})$. Now send $n \rightarrow \infty$, one gets $\lim_{n \rightarrow \infty} B_n = \bigcap_{n \geq 1} B_n = \{x\}$. By continuity of measure,

$$\mathbb{P}(X = x) = \lim_{n \rightarrow \infty} \mathbb{P}(B_n) = F(x) - \lim_{n \rightarrow \infty} F(x - \frac{1}{n}) = F(x) - \lim_{y \rightarrow x^-} F(y).$$
□

🔥 **Exercise 3.1** Let $(\Omega_1, \mathcal{F}_1)$ and $(\Omega_2, \mathcal{F}_2)$ be measurable spaces with $f : \Omega_1 \rightarrow \Omega_2$.

1. Show that $\mathcal{F} := \{f^{-1}(A) : A \in \mathcal{F}_2\}$ is a σ -algebra on Ω_1 .
2. Show that if \mathcal{G} is a σ -algebra on Ω_1 such that f is \mathcal{G} -measurable, then $\mathcal{G} \supseteq \mathcal{F}$.

Proof. 1. Trivial.

2. Take $B \in \mathcal{F}$ then $B = f^{-1}(A)$ for some $A \in \mathcal{F}_2$. Since f is \mathcal{G} -measurable, $B = f^{-1}(A) \in \mathcal{G}$, and thus $\mathcal{G} \supseteq \mathcal{F}$. □

🔥 **Exercise 3.2** Let $(\Omega_1, \mathcal{F}_1)$ and $(\Omega_2, \mathcal{F}_2)$ be measurable spaces with $f : \Omega_1 \rightarrow \Omega_2$.

1. Show that $\tilde{\mathcal{F}} := \{A \subseteq \Omega_2 : f^{-1}(A) \in \mathcal{F}_1\}$ is a σ -algebra on Ω_2 .

2. Show that if \mathcal{G} is a σ -algebra on Ω_2 such that f is \mathcal{G} -measurable, then $\mathcal{G} \subseteq \tilde{\mathcal{F}}$.

Proof. 1. Trivial.

2. Take $B \in \mathcal{G}$ then since f is \mathcal{G} -measurable, $f^{-1}(B) \in \mathcal{F}_1$, i.e., $B \in \tilde{\mathcal{F}}$ and thus $\mathcal{G} \subseteq \tilde{\mathcal{F}}$. □

3.2 Properties of Borel functions

For random variable X and Y , we use the shorthand $\{X < Y\} := \{\omega : \omega \in \Omega : X(\omega) < Y(\omega)\}$.

Lemma 3.4

Let (Ω, \mathcal{F}) be a measurable space and X, Y be Borel functions. Then

1. $\{X < Y\}, \{X \geq Y\}, \{X = Y\}, \{X \neq Y\} \in \mathcal{F}$;
2. $cX : c \in \mathbb{R}, X + Y, X - Y$ and XY are Borel functions. ♥

Proof. Recall Corollary 3.1, X is a Borel functions if and only if $\{X < a\} \in \mathcal{F}$ for any $a \in \mathbb{R}$.

1. Using \mathbb{Q} is dense in \mathbb{R} , one gets $\{X < Y\} = \bigcup_{q \in \mathbb{Q}} \{X < q\} \cap \{q < Y\} \in \mathcal{F}$. For the remaining terms, note that $\{X \leq Y\} = \{X < Y\}^c, \{X = Y\} = \{X \leq Y\} \cap \{X \geq Y\}$ and that $\{X \neq Y\} = \{X = Y\}^c$.
2. For $t \in \mathbb{R}$, if $c > 0$,

$$\{X > t\} = \{X > c^{-1}t\}.$$

If $c < 0$,

$$\{X < t\} = \{X < c^{-1}t\}$$

and thus cX is Borel. If $c = 0$, it is trivial.

For $t \in \mathbb{R}$, note that

$$\{X + Y > t\} = \bigcup_{i=1}^{\infty} (\{X > r_i\} \cap \{Y > t - r_i\})$$

where r_i are all the rational numbers. Thus $X + Y$ is Borel (same as $X - Y$). Also note that X^2 is Borel since $\{X^2 > a\} = \{X > \sqrt{a}\} \cup \{X < -\sqrt{a}\}$. Combine all these and by the transformation $XY = \frac{1}{4}((X + Y)^2 - (X - Y)^2)$, one deduces that XY is Borel. □

Lemma 3.5

Let $\{X_n\}$ be a sequence of Borel function. Then $\sup_n X_n, \inf_n X_n, \limsup_{n \rightarrow \infty} X_n, \liminf_{n \rightarrow \infty} X_n$ are Borel functions. ♥

Proof. We claim that

$$\{\sup_n X_n \leq a\} = \bigcap_{n \geq 1} \{X_n \leq a\}$$

and thus $\sup_n X_n$ is Borel. The infimum $\inf_n X_n$ is Borel since $\inf_n X_n = -\sup_n(-X_n)$. The remaining terms are Borel since

$$\begin{aligned} \limsup_{n \rightarrow \infty} X_n &= \inf_{m \geq 1} \sup_{n \geq m} X_n, \\ \liminf_{n \rightarrow \infty} X_n &= \sup_{m \geq 1} \inf_{n \geq m} X_n. \end{aligned}$$

□

Corollary 3.2

If $\lim_{n \rightarrow \infty} X_n$ exists, then $\lim_{n \rightarrow \infty} X_n$ is Borel.

♥

Lemma 3.6

Let $(\Omega_1, \mathcal{F}_1), (\Omega_2, \mathcal{F}_2), (\Omega_3, \mathcal{F}_3)$ be three measurable spaces. $X : \Omega_1 \rightarrow \Omega_2$ and $Y : \Omega_2 \rightarrow \Omega_3$ are measurable functions. Then $Y \circ X : \Omega_1 \rightarrow \Omega_3$ is also measurable.

♥

Proof. For any $B \in \mathcal{F}_3$, note that $(Y \circ X)^{-1}(B) = X^{-1} \circ Y^{-1}(B) = X^{-1}(Y^{-1}(B)) \in \mathcal{F}_1$. □

3.3 σ -algebra generated by a random variable

Definition 3.5

Let X be a random variable. Then

$$\sigma(X) := \{X^{-1}(B) : B \in \mathcal{B}(\mathbb{R})\}$$

is called **the σ -algebra generated by X** .

♣

Similarly, given measurable X , $\sigma(X)$ is the smallest σ -algebra containing X .

Definition 3.6

Given $\{X_i\}_{i \in I}$ to be a family of random variables,

$$\sigma(X_i, i \in I) := \sigma\left(\bigcup_{i \in I} \sigma(X_i)\right)$$

is called **the σ -algebra generated by $\{X_i\}_{i \in I}$** .

♣

Example 3.2 Let Ω, \mathcal{F} be measurable space and $A_1, A_2, \dots, A_n \in \mathcal{F}$ are two-by-two disjoint. Define

$$X := b_1 \mathbb{1}_{A_1} + b_2 \mathbb{1}_{A_2} + \dots + b_n \mathbb{1}_{A_n}$$

with distinct coefficients b_1, \dots, b_n . Show that $\sigma(X) = \sigma(\{A_1, A_2, \dots, A_n\})$.

Before starting the proof, we first verify a lemma:

Lemma 3.7

$$\sigma(X) = \sigma(\{X < a : a \in \mathbb{R}\}).$$

♥

Proof of the lemma. Recall that $\mathcal{B}(\mathbb{R}) = \sigma(\mathcal{E})$ where $\mathcal{E} = \{[a, b) : -\infty < a < b \leq +\infty\}$. Now for any $S \in \sigma(X)$, there is $[a, b)$ such that $S = X^{-1}([a, b)) = X^{-1}((-\infty, a)^c \cap (-\infty, b)) = \{X < a\}^c \cap \{X < b\} \in \text{RHS}$, implying $\text{LHS} \subseteq \text{RHS}$.

Conversely, for a fixed a , $\{X < a\} = X^{-1}((-\infty, a)) \in \sigma(X)$. Hence by the “smallest”, $\text{RHS} \subseteq \text{LHS}$. □

Proof of Example 3.2. For any i , note that $A_i = X^{-1}(\{b_i\}) \in \sigma(X)$ and thus $\text{RHS} \subseteq \text{LHS}$.

Conversely, for any $a \in \mathbb{R}$, $\{X \leq a\} = \text{disjoint union of } \{A_i\}_{i=1}^n \text{ such that } b_i \leq a = \left(\bigcup_{i=1}^n A_i\right)^c \in \sigma(\{A_1, A_2, \dots, A_n\})$.

Hence by the lemma, $\sigma(X) = \sigma(\{X \leq a\}) \subseteq \sigma(\{A_1, A_2, \dots, A_n\})$. □

3.4 Independence

Definition 3.7

Let I be a countable set. We say that the events $\{A_i \in \mathcal{F} : i \in I\}$ are independent if, for all finite subsets $J \in I$,

$$\mathbb{P}\left(\bigcap_{i \in J} A_i\right) = \prod_{i \in J} \mathbb{P}(A_i).$$



Definition 3.8

Intuitively, two random variables X and Y are said to be **independent** if for any $A, B \in \mathcal{B}(\mathbb{R})$,

$$P(X \in A, Y \in B) = P(X \in A)P(Y \in B).$$



Definition 3.9

Let (Ω, \mathcal{F}, P) be a probability space.

1. A sequence of $G_1, G_2, \dots, G_n \subseteq \mathcal{F}$ is called **independent** if

$$P(A_1 \cap A_2 \cap \dots \cap A_n) = P(A_1)P(A_2) \dots P(A_n),$$

for all $A_i \in G_i, i = 1, 2, \dots, n$.

2. A sequence of random variable X_1, X_2, \dots, X_n are **independent** if $\sigma(X_1), \dots, \sigma(X_n)$ are independent, where $\sigma(X_i) = \{X_i^{-1}(B) : B \in \mathcal{B}(\mathbb{R})\}$.



Remark

- Statement 2 $\iff \forall B_1, B_2, \dots, B_n \in \mathcal{B}(\mathbb{R})$,

$$P(X_1 \in B_1, \dots, X_n \in B_n) = \prod_{i=1}^n P(X_i \in B_i) = \prod_{i=1}^n P(X_i^{-1}(B_i)).$$

- Events A_1, \dots, A_n are independent $\iff \mathbb{1}_{A_1}, \dots, \mathbb{1}_{A_n}$ are independent random variables.

Definition 3.10

The random variables $X_n, n \in \mathbb{N}$, are called **independent identically distributed (i.i.d.)** if they are independent and, moreover, all of them have the same distribution (i.e. the distribution functions μ_{X_n} are equal, for all $n \in \mathbb{N}$).



Theorem 3.2 (Dynkin $\pi - \lambda$)

If \mathcal{C} is a π -system and \mathcal{L} is a λ -system such that $\mathcal{C} \subseteq \mathcal{L}$, then $\sigma(\mathcal{C}) \subseteq \mathcal{L}$



Theorem 3.3

Suppose $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_n$ are independent π -systems. Then $\sigma(\mathcal{A}_1), \dots, \sigma(\mathcal{A}_n)$ are independent.



Proof. We prove if $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_n$ are independent, then $\sigma(\mathcal{A}_1), \mathcal{A}_2, \dots, \mathcal{A}_n$ are independent and iterate.

Take $A_2 \in \mathcal{A}_2, \dots, A_n \in \mathcal{A}_n$. Set $B := A_2 \cap A_3 \dots \cap A_n$. Define $\mathcal{L} = \{A \in \Omega : P(A \cap B) = P(A)P(B)\}$. If $A \in \mathcal{A}_1$, then $P(A \cap B) = P(A)P(B)$, implying that $\mathcal{L} \supseteq \mathcal{A}_1$. We claim that \mathcal{L} is a λ -system. Clearly $\Omega \in \mathcal{L}$ since $P(\Omega \cap B) = P(\Omega)P(B)$. If $A_1, A_2 \in \mathcal{L} : A_1 \subseteq A_2$, then

$$\begin{aligned} P((A_2 \setminus A_1) \cap B) &= P(A_2 \cap B) - P(A_1 \cap B) \\ &= P(A_2)P(B) - P(A_1)P(B) \end{aligned}$$

$$= P(A_2 \setminus A_1)P(B).$$

If $A_k \in \mathcal{L}$ and $A_k \uparrow A$, then

$$\begin{aligned} P(A \cap B) &= \lim_{k \rightarrow \infty} P(A_k \cap B) \\ &= \lim_{k \rightarrow \infty} P(A_k)P(B) \\ &= P(A)P(B). \end{aligned}$$

By Dynkin $\pi - \lambda$ Theorem, $\mathcal{L} \subseteq \sigma(\mathcal{A}_1)$, implying for any $A \in \sigma(\mathcal{A}_1)$,

$$\begin{aligned} P(A \cap (A_2 \cap A_3 \cap \cdots \cap A_n)) &= P(A)P(A_2 \cap A_3 \cap \cdots \cap A_n) \\ &= \prod_{i=1}^n P(A_i). \end{aligned}$$

□

Corollary 3.3

X_1, X_2, \dots, X_n are independent random variables if and only if for any $x_1, x_2, \dots, x_n \in \mathbb{R}$,

$$P[X_1 \leq x_1, \dots, X_n \leq x_n] = \prod_{i=1}^n P(X_i \leq x_i). \quad (3.1)$$

♥

Proof. Let $\mathcal{A}_i = \{\{X_i \leq x\} : x \in \mathbb{R}\}$, then each \mathcal{A}_i is a π -system since $\{X_i \leq x\} \cap \{X_i \leq y\} = \{X_i \leq x \wedge y\}$. Furthermore, $\sigma(\mathcal{A}_i) = \sigma(X_i)$ by Lemma 3.7. Now (3.1) implies $\mathcal{A}_1, \dots, \mathcal{A}_n$ are independent and thus by Theorem 3.3, $\sigma(X_1), \dots, \sigma(X_n)$ are independent. □

🔗 **Exercise 3.3** Let $\Omega := [0, 1] \times [0, 1]$, $\mathcal{F} = \mathcal{B}(\mathbb{R}^2) \cap [0, 1]^2$ and m denote the Lebesgue measure. Define $\mathcal{A}_1 = \{[0, 1] \times A : A \in \mathcal{B}(\mathbb{R}) \cap [0, 1]\}$ and $\mathcal{A}_2 = \{A \times [0, 1] : A \in \mathcal{B}(\mathbb{R}) \cap [0, 1]\}$. Show that \mathcal{A}_1 and \mathcal{A}_2 are σ -algebra and they are independent with respect to m .

Proof.

□

Theorem 3.4 (Uniqueness of extension)

Let \mathcal{G} be a π -system. Let μ_1, μ_2 be two σ -finite measure on $(\Omega, \sigma(\mathcal{G}))$ such that $\mu_1(\Omega) = \mu_2(\Omega)$ and $\mu_1 = \mu_2$ on \mathcal{G} . Then $\mu_1 = \mu_2$ on $\sigma(\mathcal{G})$.

♥

Proof. Let $\mathcal{D} := \{A \in \sigma(\mathcal{G}) : \mu_1(A) = \mu_2(A)\}$. Then we have $\mathcal{D} \supseteq \mathcal{G}$. We claim that \mathcal{D} is a λ -system. Indeed,

- $\Omega \in \mathcal{D}$ since $\mu_1(\Omega) = \mu_2(\Omega)$;
- If $A, B \in \mathcal{D} : A \subset B$, note that

$$\mu_2(B \setminus A) = \mu_2(B) - \mu_2(A) = \mu_1(B) - \mu_1(A) = \mu_1(B \setminus A)$$

- If $A, A_n \in \mathcal{D} : A_n \uparrow A$, note thta

$$\mu_2(A) = \lim_{n \rightarrow \infty} \mu_2(A_n) = \lim_{n \rightarrow \infty} \mu_1(A_n) = \mu_1(A).$$

Hence by Dynkin's Theorem, $\mathcal{D} \supseteq \sigma(\mathcal{G})$ and thus $\mu_1 = \mu_2$ on $\sigma(\mathcal{G})$. □

Constructing independent random variables

Let F_1, F_2, \dots, F_n be distribution functions. How to construct independent variables X_1, X_2, \dots, X_n such that $P(X_i \leq x) = F_i(x)$?

Let $\Omega = \mathbb{R}^n$ and $\mathcal{F} = \mathcal{B}(\mathbb{R}^n)$ and X_i be the projection mapping:

$$\begin{aligned} X_i : \mathbb{R}^n &\longrightarrow \mathbb{R} \\ \omega &\longmapsto \omega_i \end{aligned}$$

Let P_n be the measure on $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$ such that $P_n([a_1, b_1] \times [a_2, b_2] \times \dots \times [a_n, b_n]) = \prod_{i=1}^n (F_i(b_i) - F_i(a_i))$.

Since $\{[a_1, b_1] \times \dots \times [a_n, b_n]\}$ is a π -system that produces $\mathcal{B}(\mathbb{R}^n)$, by extension theorem, P_n extends uniquely to a measure on $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n))$. Indeed, $P_n = m_{F_1} \otimes m_{F_2} \otimes \dots \otimes m_{F_n}$ where m_{F_1}, \dots, m_{F_n} are one-dimensional Lebesgue Stieljes measures.

Constructing infinite random variables

We try a similar construction: $\Omega := \mathbb{R}^{\mathbb{N}} = \{(\omega_1, \omega_2, \dots) : \omega_i \in \mathbb{R}\}$. Our goal is to define a σ -algebra and a probability measure based on approximation on finite dimension. Define the **cylinder set** \mathcal{A} as

$$\mathcal{A} := \bigcup_{n=1}^{\infty} \{B_1 \times B_2 \times \dots \times B_n \times \mathbb{R} \times \mathbb{R} \times \dots : B_1, B_2, \dots, B_n \in \mathcal{B}(\mathbb{R})\}.$$

Note that \mathcal{A} is an algebra but not a σ -algebra: Let $A_n = \mathbb{R} \times \dots \times [0, 1] \times \mathbb{R} \times \dots$. Then $\bigcap_{n \geq 1} A_n = [0, 1]^{\mathbb{N}} \notin \mathcal{A}$.

It is natural to define a content (but countably-additive) on \mathcal{A} : for any $A \in \mathcal{A}$, there is $n \in \mathbb{N}$ such that $A = B_1 \times \dots \times B_n \times \mathbb{R} \times \dots \times \mathbb{R}$. Set

$$P(B_1 \times B_2 \times \dots \times B_n \times \mathbb{R} \times \mathbb{R} \times \dots) = P_n(B_1 \times B_2 \times \dots \times B_n)$$

Can we extend P to $\sigma(\mathcal{A})$?

Theorem 3.5 (Kolmogorov extension)

Suppose the probability measure $(\mathbb{R}^n, \mathcal{B}(\mathbb{R}^n), P_n)_{n \geq 1}$ are consistent:

$$P_{n+1}((a_1, b_1] \times \dots \times (a_n, b_n] \times \mathbb{R}) = P_n((a_1, b_1] \times \dots \times (a_n, b_n]), \quad n \geq 1$$

Then there is a unique probability measure P on $(\mathbb{R}^{\mathbb{N}}, \sigma(\mathcal{A}))$ such that

$$P((a_1, b_1] \times \dots \times (a_n, b_n] \times \mathbb{R} \times \dots) = P_n((a_1, b_1] \times \dots \times (a_n, b_n]).$$



Example 3.3 If $\{X_i\}_{i \in \mathbb{N}}$ is a random walk or a Markov chain, the joint distribution

$$(X_{t_1}, \dots, X_{t_n}), \quad n \geq 1$$

uniquely determines the law of $\{X_i\}_{i \in \mathbb{N}}$.

Chapter 4 Discrete and continuous random variables

4.1 Definitions

Definition 4.1

A random variable X is said to be **discrete** if it takes value in countable set $\{x_1, x_2, \dots\}$, where $\{x_1, x_2, \dots\}$ are called the **atoms** of X . Its **probability mass function** $f(x)$ is defined by

$$f(x) := \mathbb{P}(X = x). \quad (4.1)$$

Definition 4.2

A random variable X is **(absolutely) continuous** if

$$F_X(x) = \int_{-\infty}^x f(u) du \quad (4.2)$$

for some integrable function $f : \mathbb{R} \rightarrow [0, +\infty)$. Then f is called the **probability density function**.

Remark There is random variable such that its distribution function is not absolutely continuous:

Example 4.1 Singular variable X whose distribution F_X is the Cantor function.

The probability density function has two properties:

1. $f(x) \geq 0$;
2. $\int_{-\infty}^{\infty} f(x) dx = 1$.

The distribution and mass function are related by

$$F_X(x) = \mathbb{P}(X \leq x) = \sum_{x_i, x_i \leq x} f(x_i), \quad f(x) = F'(x) = \lim_{y \uparrow x} F(y)$$

Based on (4.2), the relation of probability density function and distribution function is also given by

$$F'(x) = f(x). \quad (4.3)$$

For a discrete random variable, its distribution is always a right-continuous step function. However, for a continuous random variable, its distribution function is always a continuous one since

$$F(x + \Delta x) - F(x) = \int_x^{x+\Delta x} f(x) dx \rightarrow 0 \quad (\Delta x \rightarrow 0).$$

Note that for an absolutely continuous random variable, $P(X = a) = \int_a^a f(x) dx = 0$ for any $a \in \mathbb{R}$ and thus

$$P(a \leq X \leq b) = P(a < X \leq b) = P(a \leq X < b) = P(a < X < b).$$

4.2 Expectation and variance

Definition 4.3

The **mean/expectation/expected value** of a **discrete** random variable X with probability mass function f is

$$\mathbb{E}(X) := \sum_{x: f(x) > 0} x f(x) = \sum_{x: f(x) > 0} x \mathbb{P}(X = x). \quad (4.4)$$

Example 4.2 Let X denote the number of heads occurring in two coin flips. Then according to (4.4),

$$\begin{aligned}\mathbb{E}(X) &= \sum x\mathbb{P}(X = x) \\ &= 0 \cdot \mathbb{P}(X = 0) + 1 \cdot \mathbb{P}(X = 1) + 2 \cdot \mathbb{P}(X = 2) \\ &= 0 + 1 \cdot \frac{1}{2} + 2 \cdot \frac{1}{4} \\ &= 1.\end{aligned}$$

Lemma 4.1 (Change of variable)

Let $g : \mathbb{R} \rightarrow \mathbb{R}$ and X be a random variable with probability mass function f . Then

$$\mathbb{E}[g(X)] = \sum_{x:f(x)>0} g(x)f(x).$$



Proof. Note that for a fixed y ,

$$\mathbb{P}(g(X) = y) = \mathbb{P}\{\omega : g(X(\omega)) = y\} = \sum_{x:g(x)=y} \mathbb{P}(X = x)$$

and thus

$$\begin{aligned}\mathbb{E}(g(X)) &= \sum_y y\mathbb{P}(g(X) = y) = \sum_y \sum_{x:g(x)=y} y\mathbb{P}(X = x) \\ &= \sum_x g(x)\mathbb{P}(X = x) = \sum_x g(x)f(x).\end{aligned}$$

□

Lemma 4.2

Let X be a random variable taking values in \mathbb{N} , then

1. $\mathbb{E}(X) = \sum_{n \in \mathbb{N}} \mathbb{P}(X \geq n)$.
2. $\sum_{k=0}^{\infty} k\mathbb{P}(X > k) = \frac{1}{2}[\mathbb{E}(X^2) - \mathbb{E}(X)]$.



Proof. 1.

$$\begin{aligned}\mathbb{E}(X) &= 1 \cdot \mathbb{P}(X = 1) + 2 \cdot \mathbb{P}(X = 2) + 3 \cdot \mathbb{P}(X = 3) + \dots \\ &= (\mathbb{P}(X = 1) + \mathbb{P}(X = 2) + \dots) + (\mathbb{P}(X = 2) + \mathbb{P}(X = 3) + \dots) + \dots \\ &= \mathbb{P}(X \geq 1) + \mathbb{P}(X \geq 2) + \dots \\ &= \sum_{n \in \mathbb{N}} \mathbb{P}(X \geq n).\end{aligned}$$

2.

$$\begin{aligned}\sum_{k=0}^{\infty} k\mathbb{P}(X > k) &= \sum_{k=0}^{\infty} k \sum_{i=k+1}^{\infty} \mathbb{P}(X = i) = \sum_{i=1}^{\infty} \sum_{k=0}^{i-1} k\mathbb{P}(X = i) \\ &= \sum_{i=1}^{\infty} \mathbb{P}(X = i) \frac{(i-1)i}{2} = \frac{1}{2} \sum_{i=1}^{\infty} i^2 \mathbb{P}(X = i) - \frac{1}{2} \sum_{i=1}^{\infty} i\mathbb{P}(X = i) \\ &= \frac{1}{2} \mathbb{E}(X^2) - \frac{1}{2} \mathbb{E}(X).\end{aligned}$$

□

Example 4.3 Let X_i be i.i.d random variable denoting the offer you receive and define T as

$$T := \text{the first time you see an offer better than } X_1.$$

Compute $\mathbb{E}(T)$.

Solution. By Lemma 4.2,

$$\begin{aligned} \mathbb{E}(T) &= \sum_{n \geq 1} \mathbb{P}(T \geq n) = \sum_{n \geq 1} \mathbb{P}(X_1 = \max\{X_1, X_2, \dots, X_n\}) \\ &= \sum_{n \geq 1} \frac{1}{n} = +\infty. \end{aligned}$$

□

Definition 4.4

Let X be a continuous random variable with probability density function $p(x)$. If

$$\int_{-\infty}^{\infty} |x|p(x)dx < \infty,$$

the value

$$\mathbb{E}(X) = \int_{-\infty}^{\infty} xp(x)dx < \infty, \quad (4.5)$$

is called the *mean/expectation/expected value* of X .



Similarly, one has

Lemma 4.3 (Change of variable)

Let $g : \mathbb{R} \rightarrow \mathbb{R}$ and X be a random variable with probability mass function f . Then

$$\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x)p(x)dx.$$



Lemma 4.4

For a non-negative continuous random variable X , one has

$$E(X) = \int_0^{\infty} P(X \geq x)dx.$$



Proof. It suffices to prove that

$$E(X) = \int_0^{\infty} [1 - F(x)]dx.$$

Note that

$$\int_0^{\infty} xf(x)dx = \int_0^{\infty} \left(\int_0^x dy \right) f(x)dx = \int_0^{\infty} \left(\int_y^{\infty} f(x)dx \right) dy = \int_0^{\infty} [1 - F(y)]dy.$$

□

Exercise 4.1 Let $X \geq 0$ be a random variable. Show that

$$E(X^n) = \int_0^{\infty} nx^{n-1}P(X > x)dx.$$

Proof. Use the conclusion above, we see

$$E(X^n) = \int_0^{\infty} P(X^n > y)dy.$$

Now set $y = x^n$, we get

$$E(X^n) = \int_0^{\infty} P(X^n > x^n) n x^{n-1} dx = \int_0^{\infty} P(X > x) n x^{n-1} dx.$$

□

Definition 4.5

The k -th moment of a random variable X is defined as $m_k := \mathbb{E}(X^k)$ whereas the **the k -th central moment** of X is defined as $\sigma_k := \mathbb{E}[(X - \mathbb{E}(X))^k]$.



Definition 4.6

The second central moment, i.e., $\mathbb{E}[(X - \mathbb{E}(X))^2]$ is defined to be the **variance** of X , denoted by σ^2 or $\text{var}(X)$.



Remark Note that

$$\begin{aligned} \text{var}(X) &= \mathbb{E}[(X - \mathbb{E}(X))^2] = \mathbb{E}[X^2 - 2X \cdot \mathbb{E}(X) + \mathbb{E}(X)^2] \\ &= \mathbb{E}(X^2) - 2\mathbb{E}(X)^2 + \mathbb{E}(X)^2 \\ &= \mathbb{E}(X^2) - \mathbb{E}(X)^2. \end{aligned}$$

Lemma 4.5

Expectation has the following properties:

1. For any $a, b \in \mathbb{R}$, $\mathbb{E}(aX + bY) = a\mathbb{E}(X) + b\mathbb{E}(Y)$;
2. $\mathbb{E}(XY) = \mathbb{E}(X)\mathbb{E}(Y)$ given X and Y independent;
3. $\text{var}(X + Y) = \text{var}(X) + \text{var}(Y)$ given X and Y independent.



Proof. 1. Trivial.

2. Note that

$$\begin{aligned} \mathbb{E}(XY) &= \sum x_i y_j \mathbb{P}(X = x_i, Y = y_j) \\ &= \sum x_i \mathbb{P}(X = x_i) \sum y_j \mathbb{P}(Y = y_j) \\ &= \mathbb{E}(X)\mathbb{E}(Y). \end{aligned}$$


3.

$$\begin{aligned} \text{var}(X) + \text{var}(Y) &= \mathbb{E}((X + Y - \mathbb{E}(X + Y))^2) \\ &= \mathbb{E}((X - \mathbb{E}(X))^2) + \mathbb{E}((Y - \mathbb{E}(Y))^2) + 2\mathbb{E}((X - \mathbb{E}(X))(Y - \mathbb{E}(Y))) \\ &= \text{var}(X) + \text{var}(Y). \end{aligned}$$

The last equality holds since by 2, $\mathbb{E}(XY) = \mathbb{E}(X)\mathbb{E}(Y)$ and

$$\begin{aligned} \mathbb{E}((X - \mathbb{E}(X))(Y - \mathbb{E}(Y))) &= \mathbb{E}(XY - X\mathbb{E}(Y) - Y\mathbb{E}(X) + \mathbb{E}(X)\mathbb{E}(Y)) \\ &= \mathbb{E}(XY) - 2\mathbb{E}(X)\mathbb{E}(Y) + \mathbb{E}(X)\mathbb{E}(Y) \\ &= 0. \end{aligned}$$

□

 **Exercise 4.2** Let X be a random variable with distribution function

$$F(x) = \begin{cases} 0, & x < 0, \\ 1/4, & 0 \leq x < 1, \\ 1/3, & 1 \leq x < 3, \\ 1/2, & 3 \leq x < 6, \\ 1, & x \geq 6. \end{cases}$$

Try to compute $P(X < 3)$, $P(X \leq 3)$.

Solution. We first observe that X is discrete and thus

$$P(X = 0) = F(0) - \lim_{y \rightarrow 0^-} F(y) = \frac{1}{4}, \quad P(X = 1) = F(1) - \lim_{y \rightarrow 1^-} F(y) = \frac{1}{12},$$

$$P(X = 3) = F(3) - \lim_{y \rightarrow 3^-} F(y) = \frac{1}{6}, \quad P(X = 6) = F(6) - \lim_{y \rightarrow 6^-} F(y) = \frac{1}{2}.$$

Thus

$$P(X < 3) = P(X = 0) + P(X = 1) = \frac{1}{3}, \quad P(X \leq 3) = 1 - P(X = 6) = \frac{1}{2}.$$

□

 **Exercise 4.3** Let X be a random variable with distribution function


$$F(x) = \begin{cases} \frac{e^x}{2}, & x < 0, \\ \frac{1}{2}, & 0 \leq x < 1, \\ 1 - \frac{1}{2}e^{-\frac{1}{2}(x-1)} & x \geq 1. \end{cases}$$

Compute $\text{Var}(X)$.

Solution. The probability density function of X is given by

$$p(x) = F'(x) = \begin{cases} \frac{e^x}{2}, & x < 0, \\ 0, & 0 \leq x < 1, \\ \frac{1}{4}e^{-\frac{1}{2}(x-1)} & x \geq 1. \end{cases}$$

□

 **Exercise 4.4** Let $g : \mathbb{R} \rightarrow \mathbb{R}$ be a non-decreasing function such that $\mathbb{E}(g(X))$ exists. Show that for every $\varepsilon > 0$,

$$\mathbb{P}(X > \varepsilon) \leq \frac{\mathbb{E}(g(X))}{g(\varepsilon)}.$$

Proof. Let $p(x)$ denote the probability density function of X . Then

$$\mathbb{P}(X > \varepsilon) = \int_{\varepsilon}^{\infty} p(x) dx = \int_{\varepsilon}^{\infty} \frac{g(x)}{g(x)} p(x) dx \leq \int_{\varepsilon}^{\infty} \frac{g(x)}{g(\varepsilon)} p(x) dx = \frac{1}{g(\varepsilon)} \int_{\varepsilon}^{\infty} g(x) p(x) dx \leq \frac{\mathbb{E}(g(X))}{g(\varepsilon)}.$$

□


 **Exercise 4.5** Let X be a non-negative continuous random variable. Prove that for $x \geq 0$,

$$\mathbb{P}(X < x) \geq 1 - \frac{\mathbb{E}(X)}{x}.$$

Proof. Let $p(x)$ denote the probability density function of X . Then

$$\begin{aligned}\mathbb{P}(X < x) &= \int_0^x p(t)dt = 1 - \int_x^\infty p(t)dt \geq 1 - \int_x^\infty \frac{t}{x}p(t)dt \\ &\geq 1 - \frac{1}{x} \int_0^\infty tp(t)dt = 1 - \frac{\mathbb{E}(X)}{x}.\end{aligned}$$

□

 **Exercise 4.6** Let X be a random variable taking values in \mathbb{N} . If $\text{var}(X)$ exists, show that

$$\text{var}(X) = 2 \sum_{n=1}^{\infty} n\mathbb{P}(X \geq n) - \mathbb{E}(X)[\mathbb{E}(X) + 1].$$

Proof. By the existence of variance, we know that the series $\sum_{n=1}^{\infty} n^2\mathbb{P}(X = n)$ is absolutely convergent and thus

$$\text{var}(X) = \mathbb{E}(X^2) - \mathbb{E}^2(X) = \mathbb{E}(X^2) + \mathbb{E}(X) - \mathbb{E}(X)[\mathbb{E}(X) + 1],$$

where

$$\begin{aligned}\mathbb{E}(X^2) + \mathbb{E}(X) &= \sum_{n=1}^{\infty} n(n+1)\mathbb{P}(X = n) = 2 \sum_{n=1}^{\infty} \left(\sum_{i=1}^n i \right) \mathbb{P}(X = n) \\ &= 2 \sum_{i=1}^{\infty} i \left[\sum_{n=i}^{\infty} \mathbb{P}(X = n) \right] = 2 \sum_{i=1}^{\infty} i\mathbb{P}(X \geq i).\end{aligned}$$

□

4.3 Lebesgue integration

Let (E, \mathcal{E}, μ) be a measure space.

We want to define, where possible, for measurable functions $f : E \rightarrow [-\infty, \infty]$, the integral of f with respect to the measure μ :

$$\mu(f) = \int_E f d\mu = \int_{x \in E} f(x)\mu(dx).$$

For random variables on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, the integral will be called the expectation of X , and written $\mathbb{E}(X)$.

Definition 4.7 (Simple functions)

A simple function is a function of the form

$$f = \sum_{k=1}^m a_k \mathbb{1}_{A_k},$$

where $0 < a_k \leq \infty$, and each $A_k \in \mathcal{E}$ is **disjoint with each other**.



Definition 4.8 (Integration of simple functions)

We define the integral of a simple function f to be

$$\mu(f) := \sum_{k=1}^m a_k \mu(A_k),$$

with the convention that $0 \cdot \infty = 0$.



Example 4.4 $\mu(\mathbb{1}_A) = \int_A 1 dx = \mu(A)$, for any measurable $A \in \mathcal{E}$.

Example 4.5 On probability space we have $\mathbb{E}(\mathbb{1}_A) = \int_A 1 d\mathbb{P} = \mathbb{P}(A)$.

Moreover for simple functions f, g and constants $\alpha, \beta \geq 0$ the following set of properties, **(P)**, holds:

1. $\mu(\alpha f + \beta g) = \mu\alpha(f) + \beta\mu(g)$;
2. $f \leq g \implies \mu(f) \leq \mu(g)$;
3. $f = 0$ a.e. $\iff \mu(f) = 0$.

Definition 4.9 (Integration of general measurable functions)

For general non-negative measurable functions f , we define the integral by

$$\mu(f) := \sup\{\mu(g) : g \text{ is simple, } g \leq f\}.$$



Note that property **(P)**(2) implies that this definition is consistent with the definition of μ for simple functions. Also the properties **(P)** hold for f, g non-negative measurable functions. Let

$$f^+ := f \vee 0 \text{ and } f^- := -f \vee 0.$$

Then

$$f = f^+ - f^- \text{ and } |f| = f^+ + f^-.$$

We say that a measurable function f is integrable if $\mu(|f|) < \infty$, and define

$$\mu(f) := \mu(f^+) - \mu(f^-).$$


Remark Combining with property **(P)**(2), we know f is integrable if and only if $\int |f| d\mu < +\infty$.

Example 4.6 $\sin x$ is not integrable with respect to Lebesgue measure since

$$\int_{\mathbb{R}} |\sin x| dx \rightarrow \infty.$$

Example 4.7 $\frac{\sin x}{x}$ is not integrable with respect to Lebesgue measure since

$$\begin{aligned} \int_{\mathbb{R}} \left| \frac{\sin x}{x} \right| dx &\geq \sum_{n=1}^{\infty} \int_{(n-1)\pi}^{n\pi} \frac{|\sin x|}{x} dx \geq \sum_{n=1}^{\infty} \frac{1}{n\pi} \int_{(n-1)\pi}^{n\pi} |\sin x| dx \\ &\geq \sum_{n=1}^{\infty} \frac{2}{n\pi}. \end{aligned}$$

 **Exercise 4.7** Prove that $\mu(A) = 0$ implies $\int_A f d\mu = 0$ for a Borel function f .

Proof. Following the standard machine. □

Proposition 4.1

Let A be a measurable subset of E , then

$$\int_A f(x) dx = \int_E f(x) \mathbb{1}_A(x) dx.$$



Proof. Note that

$$\begin{aligned} \int_A f(x) dx &= \sup_{\substack{h(x) \leq f(x) \\ x \in A}} \left\{ \int_A h(x) dx \right\} \\ &= \sup_{\substack{h(x) \mathbb{1}_A(x) \leq f(x) \mathbb{1}_A(x) \\ x \in E}} \left\{ \int_A h(x) dx \right\} = \int_E f(x) \mathbb{1}_A(x) dx. \end{aligned}$$

□

Proposition 4.2

$f(x) = 0$ a.e. if and only if $\int_E f(x)dx = 0$ (If $m(E) = 0$, then $\int_E f(x)dx = 0$).

♠

Proof. The “ \implies ” is trivial. Conversely, we define

$$E_k := \{x \in E : f(x) > 1/k\}.$$

Since

$$\frac{1}{k}m(E_k) = \int_{E_k} \frac{1}{k}dx \leq \int_{E_k} f(x)dx \leq \int_E f(x)dx = 0$$

we have $m(E_k) = 0$ ($k = 1, 2, \dots$). Note that

$$\{x \in E : f(x) > 0\} = \bigcup_{k=1}^{\infty} E_k,$$

and thus $m(\{f > 0\}) = 0$.

□

Theorem 4.1 (Monotone Convergence Theorem(M.C.T))

Let $\{f_n\}_{n \in \mathbb{N}}$ be a non-decreasing sequence of measurable functions with $f_n \geq 0$ and $\lim_{n \rightarrow \infty} f_n = f$. Then

$$\lim_{k \rightarrow \infty} \int_E f_k(x)dx = \int_E f(x)dx. \quad (4.6)$$

♡

Proof. It is easy to see that f is measurable and $\int_E f(x)dx$ is well-defined. Since

$$\int_E f_k(x)dx \leq \int_E f_{k+1}(x)dx \quad (k = 1, 2, \dots),$$

and thus $\lim_{k \rightarrow \infty} \int_E f_k(x)dx$ is well-defined. From the non-decreasing property, we see

$$\lim_{k \rightarrow \infty} \int_E f_k(x)dx \leq \int_E f(x)dx.$$

Now let $0 < c < 1$ and $h(x)$ be any non-negative simple function on \mathbb{R}^n such that $h(x) \leq f(x)$, $x \in E$. Define

$$E_k := \{x \in E : f_k(x) \geq ch(x)\} \quad (k = 1, 2, \dots),$$

then E_k is measurable and increasing with $\lim_{k \rightarrow \infty} E_k = E$. By Theorem ??,

$$\lim_{k \rightarrow \infty} c \int_{E_k} h(x)dx = c \int_E h(x)dx.$$

From inequality

$$\int_E f_k(x)dx \geq \int_{E_k} f_k(x)dx \geq \int_{E_k} ch(x)dx = c \int_{E_k} h(x)dx$$

we get

$$\lim_{k \rightarrow \infty} \int_E f_k(x)dx \geq c \int_E h(x)dx.$$

Send $c \rightarrow 1$, obtaining

$$\lim_{k \rightarrow \infty} \int_E f_k(x)dx \geq \int_E h(x)dx.$$

By the “sup” in the definition of integral, we get

$$\lim_{k \rightarrow \infty} \int_E f_k(x) dx \geq \int_E f(x) dx.$$

□

4.4 Absolute continuity and Radon-Nikodym derivative

Definition 4.10

Let $(\Omega, \mathcal{F}, \mu)$ be a measure space. If for any $A \in \mathcal{F}$, one has $v(A) = \int_A f d\mu$ for a certain Borel function $f : \Omega \rightarrow [0, +\infty)$, then f is called the **Radon-Nikodym derivative** of v with respect to μ , denoted by $f = \frac{dv}{d\mu}$.



Proposition 4.3

Let $(\Omega, \mathcal{F}, \mu)$ be a measure space and $f : \Omega \rightarrow [0, +\infty)$ be a Borel function. Then

$$v(A) = \int_A f d\mu$$

defines a measure.



Proof. • From Exercise 4.7 we know if $A = \emptyset$ then $v(A) = 0$.

• Now we continue to prove the countable additivity. Let $\{A_j\}$ be a countable disjoint set sequence. Then

$$\begin{aligned} v\left(\bigcup_{j=1}^{\infty} A_j\right) &= \int_{\bigcup_{j \geq 1} A_j} f d\mu = \int_{\Omega} f \cdot \mathbb{1}_{\bigcup_{j \geq 1} A_j} d\mu \\ &= \int_{\Omega} f \lim_{n \rightarrow \infty} \mathbb{1}_{\bigcup_{j=1}^n A_j} d\mu = \lim_{n \rightarrow \infty} \int_{\Omega} f \cdot \mathbb{1}_{\bigcup_{j=1}^n A_j} d\mu \\ &= \lim_{n \rightarrow \infty} \int_{\Omega} f \cdot \sum_{j=1}^n \mathbb{1}_{A_j} d\mu = \lim_{n \rightarrow \infty} \sum_{j=1}^n v(A_j) \\ &= \sum_{j \geq 1} v(A_j). \end{aligned}$$

□

Definition 4.11

We say v is **absolutely continuous with respect to** μ , denoted by $v \ll \mu$ if and only if $v(A) = 0$ given any $A \in \mathcal{F}$ such that $\mu(A) = 0$. In particular, if $v \ll \mu$ and $\mu \ll v$, we say they are **equivalent**, denoted by $\mu \sim v$.



Example 4.8 If $v(A) = \int_A f d\mu$ for a Borel f , then $v \ll \mu$.

Example 4.9 Define measure M to be $M := 2m$, where m is the Lebesgue measure. Then $M \ll m$ and $m \ll M$.

Example 4.10 Let m be the Lebesgue measure whereas c denote the counting measure, i.e.,

$$c(A) = \begin{cases} |A|, & |A| < +\infty, \\ +\infty, & |A| = +\infty, \end{cases}$$

for any set A . Then $m \ll c$ whereas the inversed direction is not true.

Theorem 4.2 (Radon-Nikodym)

If μ is σ -finite on Ω , i.e., Ω can be written as $\bigcup_{j \geq 1} A_j$ for $A_j \in \mathcal{F}$ and $\mu(A_j) < \infty$, and $\mu \ll \nu$, then there is a measurable function f such that

1. $\nu(A) = \int_A f d\mu$ for any $A \in \mathcal{F}$;
2. $\int_{\Omega} f d\mu = 1$;
3. $f \geq 0$ a.e. with respect to measure μ .


Proposition 4.4 (Equivalent character of absolute continuity)

$\nu \ll \mu \iff \forall \varepsilon > 0, \exists \delta > 0$ such that for any A such that $\mu(A) < \delta$, we have $\nu(A) < \varepsilon$.



Proof. \implies . Assume the contrary: $\exists \varepsilon_0 > 0, \forall \delta > 0$, there exists A with $\mu(A) < \delta$ but $\nu(A) \geq \varepsilon_0$. Now for any $n \in \mathbb{N}$, take A_n such that $\mu(A_n) \leq 1/2^n$ and $\nu(A_n) \geq \varepsilon_0$. Defining

$$B_n = \bigcup_{k \geq n} A_k,$$

$$B = \limsup A_n = \bigcap_{n \geq 1} \bigcup_{k \geq n} A_k = \bigcap_{n \geq 1} B_n,$$

we have

$$\mu(B_n) \leq \sum_{k \geq n} \mu(A_k) \leq \sum_{k \geq n} \frac{1}{2^k} = \frac{1}{2^{n-1}}$$

and

$$\nu(B_n) \geq \nu(A_n) \geq \varepsilon_0.$$

By continuity of measure, $\mu(B) = \lim_{n \rightarrow \infty} \mu(B_n) = 0$ whereas $\nu(B) = \lim_{n \rightarrow \infty} \nu(B_n) = \varepsilon_0 > 0$, contradicting with $\nu \ll \mu$. \square

Remark Absolutely random variable \iff its distribution \mathbb{P}_X is absolutely continuous with respect to Lebesgue measure.

Proposition 4.5 (Chain rule)

$\nu \ll \mu$ and f is integrable with respect to $\nu \iff f \frac{d\nu}{d\mu}$ is integrable with respect to μ and $\int f d\nu = \int f \frac{d\nu}{d\mu} \cdot d\mu$.



Remark Write in probabilistic way: if $Q \ll P$, then

$$E^Q[X] = E^P[X \cdot \frac{dQ}{dP}].$$

Proof. Only the " \implies " direction.

- If $f = \mathbf{1}_A$, where $A \in \mathcal{F}$, then

$$\int \mathbf{1}_A d\mu = \nu(A) = \int_A \frac{d\nu}{d\mu} \cdot d\mu = \int \mathbf{1}_A \frac{d\nu}{d\mu} \cdot d\mu$$

- By linearity, if $\varphi = \sum_{i=1}^n a_i \mathbb{1}_{A_i}$, $a_i \geq 0$, $A_i \cap A_j = \emptyset$, then

$$\int \varphi d\nu = \int \varphi \frac{d\nu}{d\mu} \cdot d\mu.$$

- Let $f \geq 0$ be a non-negative Borel function, then there is increasing measurable simple function sequence $\{\varphi_k\}$ such that $\varphi_k \rightarrow f$. Then

$$\int f d\nu = \lim_{k \rightarrow \infty} \int \varphi_k d\nu = \lim_{k \rightarrow \infty} \int \varphi_k \frac{d\nu}{d\mu} \cdot d\mu = \int \lim_{k \rightarrow \infty} \varphi_k \frac{d\nu}{d\mu} \cdot d\mu = \int f \frac{d\nu}{d\mu} \cdot d\mu.$$

- Let f be a general Borel function. Then write f as $f = f^+ - f^-$.

□

Example 4.11 Let (Ω, \mathcal{F}, P) be a probability space. Suppose a random variable X is $X \sim \mathcal{N}(0, 1)$. For $\theta \in \mathbb{R}$, there is a new probability measure Q such that under Q , $X + \theta \sim \mathcal{N}(0, 1)$.

Proof. Take $\frac{dQ}{dP} = e^{-\theta X - \frac{1}{2}\theta^2}$. Then

$$\begin{aligned} E^Q(e^{t(X+\theta)}) &= E^P \left[e^{t(X+\theta)} \frac{dQ}{dP} \right] = E^P [e^{tX} e^{-\theta X} e^{-\frac{1}{2}\theta^2} e^{\theta t}] \\ &= E^P [e^{(t-\theta)X}] e^{\theta t - \frac{1}{2}\theta^2} = e^{\frac{1}{2}(t-\theta)^2} e^{\theta t - \frac{1}{2}\theta^2} \\ &= e^{\frac{1}{2}t^2}. \end{aligned}$$

Thus $X + \theta \stackrel{Q}{\sim} \mathcal{N}(0, 1)$.

□

4.5 Common discrete distribution

Example 4.12(Bernoulli) Let X be a random variable such that $\mathbb{P}(X = 1) = p$ and $\mathbb{P}(X = 0) = q = 1 - p$. Then

$$\mathbb{E}(X) = 0 \cdot \mathbb{P}(X = 0) + 1 \cdot \mathbb{P}(X = 1) = p, \quad \text{var}(X) = \mathbb{E}(X^2) - \mathbb{E}(X)^2 = p - p^2 = pq.$$

Example 4.13(Binomial(n, p)) Consider an unfair coin with probability p to be a head and probability $q = 1 - p$ to be a tail after tossing up. Let X denote the number of heads in n coin flips. Then

$$\begin{aligned} f(k) &= \mathbb{P}(X = k) = \binom{n}{k} p^k q^{n-k}, \\ \mathbb{E}(X) &= \sum_{k=1}^n k \binom{n}{k} p^k q^{n-k} = np. \end{aligned}$$

In fact, consider the Binomial expression

$$(1 + x)^n = \sum_{k=0}^n \binom{n}{k} x^k.$$

Taking derivative with respect to x , obtaining

$$n(1 + x)^{n-1} = \sum_{k=1}^n \binom{n}{k} k x^{k-1}$$

Thus

$$n(1 + x)^{n-1} x = \sum_{k=1}^n \binom{n}{k} k x^k$$

Now set $x = p/q$ and multiply q^n on both sides, we get $\mathbb{E}(X) = np$.

Alternatively, set $X = Y_1 + Y_2 + \cdots + Y_n$, where Y_i takes value 1 if a head occurs in toss i and takes value 0 if a tail occurs in toss i . Then Y_i are i.i.d random variable obeying Bernoulli distribution and thus

$$\begin{aligned}\mathbb{E}(X) &= \mathbb{E}\left(\sum_{i=1}^n Y_i\right) = \sum_{i=1}^n \mathbb{E}(Y_i) = np; \\ \text{var}(X) &= \mathbb{E}((X - \mathbb{E}(X))^2) = \mathbb{E}\left[\left(\sum_{i=1}^n (Y_i - \mathbb{E}(Y_i))\right)^2\right] \\ &= \sum_{i=1}^n \mathbb{E}[(Y_i - \mathbb{E}(Y_i))^2] + 2 \sum_{i \neq j} \mathbb{E}[(Y_i - \mathbb{E}(Y_i))(Y_j - \mathbb{E}(Y_j))] \\ &= \sum_{i=1}^n \text{var}(Y_i) = npq.\end{aligned}$$

Example 4.14 A random variable X taking value in \mathbb{N} is said to be a **Poisson random variable** with parameter λ if

$$f(i) = \mathbb{P}(X = i) = \frac{\lambda^i}{i!} e^{-\lambda}, \quad \mathbb{E}(X) = \lambda.$$

In fact, Poisson arises from Binomial in the limit $n \rightarrow \infty$ but $np = \lambda$:

$$\begin{aligned}\mathbb{P}(X = i) &= \binom{n}{i} p^i (1-p)^{n-i} = \binom{n}{i} \left(\frac{\lambda}{n}\right)^i \left(1 - \frac{\lambda}{n}\right)^{n-i} \\ &= \frac{n(n-1) \cdots (n-i+1)}{i!} \frac{\lambda^i}{n^i} \left(1 - \frac{\lambda}{n}\right)^n \left(1 - \frac{\lambda}{n}\right)^{-i} \\ &\stackrel{n \rightarrow \infty}{\approx} \frac{\lambda^i}{i!} e^{-\lambda}.\end{aligned}$$

Exercise 4.8 Find the expectation of $X(X-1) \cdots (X-k+1)$, where X is a random variable obeying the Poisson distribution.

Solution. To evaluate the moments of the Poisson random variable, we use a little inspiration to observe that for $k \geq 1$,

$$\begin{aligned}E(X(X-1) \cdots (X-k+1)) &= \sum_{j=k}^{\infty} j(j-1) \cdots (j-k+1) e^{-\lambda} \frac{\lambda^j}{j!} \\ &= \lambda^k \sum_{j=k}^{\infty} e^{-\lambda} \frac{\lambda^{j-k}}{(j-k)!} = \lambda^k\end{aligned}$$

where the equalities follow from (i) the facts that $j(j-1) \cdots (j-k+1) = 0$ when $j < k$, (ii) cancelling part of the factorial, and (iii) the fact that Poisson distribution has total mass 1. Using the last formula, it follows that $EX = \lambda$ while

$$\text{var}(X) = EX^2 - (EX)^2 = E(X(X-1)) + EX - \lambda^2 = \lambda$$

□

Example 4.15 A random variable X is said to be **Geometric**(p) if

$$\mathbb{P}(X = k) = (1-p)^{k-1} \cdot p \quad k \in \mathbb{N},$$

i.e., denoting the first success in independent trials with success probability p . Show that a geometric random variable X has expectation $\mathbb{E}(X) = 1/p$ and variance $\text{var}(X) = 1 - p/p^2$.

Proof. Let $q := 1 - p$. Then

$$\begin{aligned} E(X) &= \sum_{k=1}^{\infty} k p q^{k-1} = p \sum_{k=1}^{\infty} k q^{k-1} = p \sum_{k=1}^{\infty} \frac{dq^k}{dq} \\ &= p \frac{d}{dq} \left(\sum_{k=1}^{\infty} q^k \right) = p \frac{d}{dq} \left(\frac{1}{1-q} \right) = \frac{p}{(1-q)^2} = \frac{1}{p}. \end{aligned}$$

Note that

$$\begin{aligned} E(X^2) &= \sum_{k=1}^{\infty} k^2 p q^{k-1} = p \left[\sum_{k=1}^{\infty} k(k-1) q^{k-1} + \sum_{k=1}^{\infty} k q^{k-1} \right] \\ &= p q \sum_{k=1}^{\infty} k(k-1) q^{k-2} + \frac{1}{p} = p q \sum_{k=1}^{\infty} \frac{d^2}{dq^2} q^k + \frac{1}{p} \\ &= p q \frac{d^2}{dq^2} \left(\sum_{k=0}^{\infty} q^k \right) + \frac{1}{p} = p q \frac{d^2}{dq^2} \left(\frac{1}{1-q} \right) + \frac{1}{p} \\ &= p q \frac{2}{(1-q)^3} + \frac{1}{p} = \frac{2q}{p^2} + \frac{1}{p} \end{aligned}$$

and thus

$$\text{var}(X) = E(X^2) - E^2(X) = \frac{2q}{p^2} + \frac{1}{p} - \frac{1}{p^2} = \frac{1-p}{p^2}.$$

□

Example 4.16 Consider tossing an unfair coin with probability p ($0 < p < 1$) to get a head. What is the expectation of tossing times such that a head and a tail have both occurred?

Solution. We use X to denote the number of tossing times when a head and a tail have both occurred. Then X take values from 2, 3, \dots , and

$$P(X = k) = (1-p)^{k-1} p + p^{k-1} (1-p), \quad k = 2, 3, \dots$$

Thus

$$\begin{aligned} E(X) &= \sum_{k=2}^{\infty} k [(1-p)^{k-1} p + p^{k-1} (1-p)] \\ &= \sum_{k=1}^{\infty} k (1-p)^{k-1} p - p + \sum_{k=1}^{\infty} k p^{k-1} (1-p) - (1-p) \\ &= \frac{1}{p} + \frac{1}{1-p} - 1 = \frac{1}{p(1-p)} - 1. \end{aligned}$$

□

Example 4.17(Coupon collector) Suppose there are N types of coupons. Define X to be

$$X := \text{the first time to get a complete set } \{1, 2, \dots, N\}.$$

Try to compute $\mathbb{E}(X)$.

Solution. Define $Y_k :=$ first time to get k distinct coupons and thus $X = Y_N = \sum (Y_k - Y_{k-1}) + Y_1$. Note that $Y_k - Y_{k-1} \sim \text{Geo} \left(\frac{N - (k-1)}{N} \right)$ and thus

$$\mathbb{E}(Y_N) = \sum_{k=2}^N \mathbb{E}(Y_k - Y_{k-1}) + \mathbb{E}(Y_1) = \sum_{k=2}^N \frac{N}{N - (k-1)} + 1$$

$$= 1 + N\left(1 + \frac{1}{2} + \cdots + \frac{1}{N-1}\right) \sim N \ln N + \gamma N + O\left(\frac{1}{N}\right),$$

where $\gamma \approx 0.5772156649$ is the Euler-Mascheroni constant. \square

Example 4.18(Exponential) Let X be a random variable on measure space $(\mathbb{R}, \mathcal{B}(\mathbb{R}), m)$ with probability density function $f(x) = \lambda e^{-\lambda x} \mathbb{1}_{[0, +\infty)}(x)$ and distribution function $F(x) = 1 - e^{-\lambda x}$. Then

$$\mathbb{P}_\lambda(B) := \int_B f(x) dx$$

defines a probability measure with

$$\mathbb{P}_\lambda(\mathbb{R}) := \int_0^\infty e^{-\lambda x} dx = 1.$$

X is then called obeying an **exponential**(λ) distribution, denoted by $X \sim \text{Exp}(\lambda)$.

Exercise 4.9 Show that if $X \sim \text{Exp}(\lambda)$, then $\mathbb{E}(X) = 1/\lambda$ and $\text{var}(X) = 1/\lambda^2$.

Solution.

$$\begin{aligned} \mathbb{E}(X) &= \int_0^\infty x \lambda e^{-\lambda x} dx = \frac{\lambda}{-\lambda} \int_\infty^0 x de^{-\lambda x} \\ &= \int_0^\infty x de^{-\lambda x} = x e^{-\lambda x} \Big|_0^\infty - \int_0^\infty e^{-\lambda x} dx \\ &= \frac{1}{\lambda}. \end{aligned}$$

Recall Lemma 4.3,

$$\mathbb{E}(X^2) = \int_0^\infty x^2 \lambda e^{-\lambda x} dx = \int_0^\infty x^2 d(-e^{-\lambda x}) = -x^2 e^{-\lambda x} \Big|_0^\infty + 2 \int_0^\infty x e^{-\lambda x} dx = \frac{2}{\lambda^2}.$$

Thus

$$\text{var}(X) = \mathbb{E}(X^2) - \mathbb{E}^2(X) = \frac{2}{\lambda^2} - \frac{1}{\lambda^2} = \frac{1}{\lambda^2}. \quad \square$$

Exercise 4.10 Let $X \sim \text{Exp}(\lambda)$.

1. (Lack of memory) Show that for any $s, t > 0$, $\mathbb{P}[X > t + s \mid X > s] = \mathbb{P}[X > t]$.
2. Find all continuous random variable such that $\mathbb{P}[X > t + s \mid X > s] = \mathbb{P}[X > t]$.

Solution. 1. Note that

$$\mathbb{P}(X \geq t) = \int_t^\infty e^{-\lambda x} dx = e^{-\lambda t}$$

and

$$\begin{aligned} \mathbb{P}[X > t + s \mid X > s] &= \frac{\mathbb{P}(X > t + s \cap X > s)}{\mathbb{P}(X > s)} = \frac{\mathbb{P}(X > t + s)}{\mathbb{P}(X > s)} \\ &= \frac{\int_{t+s}^\infty \lambda e^{-\lambda x} dx}{\int_s^\infty \lambda e^{-\lambda x} dx} = \frac{e^{-\lambda(t+s)}}{e^{-\lambda s}} \\ &= e^{-\lambda t}. \end{aligned}$$

2. By condition we have

$$\frac{\mathbb{P}(X > t + s \cap X > s)}{\mathbb{P}(X > s)} = \frac{\mathbb{P}(X > t + s)}{\mathbb{P}(X > s)} = \mathbb{P}(X > t)$$

and thus $H(x) := 1 - F(x)$, where $F(x)$ denote the distribution function of X , satisfies

$$H(t + s) = H(x) \cdot H(s), \quad s, t > 0.$$

Define $G = \ln H$ and then

$$G(t + s) = G(t) + G(s) \quad s, t > 0,$$

implying that G is indeed a linear function, i.e., $G(x) = cx$ for some $c \in \mathbb{R}$. It follows that $F(x) = 1 - H(x) = 1 - e^{cx}$. □

Example 4.19 Let $[0, t]$ be a given time period and $N(t)$ denote the number of times that a breakdown occurs in the period $[0, t]$. If $N(t)$ obeys Poisson distribution with parameter λt , show that T , the interval between two different breakdowns, obeys the exponential distribution.

Proof. By condition, $N(t) \sim \text{Poisson}(\lambda t)$, i.e.,

$$P(N(t) = k) = \frac{(\lambda t)^k}{k!} e^{-\lambda t}, \quad k = 0, 1, \dots$$

Note that $\{T \geq t\} = \{N(t) = 0\}$, and thus

- for $t < 0$, $F_T(t) = P(T \leq t) = 0$;
- for $t \geq 0$,


$$F_T(t) = P(T \leq t) = 1 - P(T > t) = 1 - P(N(t) = 0) = 1 - e^{-\lambda t},$$

implying $T \sim \exp(\lambda)$. □

Example 4.20 (Gamma distribution) Consider the probability mass function $P_{\lambda, t}(A) := \int_A \frac{1}{\Gamma(t)} \lambda^t x^{t-1} e^{-\lambda x} dx$

for $\lambda, t \geq 0$, where $\Gamma(t) := \int_0^\infty x^{t-1} e^{-x} dx$ is called the **Gamma function**. The Gamma function has the following properties:

- $\Gamma(1) = 1$, $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$ and $f_{\lambda, 1} = \lambda e^{-\lambda x}$;
- $\Gamma(t + 1) = t\Gamma(t)$ for $t > 0$;
- $\Gamma(n) = (n - 1)\Gamma(n - 1) = (n - 1)!$ and $f_{\lambda, n} = \frac{\lambda^n}{(n - 1)!} e^{-\lambda x}$;

 **Exercise 4.11** Compute $E(X)$ and $\text{var}(X)$ for a Gamma distributed random variable X .

Solution. Using properties of a Gamma function,

$$E(X) = \frac{\lambda^t}{\Gamma(t)} \int_0^\infty x^t e^{-\lambda x} dx = \frac{\Gamma(t + 1)}{\Gamma(t)} \frac{1}{\lambda} = \frac{t}{\lambda}.$$

Furthermore,

$$E(X^2) = \frac{\lambda^t}{\Gamma(t)} \int_0^\infty x^{t+1} e^{-\lambda x} dx = \frac{\Gamma(t + 2)}{\lambda^2 \Gamma(t)} = \frac{t(t + 1)}{\lambda^2}.$$

Thus

$$\text{var}(X) = E(X^2) - E^2(X) = \frac{t(t + 1)}{\lambda^2} - \left(\frac{t}{\lambda}\right)^2 = \frac{t}{\lambda^2}. \quad \square$$

Theorem 4.3 (Convolution)

Let X and Y be two independent random variable with density function $f_X(x)$ and $f_Y(y)$, respectively.

Then $Z := X + Y$ has a density function

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(z - y) f_Y(y) dy = \int_{-\infty}^{\infty} f_X(x) f_Y(z - x) dx. \quad (4.7)$$



Proof. Note that by independence

$$\begin{aligned} F_Z(z) &= P(X + Y \leq z) = \iint_{x+y \leq z} f_X(x)f_Y(y)dx dy \\ &= \int_{-\infty}^{\infty} \left[\int_{-\infty}^{z-y} f_X(x)dx \right] f_Y(y)dy = \int_{-\infty}^{\infty} \int_{-\infty}^z f_X(t-y)f_Y(y)dt dy \\ &= \int_{-\infty}^z \left(\int_{-\infty}^{\infty} f_X(t-y)f_Y(y)dy \right) dt. \end{aligned}$$

Hence the density function of Z is


$$f_Z(z) = \int_{-\infty}^{\infty} f_X(z-y)f_Y(y)dy.$$

Now set $z - y = x$, we get

$$f_Z(z) = \int_{-\infty}^{\infty} f_X(x)f_Y(z-x)dx.$$

□

Proposition 4.6

Let X_1, X_2, \dots, X_n be independent random variables obeying the exponential(λ) distribution. Then $X_1 + X_2 + \dots + X_n$ is a random variable with Gamma (λ, n) distribution. 

Proof. First consider the case of discrete. Note that for a fixed $z \in \mathbb{R}$,

$$\begin{aligned} P(X + Y = z) &= \sum_x P(X = x, Y = z - x) = \sum_x P(X = x)P(Y = z - x) \\ &= \sum_x f_X(x)f_Y(z - x) \end{aligned}$$


Furthermore under the condition of continuous,

$$f_{X+Y}(z) = \int_{-\infty}^{\infty} f_X(x)f_Y(z-x)dx.$$

□


Definition 4.12

A **Poisson process** $(N_s)_{s \geq 0}$ with rate λ satisfies

1. $N_0 = 0$;
2. $N_t - N_s \sim \text{Poisson}(\lambda(t-s))$ for any $0 \leq s \leq t$;
3. $N_{t_1}, N_{t_2} - N_{t_1}, \dots, N_{t_n} - N_{t_{n-1}}$ are independent for any $0 \leq t_1 \leq t_2 \leq \dots \leq t_n$. 

Alternative construction We will see an alternative construction to generate a Poisson process:

Proposition 4.7

Let $\tau_1, \tau_2, \dots, \tau_n$ be independent random variables with exponential(λ) distribution. Define $T_n = \sum_{i=1}^n \tau_i$ and $N_s = \max\{n : T_n \leq s\}$. Then $(N_s)_{s \geq 0}$ is a Poisson process with rate λ . 

Example 4.21 In figure ??, N_s is equal to 2.

Proof of Proposition 4.7. • $N_0 = 0$ is trivial;


- Now assume $s = 0$. Then

$$\begin{aligned}
 P(N_t = n) &= P(T_n \leq t, T_{n+1} > t) = \int_0^t f_{T_n}(s)P(T_{n+1} > t | T_n = s)ds \\
 &= \int_0^t f_{T_n}(s)P(\tau_{n+1} > t - s)ds \\
 &= \int_0^t \frac{\lambda^n}{(n-1)!} s^{n-1} e^{-\lambda s} e^{-\lambda(t-s)} ds \\
 &= \frac{\lambda^n}{(n-1)!} e^{-\lambda t} \int_0^t s^{n-1} ds \\
 &= \frac{\lambda^n t^n}{n!} e^{-\lambda t}.
 \end{aligned}$$

- Finally we show $N_{t+s} - N_s \sim \text{Poisson}(\lambda, t)$. Assume $N_s = n$ and by lack of memory,

$$P(\tau_{n+1} > s + t - T_n | \tau_{n+1} > s - T_n) = P(\tau_{n+1} > t) = P(\tau_1 > t) = e^{-\lambda t}.$$

□

 **Exercise 4.12** For $X \sim \text{Poisson}(\lambda)$, show that

$$E(X^n) = \lambda E[(X + 1)^{n-1}].$$

Proof.

$$\begin{aligned}
 E(X^n) &= \sum_{k=0}^{\infty} k^n \frac{\lambda^k}{k!} e^{-\lambda} = \lambda e^{-\lambda} \sum_{k=1}^{\infty} k^{n-1} \frac{\lambda^{k-1}}{(k-1)!} \\
 &\stackrel{k'=k-1}{=} \lambda e^{-\lambda} \sum_{k'=0}^{\infty} (k'+1)^{n-1} \frac{\lambda^{k'}}{k'!} \\
 &= \lambda E[(X + 1)^{n-1}].
 \end{aligned}$$

□

Example 4.22(Uniform distribution) A random variable X is **uniformly distributed** if

$$f_X(x) = \begin{cases} 1, & x \in [0, 1] \\ 0, & \text{otherwise.} \end{cases}$$

Then $P(a \leq X \leq b) = b - a$.

Example 4.23 Let $X \sim \text{unif}[0, 1]$ and $Y \sim \text{unif}[0, 1]$ be independent. What is $f_{X+Y}(a)$?

Solution.

$$f_{X+Y}(a) = \int_{-\infty}^{+\infty} f_X(x) f_Y(a-x) dx = \begin{cases} \int_0^a 1 dx = a, & 0 < a < 1, \\ \int_{a-1}^1 1 dx = 2 - a, & 1 < a < 2. \end{cases}$$

□

Example 4.24 Let X_1, X_2, \dots, X_n be independent uniformly zero-one distributed.

1. Show that $P(X_1 + \dots + X_n \leq x) = \frac{x^n}{n!}$ for $0 \leq x \leq 1$;
2. Let $N := \{n : X_1 + \dots + X_n \geq 1\}$. Compute $E(N)$.

Proof. 1. By induction, assume $F_{n-1}(x) = \frac{x^{n-1}}{(n-1)!}$ for $x \in [0, 1]$. Then

$$\begin{aligned} F_n(z) &= \int_0^1 f_{X_n}(x)F_{n-1}(z-x)dx = \int_0^1 1 \cdot \frac{(z-x)^{n-1}}{(n-1)!} \mathbf{1}_{\{x \leq z\}} dx \\ &= \int_0^z \frac{(z-x)^{n-1}}{(n-1)!} dx = \frac{z^n}{n!}. \end{aligned}$$

2. Note that $P(N > n) = P(X_1 + \cdots + X_n \leq 1) = F_n(1)$. Then

$$E_N = \sum_{n=0}^{\infty} P(N > n) = \sum_{n=0}^{\infty} \frac{1}{n!} = e.$$

□

Example 4.25 (Normal distribution)

• We say a random variable X satisfies the **normal zero-one distribution**, denoted by $X \sim \mathcal{N}(0, 1)$ if its probability density function $f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$. Note that $\int_{-\infty}^{\infty} e^{-x^2/2} dx = \sqrt{2\pi}$.

• We say a random variable X satisfies the **normal distribution**, denoted by $X \sim \mathcal{N}(\mu, \sigma^2)$ if its probability density function $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$. We now have $E(X) = \mu$ and $\text{var}(X) = \sigma^2$. In fact,

$$\begin{aligned} E(X) &= \int_{-\infty}^{\infty} x \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx \\ &\stackrel{y=x-\mu}{=} \int_{-\infty}^{\infty} (y+\mu) \frac{1}{\sqrt{2\pi\sigma^2}} e^{-y^2/2\sigma^2} dy \\ &= \int_{-\infty}^{\infty} y \frac{1}{\sqrt{2\pi\sigma^2}} e^{-y^2/2\sigma^2} dy + \mu \\ &= \mu \end{aligned}$$

$$\begin{aligned} \text{var}(X) &= E(X^2) - E^2(X) = \int_{-\infty}^{\infty} \frac{x^2 e^{-\lambda x^2}}{\sqrt{2\pi}} dx - \mu^2 = \frac{d}{d\lambda} \int_{-\infty}^{\infty} \frac{-e^{-\lambda x^2}}{\sqrt{2\pi}} dx - \mu^2 \\ &= \frac{\sqrt{2}}{4} \frac{1}{\lambda\sqrt{\lambda}} - \mu^2 = \sigma^2. \end{aligned}$$

Proposition 4.8

If $X \sim \mathcal{N}(\mu, \sigma^2)$, then $Y = \frac{X - \mu}{\sigma} \sim \mathcal{N}(0, 1)$.



Proof. Note that

$$\begin{aligned} P(Y \leq a) &= P(X \leq a\sigma + \mu) = \int_{-\infty}^{a\sigma + \mu} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/2\sigma^2} dx \\ &\stackrel{y=\frac{x-\mu}{\sigma}}{=} \int_{-\infty}^a \frac{1}{\sqrt{\pi}} e^{-y^2/2} dy. \end{aligned}$$

□

Proposition 4.9

If $X \sim \mathcal{N}(0, 1)$, then $E(e^{tX}) = e^{t^2/2}$.



Proof. Note that

$$E(e^{tX}) = \int_{-\infty}^{\infty} \frac{e^{tx} e^{-x^2/2}}{\sqrt{2\pi}} dx$$

$$\begin{aligned}
&= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{1}{2}(x-t)^2} e^{\frac{t^2}{2}} dx \\
&= e^{t^2/2}.
\end{aligned}$$

□

🔥 **Exercise 4.13** Given $X_1 \sim \text{Exp}(\lambda_1)$, $X_2 \sim \text{Exp}(\lambda_2)$ being independent. Compute:

1. the distribution and density function of $\min\{X_1, X_2\}$;
2. $\mathbb{P}(X_1 < X_2)$.

Proof. 1. From the independence, for $t \geq 0$,

$$\begin{aligned}
\mathbb{P}(\min(X_1, X_2) \geq t) &= \mathbb{P}(\{X_1 \geq t\} \cap \{X_2 \geq t\}) \\
&= \mathbb{P}(\{X_1 \geq t\}) \cdot \mathbb{P}(\{X_2 \geq t\}) \\
&= \int_t^{\infty} \lambda_1 e^{-\lambda_1 x} dx \cdot \int_t^{\infty} \lambda_2 e^{-\lambda_2 x} dx \\
&= e^{-\lambda_1 t} \cdot e^{-\lambda_2 t} \\
&= e^{-(\lambda_1 + \lambda_2)t}
\end{aligned}$$

Thus

$$\begin{aligned}
F_{\min\{X_1, X_2\}}(t) &= \begin{cases} 0, & t < 0 \\ 1 - e^{-(\lambda_1 + \lambda_2)t}, & t \geq 0. \end{cases} \\
f_{\min\{X_1, X_2\}}(t) &= \begin{cases} 0, & t < 0, \\ (\lambda_1 + \lambda_2)e^{-(\lambda_1 + \lambda_2)t}, & t \geq 0. \end{cases}
\end{aligned}$$

2. Only a sketch solution:

$$\begin{aligned}
\mathbb{P}(X_1 < X_2) &= \int_0^{\infty} f_{X_1}(t) \mathbb{P}[X_2 > t \mid X_1 = t] dt \\
&= \int_0^{\infty} f_{X_1}(t) \frac{\mathbb{P}(X_2 > t) \cdot \mathbb{P}(X_1 = t)}{\mathbb{P}(X_1 = t)} dt \\
&= \int_0^{\infty} f_{X_1}(t) \mathbb{P}(X_2 > t) dt \\
&= \int_0^{\infty} \lambda_1 e^{-\lambda_1 t} e^{-\lambda_2 t} dt \\
&= \frac{\lambda_1}{\lambda_1 + \lambda_2}.
\end{aligned}$$


□

Example 4.26 Let $X \sim \mathcal{N}(0, 1)$ and $Y \sim \mathcal{N}(0, 1)$ be independent. Show that $X + Y \sim \mathcal{N}(0, 2)$.

Proof.

$$\begin{aligned}
f_{X+Y}(z) &= \int_{-\infty}^{\infty} f_X(x) f_Y(z-x) dx = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} \frac{1}{\sqrt{2\pi}} e^{-\frac{(z-x)^2}{2}} dx \\
&= \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{4}z^2} \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-(x-\frac{z}{2})^2} dx \\
&= \frac{1}{\sqrt{2\pi}\sqrt{2}} e^{-\frac{1}{4}z^2}.
\end{aligned}$$

□


 **Exercise 4.14** Let $X \sim \mathcal{N}(\mu_1, \sigma_1^2)$ and $Y \sim \mathcal{N}(\mu_2, \sigma_2^2)$ be independent. Show that $X + Y \sim \mathcal{N}(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$.

Proof. □

Theorem 4.4 (Bounded Convergence)

Let $(\Omega, \mathcal{F}, \mu)$ be a measure space such that $\mu(\Omega) < \infty$. Then if $f_n \rightarrow f$ a.e. and $\sup_{n \geq 1} |f_n| < \infty$, then

$$\lim_{n \rightarrow \infty} \int f_n d\mu = \int f d\mu.$$

 **Exercise 4.15** Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $X : \Omega \rightarrow [0, \infty)$ is a random variable such that $E(X^n) \leq M$ for some M and for any $n \in \mathbb{N}$.

1. Prove that $P(X > 1) = 0$ and $P(X = 1) \leq M$;
2. Compute $\lim_{n \rightarrow \infty} E(X^n)$;
3. If $E(X^n) = M$ for any $n \in \mathbb{N}$, show that $P(X \in \{0, 1\}) = 1$.

Solution. 1. For any $\varepsilon > 0$, note that

$$M \geq E(X^n) = \int_{\{0 \leq X\}} X^n d\mathbb{P} \geq \int_{\{X > 1 + \varepsilon\}} X^n d\mathbb{P} \geq (1 + \varepsilon)^n \int_{\{X > 1 + \varepsilon\}} d\mathbb{P} = (1 + \varepsilon)^n P(X > 1 + \varepsilon)$$

Send $n \rightarrow \infty$, we get $P(X > 1 + \varepsilon) = 0$. By continuity of measure,

$$P(X > 1) = \lim_{k \rightarrow \infty} P(X > 1 + \frac{1}{k}) = 0.$$

Furthermore,

$$M \geq E(X) \geq \int_{\{X=1\}} d\mathbb{P} = P(X = 1).$$

2. Note that for $0 < X < 1$, $X^n \downarrow 0$. By M.C.T,

$$\lim_{n \rightarrow \infty} \int_{\{0 < X < 1\}} X^n d\mathbb{P} = \int_{\{0 < X < 1\}} \lim_{n \rightarrow \infty} X^n d\mathbb{P} = 0$$

Follow the same process as 1, for a fixed $\varepsilon > 0$ and any $n, m \in \mathbb{N}$, we have

$$M \geq E(X^{n+m}) \geq (1 + \varepsilon)^m P(X^n > 1 + \varepsilon)$$

By choosing sufficiently large m , we obtain

$$P(X^n > 1) = 0 \quad \text{for any } n.$$

Thus

$$\lim_{n \rightarrow \infty} E(X^n) = \lim_{n \rightarrow \infty} \left(\int_{\{0 < X < 1\}} X^n d\mathbb{P} + \int_{\{1=X\}} X^n d\mathbb{P} + \int_{\{1 < X\}} X^n d\mathbb{P} \right) = P(X^n = 1).$$

3. It suffices to show that $P(0 < X < 1) = 0$. We claim that $P(\varepsilon < X < 1 - \varepsilon) = 0$ for any $\varepsilon > 0$.

Indeed, note that

$$\begin{aligned} E(X^n) &\geq \int_{\{X=1\}} X^n dP + \int_{\{\varepsilon < X < 1 - \varepsilon\}} X^n dP \\ &\geq P(X = 1) + \varepsilon^n P(\varepsilon < X < 1 - \varepsilon). \end{aligned}$$

Since $E(X^n)$ is a constant, but by 2, $\lim_{n \rightarrow \infty} E(X^n) = P(X = 1)$ and thus $E(X^n) = P(X = 1)$. Hence we conclude $P(\varepsilon < X < 1 - \varepsilon) = 0$. By continuity of measure,

$$P(0 < X < 1) = \lim_{k \rightarrow \infty} P(\frac{1}{k} < X < 1 - \frac{1}{k}) = 0.$$

□

4.6 Inequalities

4.6.1 Chebychev

We use the shorthand

$$\{f \geq \lambda\} := \{x \in E : f(x) \geq \lambda\}.$$

Since

$$\lambda \mathbf{1}_{\{f \geq \lambda\}} \leq f,$$

it follows that

$$\lambda \mu(\{f \geq \lambda\}) \leq \mu(f)$$

which is known as Chebychev's inequality.

Example 4.27 Given a random variable X with finite expectation and variance. Then for every real number t and integer k ,

$$\mathbb{P}(|X| > t) \leq \frac{\mathbb{E}(|X|^k)}{t^k}.$$

Proof. Note that

$$EX^k = \int X^k dP \geq \int_{\{|X|>t\}} X^k dP \geq t^k \int_{\{|X|>t\}} dP = t^k P(|X| > t).$$

□

Example 4.28 Given a random variable X with finite expectation and variance. Then for every real number a ,

$$\mathbb{P}(|X - \mathbb{E}(X)| \geq a) \leq \frac{\text{var}(X)}{a^2}.$$

Proof. Note that

$$\mathbb{P}(|X - \mathbb{E}(X)| \geq a) = \mathbb{P}((X - \mathbb{E}(X))^2 \geq a^2) \leq \frac{\mathbb{E}((X - \mathbb{E}(X))^2)}{a^2} = \frac{\text{var}(X)}{a^2}.$$

□

Example 4.29 If $E(e^{tX}) < \infty$ for any $t \in \mathbb{R}$. Then $P(X > a) = P(e^{tX} > e^{ta}) \leq e^{-at} E(e^{tX})$.

Exercise 4.16 Let X_1, \dots, X_n are independent variables such that $X_i \sim \text{Bernoulli}(p)$. Let $X = \sum_{i=1}^n X_i$, $\mu = np$. Show that for any $\delta > 0$,

$$P(X \geq (1 + \delta)\mu) \leq \left(\frac{e^\delta}{(1 + \delta)^{1+\delta}} \right)^\mu,$$

$$P(X \leq (1 - \delta)\mu) \leq \left(\frac{e^{-\delta}}{(1 - \delta)^{1-\delta}} \right)^\mu.$$

Proof. By Example 4.29, for any $t \in \mathbb{R}$,

$$P(X \geq (1 + \delta)\mu) = P(e^{tX} \geq e^{t(1+\delta)\mu}) \leq e^{-t(1+\delta)\mu} E(e^{tX})$$

and

$$\begin{aligned} E(e^{tX}) &= E[e^{tX_1} \dots e^{tX_n}] = [Ee^{tX_1}]^n \\ &= (1 - p + e^t \cdot p)^n \leq e^{p(e^t - 1) \cdot n} \\ &= e^{\mu(e^t - 1)} \end{aligned}$$

Hence

$$P(X \geq (1 + \delta)\mu) \leq \frac{e^{\mu(e^t - 1)}}{e^{\mu(1 + \delta)t}}.$$

Note that to minimize $\mu(e^t - 1 - t(1 + \delta))$, the optimized t is $t = \ln(1 + \delta)$. \square

Corollary 4.1

In the same setting,

$$P(|X - \mu| \geq \delta\mu) \leq e^{-\frac{\delta^2}{3}\mu}.$$



4.6.2 Cauchy-Schwarz

Theorem 4.5

Let X, Y be random variables and then

$$\mathbb{E}^2[XY] \leq \mathbb{E}(X^2)\mathbb{E}(Y^2). \quad (4.8)$$



Proof. Note that

$$\mathbb{E}[(aX - bY)^2] = a^2\mathbb{E}(X^2) - 2ab\mathbb{E}(XY) + b^2\mathbb{E}(Y^2) \geq 0, \quad \forall a, b$$

Thus its discriminant $4(\mathbb{E}^2[XY] - \mathbb{E}^2(X)\mathbb{E}^2(Y)) \leq 0$. \square

Remark The equality holds if and only if $X = cY$ for some $c \in \mathbb{R}$.

Example 4.30 Let $Y \geq 0$ with $E(Y^2) < \infty$. Apply the Cauchy-Schwarz inequality to $Y\mathbb{1}_{\{Y>0\}}$ and conclude

$$P(Y > 0) \geq (EY)^2/E(Y^2).$$

Solution. Following the hint and by Cauchy-Schwarz,

$$\begin{aligned} E(Y^2)E(\mathbb{1}_{\{Y>0\}}^2) &= E(Y^2)E(\mathbb{1}_{\{Y>0\}}) = E(Y^2)P(Y > 0) \\ &\geq E^2(Y\mathbb{1}_{\{Y>0\}}) \stackrel{Y \geq 0}{=} E^2(Y). \end{aligned}$$

\square

Corollary 4.2

$$|\text{Cov}(X, Y)| \leq \sqrt{\text{var}(X)\text{var}(Y)}. \quad (4.9)$$



Proof. Note that

$$\begin{aligned} \text{Cov}(X, Y) &= \mathbb{E}(X - \mathbb{E}(X))\mathbb{E}(Y - \mathbb{E}(Y)) \\ &\leq \sqrt{\mathbb{E}[(X - \mathbb{E}(X))^2]\mathbb{E}[(Y - \mathbb{E}(Y))^2]} \end{aligned}$$

\square

Example 4.31 Let $X \geq 0$ be a random variable with $\mathbb{E}(X) = 1$. Show that for a fixed $t \in (0, 1)$, $\mathbb{P}(X \geq t) \geq \frac{(1 - t)^2}{\mathbb{E}(X^2)}$.

Proof. Define $Y := \mathbb{1}_{\{X \geq t\}}$. Note that $\mathbb{E}(Y^2) = \mathbb{E}(\mathbb{1}_{\{X \geq t\}}^2) = \mathbb{E}(\mathbb{1}_{\{X \geq t\}}) = \mathbb{P}(\{X \geq t\})$. Then

$$\begin{aligned} \mathbb{P}(X \geq t) \cdot \mathbb{E}(X^2) &= \mathbb{E}(Y^2) \cdot \mathbb{E}(X^2) \geq (\mathbb{E}[XY])^2 = \mathbb{E}^2(X\mathbb{1}_{\{X \geq t\}}) \\ &= (\mathbb{E}X - \mathbb{E}(X\mathbb{1}_{\{X < t\}}))^2 \geq (1 - t)^2. \end{aligned}$$

\square

Corollary 4.3 (Payley-Zygmund)

Let $Z \geq 0$ be a random variable and $t \in (0, 1)$. Then

$$\mathbb{P}[Z \geq t\mathbb{E}(Z)] \geq (1-t)^2 \frac{\mathbb{E}^2(Z)}{\mathbb{E}(Z^2)}. \quad (4.10)$$



Chapter 5 Multi-dimensional random variables

5.1 Joint distribution function

The joint distribution function of random variables X and Y are defined to be

$$F : \mathbb{R}^2 \longrightarrow [0, 1]$$

$$(x, y) \longmapsto P(X \leq x, Y \leq y).$$

Definition 5.1

If X and Y are absolutely continuous random variables, then the joint density function $f : \mathbb{R}^2 \rightarrow [0, +\infty)$ satisfies

$$F(x, y) = \int_{-\infty}^x \int_{-\infty}^y f(u, v) du dv$$

and thus

$$f(x, y) = \frac{\partial^2}{\partial x \partial y} F(x, y).$$

Moreover,

$$P(X \in [a, b], Y \in [c, d]) = \int_a^b \int_c^d f(u, v) du dv$$

$$= F(b, d) - F(b, c) - F(a, d) + F(a, c).$$



By the extension theorem, for $A, B \in \mathcal{B}(\mathbb{R})$,

$$P(X \in A, Y \in B) = \int_A \int_B f(u, v) du dv.$$

Note that the event $\{Y < \infty\}$ and $\{X < \infty\}$ are of probability one, thus

$$\lim_{y \rightarrow \infty} F(x, y) = P(X \leq x, Y < \infty) = P(X \leq x);$$

$$\lim_{x \rightarrow \infty} F(x, y) = P(X < \infty, Y < y) = P(Y \leq y),$$

that is

$$F_X(x) = F(x, \infty),$$

$$F_Y(y) = F(\infty, y).$$

Then from the definition of joint distribution, for discrete cases,

$$P(X = x) = \sum_y P(X = x, Y = y);$$

$$P(Y = y) = \sum_x P(X = x, Y = y);$$

whereas for continuous cases,

$$F_X(x) = F(x, \infty) = \int_{-\infty}^x \left(\int_{-\infty}^{\infty} f(u, v) dv \right) du = \int_{-\infty}^x f_X(u) du;$$

$$F_Y(y) = F(\infty, y) = \int_{-\infty}^y \left(\int_{-\infty}^{\infty} f(u, v) du \right) dv = \int_{-\infty}^y f_Y(v) dv;$$

and

$$f_X(x) = \int_{-\infty}^{\infty} f(x, y) dy;$$

$$f_Y(y) = \int_{-\infty}^{\infty} f(x, y) dx.$$

These results are called the marginal density functions.

Remark X and Y are independent $\iff F(x, y) = F_X(x) \cdot F_Y(y) \iff f(x, y) = f_X(x) \cdot f_Y(y)$.

Example 5.1 If X, Y have joint mass function

$$f(x, y) = \frac{\alpha^x \beta^y}{x! y!} e^{-\alpha - \beta}, \quad x, y \in \mathbb{N},$$

then X and Y are independent:


$$f_X(x) = \sum_{y \in \mathbb{N}} f(x, y) = \frac{\alpha^x}{x!} e^{-\alpha} \sum_{y \in \mathbb{N}} \frac{\beta^y}{y!} e^{-\beta} = \frac{\alpha^x}{x!} e^{-\alpha},$$

implying $X \sim \text{Poisson}(\alpha)$ and $Y \sim \text{Poisson}(\beta)$.

Definition 5.2

For a two-dimensional random variable (X, Y) , the **covariance** between X and Y is defined to be

$$\text{Cov}(X, Y) = E[(X - E(X))(Y - E(Y))]. \quad (5.1)$$

Especially, $\text{Cov}(X, X) = \text{Var}(X)$. 

Definition 5.3

We say X and Y are **uncorrelated** if $\text{Cov}(X, Y) = 0$. 

From these definitions, we know

- X and Y are positively-related if $\text{Cov}(X, Y) > 0$;
- X and Y are negatively-related if $\text{Cov}(X, Y) < 0$;
- X and Y are uncorrelated (distinguish this with independent!) if $\text{Cov}(X, Y) = 0$.

Property $\text{Cov}(X, Y) = E(XY) - E(X)E(Y)$.

Proof. Note that

$$\begin{aligned} \text{Cov}(X, Y) &= E[XY - XE(Y) - YE(X) + E(X)E(Y)] \\ &= E(XY) - E(X)E(Y). \end{aligned}$$

□

Indeed, “uncorrelated” is weaker than “independence”:

Property X and Y are independent $\implies \text{Cov}(X, Y) = 0$ whereas the inverse is not true.

Example 5.2 Let $X \sim \mathcal{N}(0, \sigma^2)$ and $Y = X^2$. Then X and Y are dependent but

$$\text{Cov}(X, Y) = \text{Cov}(X, X^2) = E(X \cdot X^2) - E(X)E(X^2) = 0$$

The equality holds since $E(X^{2m-1}) = 0, m = 1, 2, \dots$ (we will verify this later) provided $X \sim \mathcal{N}(0, \sigma^2)$.

Property


$$\text{var}(X \pm Y) = \text{var}(X) + \text{var}(Y) \pm 2 \text{Cov}(X, Y). \quad (5.2)$$

Proof. Note that

$$\text{var}(X \pm Y) = E[(X \pm Y) - E(X \pm Y)]^2$$

$$\begin{aligned}
&= E\{[X - E(X)] \pm [Y - E(Y)]\}^2 \\
&= E\{[X - E(X)]^2 + [Y - E(Y)]^2 \pm 2[X - E(X)][Y - E(Y)]\} \\
&= \text{Var}(X) + \text{Var}(Y) \pm 2 \text{Cov}(X, Y).
\end{aligned}$$

□

 **Note** Given X and Y being uncorrelated, we have

$$\text{var}(X \pm Y) = \text{var}(X) + \text{var}(Y).$$

A generalization is: for n random variables X_1, X_2, \dots, X_n , one has

$$\text{var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{var}(X_i) + 2 \sum_{i=1}^n \sum_{j=1}^{i-1} \text{Cov}(X_i, X_j). \quad (5.3)$$

Property $\text{Cov}(X, Y) = \text{Cov}(Y, X)$.

Property $\text{Cov}(X, a) = 0$ for any constant a .

Property $\text{Cov}(aX, bY) = ab \text{Cov}(X, Y)$ for constants a, b .

Property $\text{Cov}(X + Y, Z) = \text{Cov}(X, Z) + \text{Cov}(Y, Z)$.

Proof. This is because

$$\begin{aligned}
\text{Cov}(X + Y, Z) &= E[(X + Y)Z] - E(X + Y)E(Z) \\
&= E(XZ) + E(YZ) - E(X)E(Z) - E(Y)E(Z) \\
&= [E(XZ) - E(X)E(Z)] + [E(YZ) - E(Y)E(Z)] \\
&= \text{Cov}(X, Z) + \text{Cov}(Y, Z).
\end{aligned}$$

□

Definition 5.4

For an n -dimensional r.v $\mathbf{X} = (X_1, X_2, \dots, X_n)$, its expectation is given by

$$E(\mathbf{X}) = (E(X_1), E(X_2), \dots, E(X_n)).$$

Its covariance matrix $\text{cov}(\mathbf{X})$ is given by

$$\begin{aligned}
&E[(\mathbf{X} - E(\mathbf{X}))(\mathbf{X} - E(\mathbf{X}))'] \\
&= \begin{pmatrix} \text{var}(X_1) & \text{cov}(X_1, X_2) & \cdots & \text{cov}(X_1, X_n) \\ \text{cov}(X_1, X_2) & \text{var}(X_2) & \cdots & \text{cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{cov}(X_n, X_1) & \text{cov}(X_n, X_2) & \cdots & \text{var}(X_n) \end{pmatrix}
\end{aligned}$$



Theorem 5.1

$\text{cov}(\mathbf{X})$ is symmetrically positive semi-definite.



Proof. Symmetricity is trivial since $\text{cov}(X_i, X_j) = \text{cov}(X_j, X_i)$. Now pick any real vector $\mathbf{c} = (c_1, c_2, \dots, c_n)'$, note that

$$\mathbf{c}' \text{cov}(\mathbf{X}) \mathbf{c} = (c_1, c_2, \dots, c_n) \begin{pmatrix} \text{var}(X_1) & \text{cov}(X_1, X_2) & \cdots & \text{cov}(X_1, X_n) \\ \text{cov}(X_1, X_2) & \text{var}(X_2) & \cdots & \text{cov}(X_2, X_n) \\ \vdots & \vdots & \ddots & \vdots \\ \text{cov}(X_n, X_1) & \text{cov}(X_n, X_2) & \cdots & \text{var}(X_n) \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ \cdots \\ c_n \end{pmatrix}$$

$$\begin{aligned}
&= \sum_{i=1}^n \sum_{j=1}^n c_i c_j \operatorname{cov}(X_i, X_j) \\
&= \sum_{i=1}^n \sum_{j=1}^n E \{ [c_i (X_i - E(X_i))] [c_j (X_j - E(X_j))] \} \\
&= E \left\{ \sum_{i=1}^n \sum_{j=1}^n [c_i (X_i - E(X_i))] [c_j (X_j - E(X_j))] \right\} \\
&= E \left\{ \left[\sum_{i=1}^n c_i (X_i - E(X_i)) \right] \left[\sum_{j=1}^n c_j (X_j - E(X_j)) \right] \right\} \\
&= E \left[\sum_{i=1}^n c_i (X_i - E(X_i)) \right]^2 \geq 0.
\end{aligned}$$

□

Definition 5.5

The *correlation coefficient* of X and Y is defined to be

$$\operatorname{Corr}(X, Y) = \frac{\operatorname{Cov}(X, Y)}{\sqrt{\operatorname{Var}(X)} \sqrt{\operatorname{Var}(Y)}}. \quad (5.4)$$



Remark Let X and Y have expectation μ_X, μ_Y respectively, considering the normalization:

$$X^* = \frac{X - \mu_X}{\sigma_X}, \quad Y^* = \frac{Y - \mu_Y}{\sigma_Y},$$

one has

$$\operatorname{Corr}(X^*, Y^*) = \operatorname{Cov}\left(\frac{X - \mu_X}{\sigma_X}, \frac{Y - \mu_Y}{\sigma_Y}\right) = \frac{\operatorname{Cov}(X, Y)}{\sigma_X \sigma_Y} = \operatorname{Corr}(X, Y).$$

Example 5.3 (Multi-dimensional uniform distribution) Let D be a bounded region in \mathbb{R}^n with volume (area) S_D . If the random variable (X_1, X_2, \dots, X_n) has a joint density function

$$f(x_1, x_2, \dots, x_n) = \begin{cases} \frac{1}{S_D}, & (x_1, x_2, \dots, x_n) \in D, \\ 0, & \text{otherwise,} \end{cases} \quad (5.5)$$

it is called (multi-dimensionally) **uniformly-distributed**.

Now for a sub-region $G \subset D$, one has

$$P((X, Y) \in G) = \iint_G f(x, y) dx dy = \iint_G \frac{1}{S_D} dx dy = \frac{S_G}{S_D}.$$

Example 5.4 (Multi-dimensional normal distribution) Let an n -dimensional r.v. vector $\mathbf{X} = (X_1, X_2, \dots, X_n)$ have covariance matrix $\mathbf{V} = \operatorname{cov}(X)$ and expectation vector $\mathbf{a} = (a_1, a_2, \dots, a_n)'$. Then the distribution with density function

$$f(x_1, x_2, \dots, x_n) = f(\mathbf{x}) = \frac{1}{\sqrt{(2\pi)^n \det \mathbf{V}}} \exp \left\{ -\frac{1}{2} (\mathbf{x} - \mathbf{a})' \mathbf{V}^{-1} (\mathbf{x} - \mathbf{a}) \right\}$$

is called a n -dimensional normal distribution. If we set $\mathbf{V}^{-1} = (r_{ij})$, then we also have

$$f(x_1, x_2, \dots, x_n) = \frac{1}{\sqrt{(2\pi)^n \det \mathbf{V}}} \exp \left\{ -\frac{1}{2} \sum_{i,j=1}^n r_{ij} (x_i - a_i) (x_j - a_j) \right\}.$$

Example 5.5 (Buffon's needle) A bunch of parallel straight lines are placed on the ground with a gap of length 1 and one throws a needle with length 1 on the ground. Try to compute P (the needle intersects with some lines).

Solution. We use Z to denote the distance from the midpoint of the needle to the nearest line whereas Θ to denote the angle formed by the needle and the lines. Then $Z \sim \text{unif}[0, 1/2]$ and $\Theta \sim \text{unif}[0, \pi]$. The joint density function of (Z, Θ) is

$$f(z, \theta) = \begin{cases} \frac{2}{\pi}, & (z, \theta) \in [0, \frac{1}{2}] \times [0, \pi] \\ 0, & \text{otherwise} \end{cases}$$


Note that the event “intersection” is given by

$$B := \{(z, \theta) : 0 < z < \frac{1}{2} \sin \theta\}$$

and thus

$$P(\text{Intersecton}) = \iint_B f(z, \theta) dz d\theta = \frac{2}{\pi} \int_0^\pi d\theta \int_0^{\frac{1}{2} \sin \theta} dz = \frac{2}{\pi} \int_0^\pi \frac{1}{2} \sin \theta d\theta = \frac{2}{\pi}.$$

□



Example 5.6(Standard bivariate normal distribution) Two random variables X and Y are said to have the **standard bivariate normal distribution** with correlation coefficient ρ (verified in later sections) if their joint probability density function is given by

$$f_{XY}(x, y) = \frac{1}{2\pi\sqrt{1-\rho^2}} e^{-\frac{x^2-2\rho xy+y^2}{2(1-\rho^2)}}, \quad -1 < \rho < 1.$$

Then

1. $\iint f(x, y) dx dy = 1$;
2. For $\rho = 0$, $f(x, y) = \frac{1}{2\pi} e^{-\frac{x^2+y^2}{2}}$, and X and Y are independent normally zero-one distributed;
3. The covariance of X and Y satisfies:

$$\text{Cov}(X, Y) = E(XY) - E(X)E(Y) = \iint xy \cdot f(x, y) dx dy = \rho.$$

Proof. Only verify 3. Indeed,

$$\begin{aligned} \text{LHS} &= \iint xy \cdot \frac{1}{\sqrt{2\pi}\sqrt{1-\rho^2}} e^{-\frac{(x-\rho y)^2}{2(1-\rho^2)}} \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} dx dy \\ &= \int y \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} dy \int x \frac{1}{\sqrt{2\pi}\sqrt{1-\rho^2}} e^{-\frac{(x-\rho y)^2}{2(1-\rho^2)}} dx \\ &= \int y \rho y \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} dy \\ &= \rho \int y^2 \frac{1}{\sqrt{2\pi}} e^{-\frac{y^2}{2}} dy \end{aligned}$$

$$= \rho.$$

□

Example 5.7 (General bivariate normal distribution) Two random variables X and Y are said to have the **general bivariate normal distribution** if their joint probability density function is given by

$$f_{XY}(x, y) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} e^{-\frac{1}{2}Q(x,y)},$$

where

$$Q(x, y) = \frac{1}{1-\rho^2} \left[\left(\frac{x-\mu_1}{\sigma_1} \right)^2 - 2\rho \frac{x-\mu_1}{\sigma_1} \frac{y-\mu_2}{\sigma_2} + \left(\frac{y-\mu_2}{\sigma_2} \right)^2 \right].$$

Then

1. Their marginal density: $X \sim \mathcal{N}(\mu_1, \sigma_1^2)$ and $Y \sim \mathcal{N}(\mu_2, \sigma_2^2)$;
- 2.

$$\rho(X, Y) = \frac{\text{Cov}(X, Y)}{\sqrt{\text{var}(X)\text{var}(Y)}} = \rho \quad (\rho = 0 \iff X, Y \text{ are independent})$$

Remark

1. If X and Y are Gaussian, then X and Y are independent $\iff X$ and Y are uncorrelated.
2. For $X \sim \mathcal{N}(0, 1)$, we know $E(e^{tX}) = e^{\frac{1}{2}t^2} = \sum_{k=0}^{\infty} \frac{1}{k!} \frac{t^{2k}}{2^k}$. Note that

$$E(e^{tX}) = E\left(\sum_{n=0}^{\infty} \frac{t^n X^n}{n!}\right) = \sum_{n=1}^{\infty} \frac{1}{n!} t^n E(X^n).$$

Observing the coefficients:

$$E(X^{2k+1}) = 0, \quad E(X^{2k}) = (2k-1)!! = (2k-1)(2k-3)\cdots 1 = \text{number of pairs of } \{1, 2, \dots, 2k\}.$$

Theorem 5.2 (Wick)

Let X_1, X_2, \dots, X_n be normal distributed r.v.s such that $E(X_i) = 0$. Then

$$E[X_1 X_2 \cdots X_n] = \sum_{\text{pairings } \pi \text{ of } \{1, 2, \dots, 2n\}} \prod_{(i,j) \in \pi} E[X_i X_j].$$



Example 5.8 Follow Wick's Theorem, $E(X_1 X_2 X_3 X_4) = E(X_1 X_2)E(X_3 X_4) + E(X_1 X_3)E(X_2 X_4) + E(X_1 X_4)E(X_2 X_3)$.

Exercise 5.1 Randomly pick two numbers from $0, 1, \dots, n$. Try to compute the expectation of the absolute value of their difference.

Solution. We use X and Y to denote the first number and the second number, then


$$P(X = i, Y = j) = \frac{1}{(n+1)n}, \quad i, j = 0, 1, \dots, n, \quad i \neq j.$$

Therefore,

$$\begin{aligned} E(|X - Y|) &= \frac{1}{(n+1)n} \sum_{i=0}^n \left\{ \sum_{j=0}^i (i-j) + \sum_{j=i+1}^n (j-i) \right\} \\ &= \frac{1}{(n+1)n} \sum_{i=0}^n \left\{ \frac{i(i+1)}{2} + \frac{(n-i)([n-i]+1)}{2} \right\} \\ &= \frac{1}{(n+1)n} \sum_{i=0}^n \left\{ i^2 + \frac{n(n+1)}{2} - in \right\} \end{aligned}$$

$$= \frac{2n+1}{6} + \frac{n+1}{2} - \frac{n}{2} = \frac{n+2}{3}.$$

□

 **Exercise 5.2** Randomly pick n points from the interval $(0, 1)$. What is the expectation of their farthest distance?

Solution. n points will separate the interval $(0, 1)$ into $n + 1$ segments. We use Y_1, Y_2, \dots, Y_{n+1} to denote their length. Then Y_1, Y_2, \dots, Y_{n+1} are i.i.d., with the same expectation. That is

$$E(Y_1) = E(Y_2) = \dots = E(Y_{n+1}) = \frac{1}{n+1}.$$

Note that the farthest distance is precisely $Y_2 + Y_3 + \dots + Y_n$, thus the result is $\frac{n-1}{n+1}$. □

5.2 Conditional distribution

5.2.1 Discrete cases

Let X and Y be two discrete random variables.

Definition 5.6

The conditional distribution function of Y given $X = x$ is

$$F_{Y|X}(y | x) = P(Y \leq y | X = x) = \sum_{y_i \leq y} P(Y = y_i | X = x), \quad (5.6)$$

for every $x \in \mathbb{R}$, such that $P(X = x) > 0$. The conditional mass function is

$$f_{Y|X}(y | x) = P(Y = y | X = x). \quad (5.7) \quad \clubsuit$$

Remark

1. $f(x, y) = f_X(x) \cdot f_{Y|X}(y | x)$;
2. if X and Y are independent, then $f_{Y|X}(y | x) = f_Y(y)$ for all x .

Example 5.9 Let X and Y be independent r.v.s such that $X \sim \text{Poisson}(\lambda_1)$, $Y \sim \text{Poisson}(\lambda_2)$. Find the conditional distribution of X given $X + Y = n$.

Solution. Note that $X + Y \sim \text{Poisson}(\lambda_1 + \lambda_2)$ and thus

$$\begin{aligned} P(X = k | X + Y = n) &= \frac{P(X = k, X + Y = n)}{P(X + Y = n)} \\ &= \frac{P(X = k)P(Y = n - k)}{P(X + Y = n)} \\ &= \frac{\frac{\lambda_1^k}{k!} e^{-\lambda_1} \cdot \frac{\lambda_2^{n-k}}{(n-k)!} e^{-\lambda_2}}{\frac{(\lambda_1 + \lambda_2)^n}{n!} e^{-(\lambda_1 + \lambda_2)}} \\ &= \frac{n!}{k!(n-k)!} \frac{\lambda_1^k \lambda_2^{n-k}}{(\lambda_1 + \lambda_2)^n} \\ &= \binom{n}{k} \left(\frac{\lambda_1}{\lambda_1 + \lambda_2} \right)^k \left(\frac{\lambda_2}{\lambda_1 + \lambda_2} \right)^{n-k}, \quad k = 0, 1, \dots, n. \end{aligned}$$

That is, given $X + Y = n$, one has $X \sim \text{Binomial}(n, p)$, where $p = \lambda_1 / (\lambda_1 + \lambda_2)$. □

5.2.2 Continuous cases

Let X and Y be continuous r.v.s. Like previously, set $\Delta x \rightarrow 0$, considering

$$\begin{aligned} P[Y \leq y \mid x \leq X \leq x + \Delta x] &= \frac{P(Y \leq y, x \leq X \leq x + \Delta x)\Delta x}{P(x \leq X \leq x + \Delta x)\Delta x} \\ &= \frac{\int_{-\infty}^y \int_x^{x+\Delta x} f(u, v) du dv}{f_X(x)\Delta x} \\ &\approx \frac{\int_{-\infty}^y f(x, v) dv \Delta x}{f_X(x)\Delta x} \end{aligned}$$

and we define

Definition 5.7

The conditional distribution function of Y given $X = x$ is

$$F_{Y|X}(y \mid x) = \int_{-\infty}^y \frac{f(x, v)}{f_X(x)} dv, \quad (5.8)$$

for every $x \in \mathbb{R}$, such that $P(X = x) > 0$. The conditional mass function is

$$f_{Y|X}(y \mid x) = \frac{f(x, y)}{f_X(x)}. \quad (5.9)$$



Example 5.10 Suppose the joint density function of X, Y is

$$f(x, y) = \begin{cases} \frac{1}{y} e^{-x/y} e^{-y}, & x, y > 0, \\ 0, & \text{otherwise.} \end{cases}$$

Try to compute $P(X > 1 \mid Y = y)$.

Solution.

$$\begin{aligned} f_{X|Y}(x, y) &= \frac{f(x, y)}{f_Y(y)} = \frac{f(x, y)}{\int_0^{\infty} f(x, y) dx} \\ &= \frac{e^{-x/y} e^{-y}/y}{\frac{e^{-y}}{y} \int_0^{\infty} e^{-x/y} dx} = \frac{e^{-x/y} e^{-y}/y}{e^{-y}} \\ &= \frac{1}{y} e^{-x/y}. \end{aligned}$$

Thus

$$P(X > 1 \mid Y = y) = \int_1^{\infty} \frac{1}{y} e^{-x/y} dx = -e^{-x/y} \Big|_1^{\infty} = e^{-1/y}.$$

□

Exercise 5.3 Given compound random variables $X(U, V), Y(U, V)$ and their joint density function $f_{X,Y}$, try to compute the joint density function of U, V .

Solution. Let $R := (-\infty, x] \times (-\infty, y]$. Let T be the C^1 transformation whose Jacobian is nonzero and that maps region S in the uv -plane onto a region R in the xy -plane. Then

$$\iint_R f_{X,Y}(x, y) dx dy = \iint_S f_{X,Y}(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$$

and thus

$$f_{U,V}(u, v) = f_{X,Y}(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right|.$$

□

Exercise 5.4 Let $X \sim \text{Unif}(1, 2)$ and $Y \sim \exp(x)$ given $X = x$. Try to compute the joint distribution function of XY .

Solution. By condition, $Y | X = x \sim \text{Exp}(x)$ and thus

$$f(x, y) = f_X(x)f(y | x) = xe^{-xy}, \quad 1 < x < 2, y > 0.$$

Consider the transformation $\begin{cases} x = v, \\ y = \frac{u}{v}, \end{cases}$ with Jacobian

$$J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} 0 & 1 \\ \frac{1}{v} & -\frac{u}{v^2} \end{vmatrix} = -\frac{1}{v}$$

Note that we also have $\begin{cases} u = xy, \\ v = x, \end{cases}$ and thus

$$f_{U,V}(u, v) = f_{X,Y}\left(v, \frac{u}{v}\right) \left| -\frac{1}{v} \right| = ve^{-vu/v} \frac{1}{v} = e^{-u}, \quad 1 < v < 2, u > 0.$$

Then by marginal density,

$$f_U(u) = \int_1^2 e^{-u} dv = e^{-u}, \quad u > 0.$$

□

Exercise 5.5 Show that the correlation coefficient of the bivariate normal distribution $\mathcal{N}(\mu_1, \mu_2, \sigma_1^2, \sigma_2^2, \rho)$ is precisely ρ .

Proof. Start with $\text{Cov}(X, Y)$.

$$\begin{aligned} \text{Cov}(X, Y) &= E[(X - E(X))(Y - E(Y))] \\ &= \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x - \mu_1)(y - \mu_2) \cdot \\ &\quad \exp \left\{ -\frac{1}{2(1-\rho^2)} \left[\frac{(x - \mu_1)^2}{\sigma_1^2} - 2\rho \frac{(x - \mu_1)(y - \mu_2)}{\sigma_1\sigma_2} + \frac{(y - \mu_2)^2}{\sigma_2^2} \right] \right\} dx dy. \end{aligned}$$

The terms inside the square brackets are exactly

$$\left(\frac{x - \mu_1}{\sigma_1} - \rho \frac{y - \mu_2}{\sigma_2} \right)^2 + \left(\sqrt{1 - \rho^2} \frac{y - \mu_2}{\sigma_2} \right)^2,$$

Consider the transformation

$$\begin{cases} u = \frac{1}{\sqrt{1-\rho^2}} \left(\frac{x - \mu_1}{\sigma_1} - \rho \frac{y - \mu_2}{\sigma_2} \right), \\ v = \frac{y - \mu_2}{\sigma_2}, \end{cases}$$

then

$$\begin{cases} x - \mu_1 = \sigma_1 (u\sqrt{1 - \rho^2} + \rho v) \\ y - \mu_2 = \sigma_2 v \end{cases}$$

$$dx dy = |J| du dv = \sigma_1 \sigma_2 \sqrt{1 - \rho^2} du dv.$$

i.e.,

$$\text{Cov}(X, Y) = \frac{\sigma_1 \sigma_2}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (uv\sqrt{1 - \rho^2} + \rho v^2) \exp\left\{-\frac{1}{2}(u^2 + v^2)\right\} du dv.$$

Note that

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} uv \exp\left\{-\frac{1}{2}(u^2 + v^2)\right\} du dv = 0$$

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} v^2 \exp\left\{-\frac{1}{2}(u^2 + v^2)\right\} du dv = 2\pi$$

and thus

$$\text{Cov}(X, Y) = \frac{\sigma_1 \sigma_2}{2\pi} \cdot \rho \cdot 2\pi = \rho \sigma_1 \sigma_2$$

$$\text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\sigma_1 \sigma_2} = \rho$$

□

5.2.3 Total probability formula; Bayes formula

Rewrite (5.9) as

$$f(y, x) = f_X(x) f_{Y|X}(y | x), \quad (5.10)$$

$$f(x, y) = f_Y(y) f_{X|Y}(x | y), \quad (5.11)$$

and compute the marginal density, we will get

$$f_Y(y) = \int_{-\infty}^{\infty} f_X(x) f_{Y|X}(y | x) dx, \quad (5.12)$$

$$f_X(x) = \int_{-\infty}^{\infty} f_Y(y) f_{X|Y}(x | y) dy. \quad (5.13)$$

Now substitute (5.11) and (5.13) back to (5.9), obtaining the Bayes:

$$f_{Y|X}(y | x) = \frac{f_Y(y) f_{X|Y}(x | y)}{\int_{-\infty}^{\infty} f_Y(y) f_{X|Y}(x | y) dx}; \quad (5.14)$$

$$f_{X|Y}(x | y) = \frac{f_X(x) f_{Y|X}(y | x)}{\int_{-\infty}^{\infty} f_X(x) f_{Y|X}(y | x) dx}. \quad (5.15)$$

5.3 Conditional expectation

5.3.1 Discrete cases

Let X and Y be discrete r.v.s.

Definition 5.8

For any x , the conditional expectation of Y given $X = x$ is

$$\varphi(x) := E[Y | X = x] = \sum_y y f_{Y|X}(y | x). \quad (5.16)$$

The conditional expectation of Y given X is a function of X :

$$E[Y | X] := \varphi(X) \quad (5.17) \quad \clubsuit$$

Remark $E[Y | X]$ is a random variable denoting the “best guess” of Y given the information of X .

Proposition 5.1 (Towering property)

$$E[E(Y | X)] = E(Y). \quad \spadesuit$$

Proof. Note that

$$\begin{aligned} E[E(Y | X)] &= \sum_x f_X(x) E[Y | X = x] = \sum_x \sum_y f_X(x) \cdot y f_{Y|X}(y | x) \\ &= \sum_x \sum_y y f(x, y) = \sum_y y f_Y(y) \\ &= E(Y). \end{aligned}$$

□

Remark

1. Towering property is useful to compute $E(Y)$;
2. Let $\{A_i\}$ be a partition of Ω , then

$$E(Y) = \sum P(A_i) E(Y | A_i)$$

Example 5.11 For a coffee shop, we assume that N , the number of customer, satisfies $N \sim \text{Poisson}(\lambda)$ and each customer has probability p to carry a dog and probability $1 - p$ of not carrying a dog. Let K denote the number of dogs. Try to compute $E[K | N]$, $E(K)$, $E[N | K]$.

Solution. Note that $E[K | N = n] = np$ since given $N = n$, $K \sim \text{Binomial}(n, p)$. Thus $E[K | N] = pN$, $E(K) = E[E(K | N)] = p \cdot E(N) = p\lambda$. Finally,

$$\begin{aligned} f_{N|K}(n | k) &= \frac{f_{K|N}(k | n) f_N(n)}{f_K(k)} \\ &= \frac{\binom{n}{k} p^k (1-p)^{n-k} \cdot \frac{\lambda^n}{n!} e^{-\lambda}}{\sum_{n \geq k} \binom{n}{k} p^k (1-p)^{n-k} \frac{\lambda^n}{n!} e^{-\lambda}} \\ &= \frac{\frac{p^k \lambda^k}{k!} e^{-p\lambda} \frac{(1-p)^{n-k} \lambda^{n-k}}{n-k!} e^{-(1-p)\lambda}}{\frac{p^k \lambda^k}{k!} e^{-p\lambda} \sum_{n \geq k} \frac{(1-p)^{n-k} \lambda^{n-k}}{n-k!} e^{-(1-p)\lambda}} \\ &= \frac{(1-p)^{n-k}}{(n-k)!} \lambda^{n-k} e^{-(1-p)\lambda}, \end{aligned}$$

implying given $K = k$, $N - k$ is Poisson $((1-p)\lambda)$. Thus

$$E[N | K = k] = E[N - k | K = k] + k = (1-p)\lambda + k, \quad E[N | K] = (1-p)\lambda + K.$$

□

5.3.2 Continuous cases

Definition 5.9

For any x , the conditional expectation of Y given $X = x$ is

$$\varphi(x) := E[Y | X = x] = \int_{-\infty}^{\infty} y f_{Y|X}(y | x) dy. \quad (5.18)$$

The conditional expectation of Y given X is nothing but a random variable, a function of X :

$$E[Y | X] := \varphi(X) \quad (5.19) \quad \clubsuit$$

The following property holds for both discrete and continuous cases:

Proposition 5.2

For X and Y being independent, one has $E[Y | X] = E(Y)$. ♠

Proof. We prove the discrete case. Note that

$$E[Y | X = x_i] = \sum_y y f_{Y|X}(y | x_i) = \sum_y y f_Y(y) = E(Y)$$

for any x_i . □

Lemma 5.1

For X and Y being independent and for any $g : \mathbb{R} \rightarrow \mathbb{R}$, one has $E[g(Y) | X] = g(Y)$. ♥

Again, we have tower property in continuous sense:

Proposition 5.3 (Tower property)

$$E(E(Y | X)) = E(Y). \quad \spadesuit$$

Proof. Note that

$$\begin{aligned} E(Y) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f(x, y) dy dx = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f(y | x) f_X(x) dy dx \\ &= \int_{-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} y f(y | x) dy \right\} f_X(x) dx = \int_{-\infty}^{\infty} E(Y | X = x) f_X(x) dx \\ &= E(E(Y | X)). \end{aligned} \quad \square$$

Remark If $Y = \mathbb{1}_A$, we have

$$P(A) = \int_{-\infty}^{\infty} f_X(x) P(A | X = x) dx.$$

Corollary 5.1

From the proof of tower property, given X discrete, one has

$$E(Y) = \sum_x E(Y | X = x) P(X = x) \quad (5.20)$$

whereas given X continuous, one has

$$E(Y) = \int_{-\infty}^{\infty} E(Y | X = x) f_X(x) dx. \quad (5.21) \quad \heartsuit$$

Example 5.12 Recall the bivariate standard normal distribution:

$$f(x, y) = \frac{1}{2\pi\sqrt{1-\rho^2}} e^{-\frac{1}{2(1-\rho^2)}(x^2 - 2\rho xy + y^2)}.$$

Try to compute $E(Y | X)$.

Solution. Note that

$$f(x, y) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} \cdot \frac{1}{\sqrt{2\pi(1-\rho^2)}} e^{-\frac{(y-\rho x)^2}{2(1-\rho^2)}}$$

Since $f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$, one has $f_{Y|X}(y | x) = \frac{1}{\sqrt{2\pi(1-\rho^2)}} e^{-\frac{(y-\rho x)^2}{2(1-\rho^2)}}$. Then

$$E(Y | X = x) = \rho x, \quad E(Y | X) = \rho X.$$

□

Example 5.13 A man is at a three-way intersection. It will take him 3 hours to reach his destination if he chooses road one. However, he will get back to the origin after 5 and 7 hours if he chooses road two or three, respectively. Suppose that he will choose each road with a equal probability. What will be the average time to reach his destination?

Solution. Let X denote the time he needs. We use random variable Y to denote the road he chooses, i.e., $\{Y = i\}$ denotes the event that he chooses road i . By the condition,

$$P(Y = 1) = P(Y = 2) = P(Y = 3) = \frac{1}{3}.$$

Note that

$$E(X | Y = 1) = 3;$$

$$E(X | Y = 2) = 5 + E(X);$$

$$E(X | Y = 3) = 7 + E(X).$$

From (5.20),

$$E(X) = \frac{1}{3}[3 + 5 + E(X) + 7 + E(X)] = 5 + \frac{2}{3}E(X)$$

and thus $E(X) = 15$.

□

Example 5.14 Let X_1, X_2, \dots be random variables that are independent with N , another random variable only taking values in \mathbb{N} . Show that

$$E\left(\sum_{i=1}^N X_i\right) = E(X_1)E(N).$$

Proof. By Tower property and (5.20),

$$\begin{aligned} E\left(\sum_{i=1}^N X_i\right) &= E\left[E\left(\sum_{i=1}^N X_i \mid N\right)\right] = \sum_{n=1}^{\infty} E\left(\sum_{i=1}^N X_i \mid N = n\right)P(N = n) \\ &= \sum_{n=1}^{\infty} E\left(\sum_{i=1}^n X_i\right)P(N = n) = \sum_{n=1}^{\infty} nE(X_1)P(N = n) \\ &= E(X_1) \sum_{n=1}^{\infty} nP(N = n) = E(X_1)E(N). \end{aligned}$$

□

 **Exercise 5.6** Show that

1. $E(aY + bZ | X) = aE(Y | X) + bE(Z | X)$ for $a, b \in \mathbb{R}$;
2. (tower property) $E\{E(Y | X, Z) | X\} = E(Y | X) = E\{E(Y | X) | X, Z\}$

Proof. 1. We have

$$\begin{aligned} \mathbb{E}(aY + bZ | X = x) &= \sum_{y,z} (ay + bz) \mathbb{P}(Y = y, Z = z | X = x) \\ &= a \sum_{y,z} y \mathbb{P}(Y = y, Z = z | X = x) + b \sum_{y,z} z \mathbb{P}(Y = y, Z = z | X = x) \\ &= a \sum_y y \mathbb{P}(Y = y | X = x) + b \sum_z z \mathbb{P}(Z = z | X = x) \\ &= aE(Y | X = x) + bE(Z | X = x). \end{aligned}$$


2. Define the function g by

$$g(x, z) = E[Y | X = x, Z = z]$$

Then

$$\begin{aligned} E\{E(Y | X, Z) | X = x\} &= \sum_{x',z} g(x', z) P(X = x', Z = z | X = x) \\ &= \sum_z g(x, z) P(X = x | Z = z) \\ &= \sum_z E[Y | X = x, Z = z] P(X = x | Z = z) \\ &= \sum_z \sum_y y P(Y = y | X = x, Z = z) P(X = x | Z = z) \\ &= \sum_z \sum_y y \frac{\mathbb{P}(Y = y, X = x, Z = z)}{\mathbb{P}(X = x, Z = z)} \cdot \frac{\mathbb{P}(X = x, Z = z)}{\mathbb{P}(X = x)} \\ &= \sum_y y \mathbb{P}(Y = y | X = x) = \mathbb{E}(Y | X = x) \\ &= \mathbb{E}\{\mathbb{E}(Y | X) | X = x, Z = z\}. \end{aligned}$$

□


 **Exercise 5.7** If $E(Y)$ and $E[h(Y)]$ both exist, show that $E[h(Y) | Y] = h(Y)$.

Proof. We prove the discrete case. Let $\varphi(Y) = E[h(Y) | Y]$. Note that for any $y_i (i = 1, 2, \dots)$, $h(y_i)$ is a constant and hence is independent with Y . Therefore

$$\varphi(y_i) = E[h(y_i) | Y = y_i] = h(y_i)$$

and one deduces that $E[h(Y) | Y] = h(Y)$.

□

 **Exercise 5.8** Show the following properties:

1. $E[g(X)Y | X] = g(X)E(Y | X)$;
2. $E(XY) = E[XE(Y | X)]$;
3. $\text{Cov}[X, E(Y | X)] = \text{Cov}(X, Y)$.

Proof. 1. Note that $E(g(X)Y | X = x) = g(x)E(Y | X = x)$ and thus $E[g(X)Y | X] = g(X)E(Y | X)$;

2. Note that $E(XY) = E[E(XY | X)] = E[XE(Y | X)]$;

3. Note that

$$\begin{aligned}\text{Cov}[X, E(Y | X)] &= E[X \cdot E(Y | X)] - E(X) \cdot E(Y) \\ &= E(XY) - E(X) \cdot E(Y) \\ &= \text{Cov}(X, Y).\end{aligned}$$

□

For multiple conditioning, we define

Definition 5.10

$$E[X | Y, Z] = \varphi(Y, Z)$$

such that

$$\varphi(y, z) = E[X | Y = y, Z = z] = \begin{cases} \sum x P(X = x | Y = y, Z = z) \\ \int^x x f(x, y, z) dx = f_{YZ}(y, z) \end{cases}$$



Proposition 5.4 (Tower proposition)

Given g measurable,

$$E(E(Y | X) \cdot g(X)) = E(Y \cdot g(X)).$$



Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\mathcal{G} \subseteq \mathcal{F}$ be a sub- σ -algebra.

Definition 5.11

The measure theoretic definition of conditional expectation $E[X | \mathcal{G}]$ is a random variable such that

1. $\varphi_{\mathcal{G}} := E[X | \mathcal{G}]$ is \mathcal{G} -measurable.
2. For all $A \in \mathcal{G}$, we have $\int_A E[X | \mathcal{G}] d\mathbb{P} = \int_A X d\mathbb{P}$.

We define $E[X | Y] = E[X | \sigma(Y)]$.



Example 5.15 Let $\mathcal{G} = \mathcal{P}(\Omega)$, then by 2, $E[X | \mathcal{G}] = X$ a.s..

Example 5.16 Let $\mathcal{G} = \{\emptyset, \Omega\}$, then by 2, $E[X | \mathcal{G}] = E(X)$.

Chapter 6 Random Walk

1-dimensional random walk

$$S_n = S_0 + \sum_{i=1}^n X_i,$$

where X_i are i.i.d random variables. We set $P(X_i = 1) = p$ and $P(X_i = -1) = q$. If $p = q = 1/2$, it is called the symmetric simple random walk.

More generally, many results under the assumption that X_i are i.i.d and $E(X_i^2) < \infty$.

d -dimensional random walk Now we assume $S_n \in \mathbb{Z}^d$.

Lemma 6.1 (Spatial homogeneity)

$$P(S_n = j \mid S_0 = a) = P(S_n = j - a \mid S_0 = 0). \quad (6.1) \quad \heartsuit$$

Proof. Note that

$$P(S_n = j \mid S_0 = a) = P(S_n = j - a \mid S_0 = 0) = P\left(\sum_{i=1}^n X_i = j - a\right).$$

□

Lemma 6.2 (Time homogeneity)

$$P(S_n = j \mid S_0 = a) = P(S_{n+m} = j \mid S_m = a). \quad (6.2) \quad \heartsuit$$

Proof. Note that

$$\text{LHS} = P\left(\sum_{i=1}^n X_i = j - a\right) = P\left(\sum_{i=m+1}^{m+n} X_i = j - a\right) = \text{RHS}.$$

□

Proposition 6.1 (Markov property)

The position after m steps do not depend on the previous m steps:

$$P(S_{n+m} = j \mid S_0, S_1, \dots, S_m) = P(S_{n+m} = j \mid S_m).$$

♠

Recurrence and Transience Let $T_y^0 = 0$ and for $k \geq 1$, let $T_y^k := \inf\{n > T_y^{k-1} : S_n = y\}$.

Definition 6.1

We say that a random walk is **recurrent** if it visits its starting position infinitely often with probability one and **transient** if it visits its starting position finitely often with probability one. That is y is called **recurrent** if $P_y[T_y^1 < \infty] = 1$ and is called **transient** if $P_y[T_y^1 < \infty] < 1$.

♣

Remark

1. If y is recurrent, by Markov property, $\forall n \in \mathbb{N}, P(T_y^k < \infty) = 1$. Thus $P_y(S_n = y \text{ infinitely often}) = P_y(\limsup_{n \rightarrow \infty} \{S_n = y\}) = 1$.

2. If y is transient, let $N_y := \sum_{n=1}^{\infty} \mathbb{1}_{\{S_n=y\}}$ (number of times of visiting y). Then

$$E(N_y) = \sum_k P(N_y \geq k) = \sum_k P_y(T_y^k < \infty) = \sum_{k=1}^{\infty} [P_y(T_y^1 < \infty)]^k = \frac{P_y(T_y^1 < \infty)}{1 - P_y(T_y^1 < \infty)} < \infty.$$

Theorem 6.1

y is a recurrent state (or transient state) if and only if $E(N_y) = +\infty$ (or $E(N_y) < +\infty$).

Proof. Start with 1-D symmetric random walk. We define

$$\delta_y(x) = \begin{cases} 1, & \text{if } y = x \\ 0, & \text{else} \end{cases}$$

Note that

$$\begin{aligned} E(N_y) &= P(X_1 = 1) \cdot E(N_y | X_1 = 1) + P(X_1 = -1) \cdot E(N_y | X_1 = -1) + \delta_y \\ &= \frac{1}{2}E(N_{x-1}) + \frac{1}{2}E(N_{x+1}) + \delta_y \end{aligned}$$

with $E(N_0) = 0$ and $E(N_N) = 0$. Solve the recurrence,

$$E(N_y) \sim cN$$

and thus $E(N_y) \rightarrow +\infty$ as $N \rightarrow +\infty$.

For random walk on \mathbb{Z}^d ($d \geq 2$), let $G(x) := E(N_x)$ denote the mean number of returns at x before hitting the boundary. Then

$$\begin{aligned} G(x) &= \frac{1}{2d} \sum_{z \sim x} G(z) + \delta_y \\ G(z) &= 0, \quad \text{if } z \in \partial \end{aligned}$$

Let $\Delta f(x) = \frac{1}{2d} \sum_{z \sim x} (f(z) - f(x))$. Then the recurrence can be written as the **discrete Green's function**:

$$\begin{aligned} \Delta G(x) &= -\delta_y(x) \\ G(z) &= 0 \quad \text{if } z \in \partial \end{aligned}$$

Lemma 6.3

Set $y = 0$, then

$$G(0) \approx \begin{cases} \ln N, & d = 2, \\ O(1), & d \geq 3, \end{cases}$$

implying y is recurrent in $d = 2$ and transient in $d \geq 3$.

Recall solving PDE by Fourier transformation:

$$\Delta G(x) = -\delta_0(x), \quad \text{in } \mathbb{R}^d,$$

then

$$\begin{aligned} \hat{G}(k) &= \int_{\mathbb{R}^d} e^{ikx} G(x) dx \\ \Delta \hat{G}(k) &= (k_1^2 + k_2^2 + \dots + k_d^2) \hat{G}(k) \end{aligned}$$

$$= \int e^{ikx} \delta_0(x) dx = 1,$$

thus

$$\hat{G}(k) = \frac{1}{|k|^2}.$$

Actually on \mathbb{Z}^d , $f_k(x) := e^{ikx}$ are eigenfunctions of Δ with $\Delta f_k = \lambda_k f_k$, where $\lambda_k = -\sum_{j=1}^d (1 - \cos k_j)$. For example, on $d = 1$,

$$\begin{aligned} \Delta f_k(x) &= \frac{1}{2} f_k(x+1) + \frac{1}{2} f_k(x-1) - f_k(x) \\ &= f_k(x) \left(\frac{1}{2} e^{ik} + \frac{1}{2} e^{-ik} - 1 \right) \\ &= f_k(x) (\cos k - 1) \end{aligned}$$

Fourier transformation

$$\hat{G}(k) = \sum_{x \in \text{Box}_N} G(x) \cdot e^{-ikx}$$

inverse Fourier transformation:

$$G(x) = \frac{1}{|\text{Box}_N|} \sum_{k \in \text{Box}_N^*} e^{-ikx} \hat{G}(k),$$

where

$$\text{Box}_N^* = \left\{ \frac{2\pi}{N} (n_1, \dots, n_d) : n_i \in [-N, N] \cap \mathbb{Z} \right\}.$$

then

$$\Delta \cdot \sum_{k \in \text{Box}_N} G(x) e^{ikx} = \lambda_k \hat{G}(k) = 1.$$

Using inverse:

$$\begin{aligned} G(0) &= \frac{1}{|\text{Box}_N|} \sum_{k \in \text{Box}_N^*} \hat{G}(k) = \frac{1}{|\text{Box}_N|} \sum_{k \in \text{Box}_N^*} \frac{1}{\sum_{j=1}^d (1 - \cos k_j)} \\ &\rightarrow \frac{1}{(2\pi)^d} \int_{[-\pi, \pi]^d} \frac{1}{\sum_{j=1}^d (1 - \cos x_j)} dx, \end{aligned}$$

note that $1 - \cos x_j \sim |x_j|^2$ and thus \square

Example 6.1 Consider the simple random walk on \mathbb{Z} with probability p to go right and probability q to go left. If the walk starts at a , try to compute $P(S_n = b)$.

Solution.

$$P(S_n = b) = \sum_r M_n^r(a, b) p^r q^{n-r}$$

where $M_n^r(a, b) :=$ number of paths with $S_0 = a, S_n = b$ such that makes r right moves. In fact, $r - (n - r) = b - a$ and thus $r = \frac{1}{2}(n + b - a)$, $M_n^r(a, b) = \binom{n}{\frac{1}{2}(n + b - a)}$. \square

Proposition 6.2 (Reflection principle)

If $a, b > 0$, then $N_{0,n}(a, b) = N_n(-a, b)$ where $N_{0,n}(a, b)$ denotes the number of paths starting at a , ending at b , and visiting 0 somehow; $N_n(-a, b)$ denotes the number of paths starting at $-a$, ending at b without extra conditions.

Proof. Let $T = \inf\{n : S_n = 0\}$. □

In a ballot, eventually candidate A has a votes and B has b votes such that $a > b$. Now we revisit the votes one by one, try to compute $P(A \text{ is always ahead of } B \mid a > b)$.

Indeed, $S_i =$ number of votes for A at time i – number of votes for B at time i .

Theorem 6.2

Let S be a simple random walk on \mathbb{Z} with $S_0 = 0$, then

$$P(S_n = b, S_i \neq 0, i = 1, 2, \dots, n) = \frac{|b|}{n} P(S_n = b).$$

Proof. Assume $b > 0$, then the number of paths from $(0, 0)$ to (n, b) that does not visit 0 is:

$$\begin{aligned} N_{n-1}(1, b) - N_{0,n-1}(1, b) &= N_{n-1}(1, b) - N_{n-1}(-1, b) \\ &= \binom{n-1}{\frac{1}{2}(n-1+b-1)} - \binom{n-1}{\frac{1}{2}(n-1+b+1)} \\ &= \binom{n}{\frac{1}{2}(n+b)} \left(\frac{n+b}{2n} - \frac{n-b}{2n} \right) \\ &= \frac{b}{n} \binom{n}{\frac{1}{2}(n+b)} \\ &= \frac{b}{n} N_n(0, b). \end{aligned}$$

Thus $P(S_n = b, S_i \neq 0, i = 1, 2, \dots, n) = \frac{|b|}{n} P(S_n = b)$. □

The Ballot problem is now converted to a simple random walk starting at 0: If we get a vote for A , we go right. Otherwise we go left. What we want is exactly $P(S_{a+b} = a - b, S_i \neq 0, i = 1, 2, \dots, a + b) = \frac{a - b}{a + b}$.

Corollary 6.1

$$P(S_i \neq 0, i = 1, 2, \dots, n) = \frac{1}{n} E(S_n) \leq \frac{1}{n} \sqrt{\text{var}(S_n)} \leq \frac{\sqrt{n}}{n}.$$

Theorem 6.3

Suppose that $S_0 = 0$ with $p = q = \frac{1}{2}$. Then for $a > 0$,

$$P\left(\max_{1 \leq k \leq n} S_k \geq a\right) = P(S_n \geq a) + P(S_n \geq a + 1).$$

Proof. Note that

$$\text{LHS} = P\left(\max_{1 \leq k \leq n} S_k \geq a; S_n \geq a\right) + P\left(\max_{1 \leq k \leq n} S_k \geq a; S_n < a\right)$$

and

$$P\left(\max_{1 \leq k \leq n} S_k \geq a; S_n \geq a\right) = P(S_n \geq a).$$

Let $T_a = \inf\{n \geq 1 : S_n = a\}$. Define the reflected walk

$$\overline{S}_n = \begin{cases} S_n, & \text{if } n \leq T_a, \\ 2a - S_n, & \text{if } n > T_a. \end{cases}$$

Note that $2a - S_n$ has the distribution of SRW starting at a . By Markov property, \overline{S}_n has the same distribution as the SRW, i.e.,

$$P\left(\max_{1 \leq k \leq n} S_k \geq a; S_n < a\right) = P(\overline{S}_n > a) = P(S_n > a) = P(S_n \geq a + 1).$$

□

Remark We say $(W_t)_{t \geq 0}$ is a **Bronian motion** if

1. $W_0 = 0$;
2. For any $t > s > 0$, $W_t - W_s \sim \mathcal{N}(0, t - s)$;
3. $W_{t_2} - W_{t_1}, W_{t_3} - W_{t_2}, \dots, W_{t_n} - W_{t_{n-1}}$ are independent.
4. The map $t \mapsto W_t$ is continuous.

Theorem 6.4

For a Bronian motion,

$$P\left(\max_{t \in [0, T]} W_t > a\right) = 2P(W_T > a).$$

♡

Theorem 6.5 (Last Visit to 0)

Consider a one-dimensional random walk with $p = q = \frac{1}{2}$. Then

$$P(\text{last visit to 0 in } [0, 2n] \text{ is } 2k) = P(S_{2k} = 0) \cdot P(S_{2n-2k} = 0).$$

♡

Proof. Note that

$$\text{LHS} = P(S_{2k} = 0)P(S_i \neq 0, i = 2k + 1, \dots, 2n \mid S_{2k} = 0).$$

Let $2m = 2n - 2k$ (i.e., we shift the random walk $2k$ steps leftwards), then by Ballot Theorem,

$$\begin{aligned} P(S_i \neq 0, i = 1, 2, \dots, 2m) &= \frac{1}{2m} E(S_{2m}) = \frac{1}{2m} \cdot 2 \sum_{k=1}^{\infty} 2k \cdot P(S_{2m} = 2k) \\ &= \frac{2}{2m} \sum_{k=1}^{\infty} 2k \binom{2m}{m+k} \left(\frac{1}{2}\right)^{2m} \\ &= 2 \sum_{k=1}^{\infty} \left(\frac{1}{2}\right)^{2m} \left[\binom{2m-1}{m+k-1} - \binom{2m-1}{m+k} \right] \\ &= 2 \cdot \left(\frac{1}{2}\right)^{2m} \binom{2m-1}{m} \\ &= P(S_{2m} = 0). \end{aligned}$$

□

Remark Recall the Stirling Formula:

$$n! \stackrel{n \rightarrow \infty}{\sim} \left(\frac{n}{e}\right)^n \sqrt{2\pi n}. \quad (6.3)$$

Then

$$P(S_{2k} = 0) = \left(\frac{1}{2}\right)^{2k} \cdot \binom{2k}{k} \stackrel{k \rightarrow \infty}{\sim} \frac{1}{\sqrt{2\pi k}}.$$

Thus

$$\begin{aligned}
 P(\text{last visit to } 0 \text{ in } [0, 2n] \leq 2nx) &= \sum_{k \leq nx} P(S_{2k} = 0) \cdot P(S_{2n-2k} = 0) \\
 &\sim \sum_{k \leq nx} \frac{1}{\sqrt{\pi k}} \frac{1}{\sqrt{\pi(n-k)}} \\
 &\sim \int_0^x n \frac{1}{\pi \sqrt{u(1-u)}} du \\
 &= \frac{2}{\pi} \arcsin x.
 \end{aligned}$$

Exercise 6.1 Consider the random walk on $\{0, 1, \dots, N\}$ that reflects at 0. More precisely, let $S_0 = k, S_{j+1} = S_j + X_j$, where $X_j = 1$ if $S_j = 0$, otherwise $P[X_j = 1] = p$ and $P[X_j = -1] = q$. Compute the expected number of steps to first reach state N , as a function of k . Distinguish the cases $p = q$ and $p \neq q$.

Solution. Denote E_k as the expected number of steps to first reach N with $S_0 = k$. Then

$$\begin{aligned}
 E_N &= 0 \\
 E_0 &= 1 + E_1 \\
 E_k &= p(1 + E_{k+1}) + q(1 + E_{k-1}), \quad 1 \leq k \leq N-1
 \end{aligned}$$

That is

$$\begin{aligned}
 E_{k+1} &= \frac{1}{p}E_k - \frac{q}{p}E_{k-1} - \frac{1}{p} \\
 E_{k+2} &= \frac{1}{p}E_{k+1} - \frac{q}{p}E_k - \frac{1}{p} \\
 E_{k+2} - E_{k+1} &= \frac{1}{p}(E_{k+1} - E_k) - \frac{q}{p}(E_k - E_{k-1})
 \end{aligned}$$

Let $x_k = E_k - E_{k-1}$, then $x_{k+2} = \frac{1}{p}x_{k+1} - \frac{q}{p}x_k$

$$\begin{aligned}
 x_{k+2} - x_{k+1} &= \frac{q}{p}(x_{k+1} - x_k) \\
 x_{k+1} - x_k &= \left(\frac{q}{p}\right)^{k-1} (x_2 - x_1) \\
 (E_{k+1} - E_k) - (E_k - E_{k-1}) &= \left(\frac{q}{p}\right)^{k-1} (E_2 - 2E_1 + E_0)
 \end{aligned}$$

□

Exercise 6.2 In one-dimensional random walk with $p = q = \frac{1}{2}$, one has

$$E(\text{number of steps before hitting } 0, N) = k(N - k) \sim O(N^2).$$

Exercise 6.3 In $\mathbb{Z}^d (d \geq 1)$ random walk with $p = q = \frac{1}{2}$, one has

$$E(\text{number of steps before hitting boundary}) \sim O(N^2).$$

Theorem 6.6 (Loop erased random walk)

In \mathbb{Z}^2 -LERW,

$$E(\text{length of LERW}) = O(N^{5/4}).$$

In \mathbb{Z}^3 -LERW, there is α such that

$$E(\text{length of LERW}) \approx N^\alpha.$$

♡

Theorem 6.7 (Self-avoiding walks)

In \mathbb{Z}^2 -SAW, SAW_n , the distance between the destination and the origin satisfies:

$$\sqrt{\text{var}(SAW_n)} \approx O(N^{3/4}).$$



Chapter 7 Functions of random variables

7.1 Functions of random variables

Example 7.1 Let $X \sim \mathcal{N}(0, 1)$ and $Y = X^2$. Try to compute the probability density function of Y .

Solution. Note that

$$F_Y(y) = P(Y \leq y) = \begin{cases} P(-\sqrt{y} \leq X \leq \sqrt{y}) & \text{if } y \geq 0; \\ 0 & \text{else.} \end{cases}$$

Now let

$$\varphi(y) = \int_{-\infty}^y \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx$$

Then

$$P(-\sqrt{y} \leq X \leq \sqrt{y}) = \varphi(\sqrt{y}) - \varphi(-\sqrt{y}) = 2\varphi(\sqrt{y}) - 1$$

Finally,

$$f_Y(y) = 2\varphi'(\sqrt{y}) \cdot \frac{1}{2\sqrt{y}} = \frac{1}{\sqrt{2y}} e^{-y/2}, \quad y \geq 0.$$

□

The following corollary is a generalization of Exercise 5.3:

Corollary 7.1

If (X_1, X_2, \dots, X_n) has joint density function f , then $Y_1, Y_2, \dots, Y_n = T(X_1, X_2, \dots, X_n)$ has density function

$$f_{Y_1, \dots, Y_n}(y_1, \dots, y_n) = f(x_1(y_1, \dots, y_n), \dots, x_n(y_1, \dots, y_n)) \cdot \left| \frac{\partial(x_1, x_2, \dots, x_n)}{\partial(y_1, y_2, \dots, y_n)} \right|.$$

♡

Proof. Note that

$$\begin{aligned} P(Y_1, \dots, Y_n \in T(A)) &= P((X_1, \dots, X_n) \in A) = \int_A f(x_1, \dots, x_n) dx_1 \cdots dx_n \\ &= \int_A f(x_1(y_1, \dots, y_n), \dots, x_n(y_1, \dots, y_n)) \left| \frac{\partial(x_1, x_2, \dots, x_n)}{\partial(y_1, y_2, \dots, y_n)} \right| dy_1 \cdots dy_n \end{aligned}$$

□

Example 7.2 Let X_1, X_2, \dots have joint density function f . Define $X_1 = aY_1 + bY_2, X_2 = cY_1 + dY_2$ with $ad \neq bc$. Find $f_{Y_1, Y_2}(y_1, y_2)$.

Solution. The jacobian is given by:

$$|J| = \begin{vmatrix} \frac{\partial x_1}{\partial y_1} & \frac{\partial x_1}{\partial y_2} \\ \frac{\partial x_2}{\partial y_1} & \frac{\partial x_2}{\partial y_2} \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = |ad - bc|$$

and thus

$$f_{Y_1, Y_2}(y_1, y_2) = f(ay_1 + by_2, cy_1 + dy_2) \cdot |ad - bc|.$$

□

Example 7.3 Suppose independent X, Y has joint density function f . Show that $U := XY$ has density function

$$f_U(u) = \int_{-\infty}^{\infty} f\left(v, \frac{u}{v}\right) \frac{1}{|v|} dv.$$

Proof. Consider the transformation

$$\begin{cases} u = xy \\ v = x \end{cases}$$

and then

$$\begin{cases} x = v \\ y = \frac{u}{v} \end{cases}$$

with Jacobian

$$|J| = \begin{vmatrix} 0 & 1 \\ \frac{1}{v} & -\frac{u}{v^2} \end{vmatrix} = \frac{1}{|v|}.$$

Then

$$f_{U,V}(u, v) = f\left(v, \frac{u}{v}\right) \cdot \frac{1}{|v|}$$

so that

$$f_U(u) = \int_{-\infty}^{\infty} f_{X,Y}\left(v, \frac{u}{v}\right) \cdot \frac{1}{|v|} dv.$$

□

Example 7.4 Suppose independent X, Y has joint density function f . Show that $U := X/Y$ has density function

$$f_U(u) = \int_{-\infty}^{\infty} f_{X,Y}(u, uv) |v| dv.$$

Proof. Let $V = Y$, considering the transformation

$$\begin{cases} u = \frac{x}{y} \\ v = y \end{cases}$$

and then

$$\begin{cases} x = uv \\ y = v \end{cases}$$

with Jacobian

$$|J| = \begin{vmatrix} v & u \\ 0 & 1 \end{vmatrix} = |v|.$$

Thus

$$f(u, v) = f_X(uv) \cdot f_Y(v) |J| = f_X(uv) f_Y(v) |v|$$

so that

$$f_U(u) = \int_{-\infty}^{\infty} f_{X,Y}(uv, v) \cdot |v| dv.$$

□

Example 7.5 Let X_1, X_2 be independent r.v.s obeying $\exp(\lambda)$. Find the joint density function of $Y_1 = X_1 + X_2$ and $Y_2 = \frac{X_1}{X_2}$.

Solution. Note that from

$$\begin{cases} Y_1 = X_1 + X_2 \\ Y_2 = \frac{X_1}{X_2} \end{cases}$$

we get

$$\begin{cases} X_1 = \frac{Y_1 Y_2}{1 + Y_2} \\ X_2 = \frac{Y_1}{1 + Y_2} \end{cases}$$

with Jacobian

$$|J| = \begin{vmatrix} \frac{y_2}{1+y_2} & \frac{y_1}{(1+y_2)^2} \\ 1 & -\frac{y_1}{(1+y_2)^2} \end{vmatrix} = \frac{|y_1|}{(1+y_2)^2}$$

Thus

$$\begin{aligned} f_{Y_1, Y_2}(y_1, y_2) &= f_{X_1, X_2}\left(\frac{y_1 y_2}{1+y_2}, \frac{y_1}{1+y_2}\right) \cdot \frac{|y_1|}{(1+y_2)^2} \\ &= \lambda^2 e^{-\lambda y_1} |y_1| \frac{1}{(1+y_2)^2} \end{aligned}$$

□

Example 7.6 Let $X_1, X_2 \sim \mathcal{N}(0, 1)$ be independent. Then $X_1 \pm X_2$ are independent.

Example 7.7(Rayleigh Distribution) Let X, Y be independent $\mathcal{N}(0, 1)$. Find the joint density function of $R = \sqrt{X^2 + Y^2}$, $\Theta = \arctan \frac{Y}{X}$.

Solution. Note that

$$\begin{cases} X = R \cos \Theta \\ Y = R \sin \Theta \end{cases}$$

has Jacobian

$$|J| = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = r.$$

Furthermore,

$$f_{X, Y}(x, y) = \frac{1}{2\pi} e^{-\frac{x^2 + y^2}{2}}$$


Thus

$$f_{R, \Theta}(r, \theta) = f_{X, Y}(r \cos \theta, r \sin \theta) \cdot r = \frac{1}{2\pi} e^{-r^2/2} r.$$

That is,

$$\Theta \sim \text{unif}[0, 2\pi), \quad f_R(r) = r e^{-r^2/2}.$$

□

 **Exercise 7.1** Show that R^2, Θ are independent and $R^2 \sim \exp\left(\frac{1}{2}\right)$.

Solution. Let $Z = R^2$. Note that

$$\begin{cases} x = \sqrt{z} \cos \theta, \\ y = \sqrt{z} \sin \theta \end{cases}$$

has Jacobian

$$|J| = \begin{vmatrix} \frac{\cos \theta}{2\sqrt{z}} & -\sqrt{z} \sin \theta \\ \frac{\sin \theta}{2\sqrt{z}} & \sqrt{z} \cos \theta \end{vmatrix} = \frac{1}{2}.$$

Thus

$$f_{Z,\Theta}(z, \theta) = f_{X,Y}(\sqrt{z} \cos \theta, \sqrt{z} \sin \theta) \cdot \frac{1}{2} = \frac{1}{2\pi} \frac{1}{2} e^{-z/2}$$

and $\Theta \sim \text{unif}[0, 2\pi)$, $R^2 \sim \exp(1/2)$. □

The Rayleigh Distribution suggests that one can use uniform random variables to generate standard normals:

Exercise 7.2 Let $U_1, U_2 \sim \text{unif}[0, 1]$ be independent, then

1. $Z_1 = -2 \ln U_1 \sim \exp\left(\frac{1}{2}\right)$: $P(-2 \ln U_1 > x) = P(U_1 < e^{-x/2}) = e^{-x/2}$ for $x > 0$;
2. (Box-Muller transformation) $Z_2 = 2\pi U_2 \sim \text{unif}[0, 2\pi)$. Take

$$\begin{cases} R^2 = -2 \ln U_1, \\ \Theta = 2\pi U_2 \end{cases}$$

and then

$$\begin{cases} X = \sqrt{-2 \ln U_1} \cos(2\pi U_2), \\ Y = \sqrt{-2 \ln U_1} \sin(2\pi U_2), \end{cases}$$

with $(X, Y) \sim \mathcal{N}(0, 1)$ being independent.

Solution. Let $z_1 = -2 \ln u_1$, then $u_1 = e^{-\frac{1}{2}z_1}$, $\frac{du_1}{dz_1} = -\frac{1}{2}e^{-\frac{1}{2}z_1}$. Hence for $z_1 > 0$, $Z_1 = -2 \ln U_1$ has density function

$$f_{Z_1}(z_1) = f_{U_1}(e^{-\frac{1}{2}z_1}) \left| \frac{du_1}{dz_1} \right| = \frac{1}{2} e^{-\frac{1}{2}z_1}$$

That is $Z_1 \sim \exp(1/2)$. Similarly, one deduces that $Z_2 \sim \text{unif}(0, 2\pi)$.

From $x^2 + y^2 = -2 \ln u_1$, $\frac{y}{x} = \tan(2\pi u_2)$, we have $u_1 = \exp\left\{-\frac{1}{2}(x^2 + y^2)\right\}$ and $u_2 = \frac{1}{2\pi} \arctan \frac{y}{x}$.

Thus the Jacobian is

$$J = \begin{vmatrix} \frac{\partial u_1}{\partial x} & \frac{\partial u_1}{\partial y} \\ \frac{\partial u_2}{\partial x} & \frac{\partial u_2}{\partial y} \end{vmatrix} = -\frac{1}{2\pi} \exp\left\{-\frac{1}{2}(x^2 + y^2)\right\}$$

Finally,

$$f_{X,Y}(x, y) = f_{U_1,U_2}(u_1, u_2) \cdot |J| = \frac{1}{2\pi} \exp\left\{-\frac{1}{2}(x^2 + y^2)\right\}, \quad -\infty < x, y < \infty$$

implying $(X, Y) \sim \mathcal{N}(0, 1)$ being independent. □

Example 7.8 Let $X, Y \sim \mathcal{N}(0, 1)$ be independent and define

$$\begin{cases} U = \sigma_1 X, \\ V = \sigma_2 \rho X + \sigma_2 \sqrt{1 - \rho^2} Y. \end{cases}$$

Show that their joint density function is

$$f_{U,V}(u, v) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} e^{-\frac{1}{2}Q(u,v)},$$

where

$$Q(u, v) = \frac{1}{1-\rho^2} \left(\left(\frac{u}{\sigma_1} \right)^2 - 2\rho \frac{u}{\sigma_1} \frac{v}{\sigma_2} + \left(\frac{v}{\sigma_2} \right)^2 \right).$$

Example 7.9 Let U, V be bivariate normal. Compute $E(UV)$ and $E(U | V)$.

Solution. As in Example 7.8, we normalize U and V to obtain X and Y which are independent and normal distributed. Then

$$E(UV) = \sigma_1\sigma_2\rho E(X^2) + \sigma_1\sigma_2\sqrt{1-\rho^2}E(XY) = \sigma_1\sigma_2\rho$$

since by independence, $E(XY) = E(X) \cdot E(Y) = 0 \cdot 0 = 0$.

Now given $U = u$, we have $V = \sigma_2\rho\frac{u}{\sigma_1} + \sigma_2\sqrt{1-\rho^2}Y \sim \mathcal{N}\left(\frac{\sigma_2}{\sigma_1}\rho u, \sigma_2^2(1-\rho^2)\right)$. Therefore,

$$E(V | U = u) = \frac{\sigma_2}{\sigma_1}\rho u, \quad E(V | U) = \frac{\sigma_2}{\sigma_1}\rho U, \quad \text{var}(V | U) = \sigma_2^2(1-\rho^2).$$

□

Exercise 7.3 Given X_1, X_2, \dots, X_n being independent and $X_i \sim \exp(\lambda_i)$, show that

$$P(X_i = \min\{X_1, X_2, \dots, X_n\}) = \frac{\lambda_i}{\lambda_1 + \lambda_2 + \dots + \lambda_n}.$$

Proof. By independence, we know (X_1, X_2, \dots, X_n) has density function

$$f(x_1, x_2, \dots, x_n) = \prod_{j=1}^n \lambda_j e^{-\lambda_j x_j}$$

Note that

$$\begin{aligned} & \{X_i = \min\{X_1, X_2, \dots, X_n\}\} \\ &= \{X_1 \geq X_i, \dots, X_{i-1} \geq X_i, 0 < X_i < \infty, X_{i+1} \geq X_i, \dots, X_n \geq X_i\}. \end{aligned}$$

Thus

$$\begin{aligned} & P(X_i = \min\{X_1, X_2, \dots, X_n\}) \\ &= \int_0^\infty \int_{x_i}^\infty \dots \int_{x_i}^\infty \int_{x_i}^\infty \dots \int_{x_i}^\infty \prod_{j=1}^n \lambda_j e^{-\lambda_j x_j} dx_1 \dots dx_{i-1} dx_{i+1} \dots dx_n dx_i \\ &= \int_0^\infty \lambda_i e^{-(\lambda_1 + \lambda_2 + \dots + \lambda_n)x_i} dx_i = \frac{\lambda_i}{\lambda_1 + \lambda_2 + \dots + \lambda_n}. \end{aligned}$$

□

Exercise 7.4 Given X_1, X_2, \dots, X_n being continuous and i.i.d, show that

$$P(X_n > \max\{X_1, X_2, \dots, X_{n-1}\}) = \frac{1}{n}.$$

Proof. Let $X_i (i = 1, 2, \dots, n)$ has probability density function $f(x)$. Then by independence, their joint density function is

$$f(x_1, x_2, \dots, x_n) = \prod_{i=1}^n f(x_i).$$

Note that

$$\{X_n > \max\{X_1, X_2, \dots, X_n\}\} = \{X_1 < X_n, X_2 < X_n, \dots, X_{n-1} < X_n, -\infty < X_n < \infty\}$$

and thus

$$P(X_n > \max\{X_1, X_2, \dots, X_n\}) = \int_{-\infty}^{\infty} \int_{-\infty}^{x_n} \dots \int_{-\infty}^{x_n} \prod_{i=1}^n f(x_i) dx_1 dx_2 \dots dx_{n-1} dx_n$$

The distribution of X_i is given by $F(x) = \int_{-\infty}^x f(t) dt$ and hence

$$P(X_n > \max\{X_1, X_2, \dots, X_n\}) = \int_{-\infty}^{\infty} F^{n-1}(x_n) dF(x_n) = \left[\frac{1}{n} F^n(x_n) \right]_{-\infty}^{\infty} = \frac{1}{n}.$$

□

7.2 Generating functions

Definition 7.1

A sequence $a = \{a_i : i = 0, 1, 2, \dots\}$ of real numbers may contain a lot of information. One concise way of storing this information is to wrap up the numbers together in a “generating function”. For example, the (ordinary) generating function of the sequence a is the function G_a defined by

$$G_a(s) := \sum_{n \geq 0} a_n s^n$$

and a_n can be computed by $a_n = \frac{1}{n!} G_a^{(n)}(0)$.



Example 7.10 Let $a_n = \binom{N}{n}$, then

$$G_a(s) = \sum_n \binom{N}{n} s^n = (1 + s)^N.$$

Example 7.11 Let $a_n = e^{in\theta}$, forming an orthonormal basis of $L^2[0, 2\pi)$, then

$$G_a(s) = \sum_{n \geq 0} e^{in\theta} s^n = \frac{1}{1 - e^{i\theta} s}.$$

Theorem 7.1 (Convolution of sequences)

Given $\{a_n\}, \{b_n\}$, we define $c_n = a_0 b_n + a_1 b_{n-1} + \dots + a_n b_0$, then

$$G_c(s) = G_a(s) \cdot G_b(s).$$



Proof. Note that

$$\begin{aligned} G_c(s) &= \sum_{n \geq 0} c_n s^n = \sum_{n=0}^{\infty} \sum_{k=0}^n a_k b_{n-k} s^n \\ &= \sum_{k=0}^{\infty} a_k s^k \sum_{n=k}^{\infty} b_{n-k} s^{n-k} \end{aligned}$$

$$= G_a(s)G_b(s).$$

□

Definition 7.2

Let X be a discrete random variable taking value in \mathbb{N} . The probability generating function of X is given by:

$$G_X(s) = E(s^X) = \sum_{n \geq 0} P(X = n) \cdot s^n.$$



Example 7.12 For $X \sim \text{Bernoulli}(p)$, $G_X(s) = (1 - p) + ps$.

Example 7.13 For $X \sim \text{Geometric}(p)$,

$$G_X(s) = \sum_{n \geq 0} (1 - p)^{n-1} ps^n = ps \sum_{n \geq 0} (1 - p)^{n-1} s^{n-1} = \frac{ps}{1 - (1 - p)s}.$$

Example 7.14 For $X \sim \text{Binomial}(n, p)$, we have the decomposition $X = Y_1 + \dots + Y_n$ for each $Y_i \sim \text{Bernoulli}(p)$ being i.i.d. Then

$$G_X(s) = G_{Y_1}(s) \cdots G_{Y_n}(s) = (1 - p + ps)^n.$$

Example 7.15 For $X \sim \text{Poisson}(\lambda)$,

$$G_X(s) = E(s^X) = \sum_k s^k \frac{\lambda^k}{k!} e^{-\lambda} = e^{\lambda(s-1)}.$$

Corollary 7.2

If X and Y are independent, then

$$G_{X+Y}(s) = G_X(s) \cdot G_Y(s).$$



Example 7.16 Let $X \sim \text{Poisson}(\lambda)$ and $Y \sim \text{Poisson}(\mu)$ be independent. Show that $X + Y \sim \text{Poisson}(\lambda + \mu)$.

Proof. Note that

$$G_X(s) = e^{\lambda(s-1)}, \quad G_Y(s) = e^{\mu(s-1)},$$

$$G_{X+Y}(s) = G_X(s) \cdot G_Y(s) = e^{(\lambda+\mu)(s-1)}$$

That is $X + Y \sim \text{Poisson}(\lambda + \mu)$. □

Definition 7.3

In general, the probability generating function of a random variable X is $G_X(s) := E(s^X)$.



Note that G does indeed generate the sequence $\{f(i) : i \geq 0\}$ since

$$E(s^X) = \sum_k s^k P(X = k) = \sum_i s^i f(i).$$

Remark

1. There is a radius of convergence R such that $G_X(s)$ converges absolutely for $|s| < R$ and diverges for $|s| > R$ where $R \geq 1$ (because $G_X(1) = 1$). The sum is uniformly convergent on sets of the form $\{s : |s| \leq R'\}$ for any $R' < R$.
2. For $|s| < R$, one can differentiate and integrate term by term.
3. If $G_a(s) = G_b(s)$ for $s \in (-\delta, \delta)$ for some $\delta > 0$, then $G_a \equiv G_b$ and $a_n = \frac{1}{n!} G^{(n)}(0) = b_n$.

4. (Abel's Theorem) If $a_i \geq 0$ and $G_a(s)$ converges absolutely for $|s| < 1$, then $\lim_{s \uparrow 1} G_a(s) = G_a(1) = \sum_{n=0}^{\infty} a_n$.

Lemma 7.1

If X has generating function $G(s)$, then

1. $G'(1) = E(X)$;
2. $G^{(k)}(1) = E[X(X-1)\cdots(X-k+1)]$.



Proof. Note that

$$\begin{aligned} \frac{d}{ds} E(s^X) &= E(Xs^{X-1}), \\ \frac{d}{ds} G^{(k)}(s) &= E[X(X-1)\cdots(X-k+1)s^{X-k}]. \end{aligned}$$

We know the radius of convergence $R \geq 1$ and thus one can apply Abel's Theorem to $s \uparrow 1$. □

Example 7.17 We now have

1. $E(X) = G'_X(1)$;
2. $E(X^2) = E(X(X-1)) + E(X) = G''(1) + G'(1)$;
- 3.

$$\begin{aligned} \text{var}(X) &= E(X^2) - E^2(X) = E(X(X-1)) + E(X) - E^2(X) \\ &= G''(1) + G'(1) - (G'(1))^2. \end{aligned}$$

Example 7.18 Consider the coin flip satisfying Bernoulli(p). Palyer A wins if the m -th head occurs before the n -th tail. Try to compute $P(\text{A wins})$?

Solution. We use P_{mn} to denote the probability of event “the m -th head occurs before the n -th tail”. It is elementary, by conditioning on the outcome of the first toss, that the probability P_{mn} , that A wins, satisfies

$$\begin{cases} P_{mn} = p \cdot P_{m-1,n} + q \cdot P_{m,n-1} \\ P_{m0} = 0 \\ P_{0n} = 1 \end{cases}$$

Consider the generating function $G(x, y) = \sum_{\substack{m \geq 0 \\ n \geq 0}} P_{mn} x^m y^n$:

$$\begin{aligned} G(x, y) &= \sum_{\substack{m \geq 1 \\ n \geq 0}} p \cdot P_{m-1,n} x^m y^n + \sum_{\substack{m \geq 0 \\ n \geq 1}} q \cdot P_{m,n-1} x^m y^n \\ &\quad + \sum_{n \geq 0} P_{0n} y^n + \sum_{m \geq 0} P_{m0} x^m \\ &= \sum_{\substack{m \geq 1 \\ n \geq 0}} px \cdot P_{m-1,n} x^{m-1} y^n + \sum_{\substack{m=0 \\ n \geq 0}} P_{0n} y^n \\ &\quad + \sum_{\substack{m \geq 0 \\ n \geq 1}} qy \cdot P_{m,n-1} x^m y^{n-1} + \sum_{\substack{m \geq 0 \\ n=0}} P_{m0} x^m \end{aligned}$$


Thus

$$G(x, y) = (px + qy)G(x, y) + \frac{1}{1-y}$$

$$G(x, y) = \frac{1}{(1-y)(1-px-xy)}$$

from which one may derive the required information by expanding in powers of x and y and finding the coefficient of $x^m y^n$. \square

Definition 7.4

The moment generating function of X is given by $M_X(t) := E(e^{tX}) = G_X(e^t)$. 

If $t < R$, then

$$M_X(t) = \sum_{n=0}^{\infty} \frac{t^n}{n!} E(X^n); \quad E(X^n) = \frac{1}{n!} M_X^{(n)}(0).$$

Example 7.19 For $X \sim \text{Poisson}(\lambda)$, one has

$$M_X(t) = G_X(e^t) = e^{\lambda(e^t-1)}.$$

Example 7.20 For $X \sim \mathcal{N}(0, 1)$, one has


$$M_X(t) = e^{\frac{1}{2}t^2}.$$

For $y \sim \mathcal{N}(\mu, \sigma^2)$, one has

$$M_Y(t) = E(e^{t(\sigma x + \mu)}) = e^{\mu t} e^{\frac{1}{2}\sigma^2 t^2}.$$

Theorem 7.2 (Random sum formula)

For a series of i.i.d. random variables $\{X_n\}$ with generating function G_X , we have the random sum

$$S_N = \sum_{i=1}^N X_i \text{ for } N \text{ being independent of } X_i. \text{ Then the generating function of } S_N \text{ is } G_N(G_X(s)).$$


Proof.

$$\begin{aligned} G_{S_N}(s) &= E[s^{S_N}] = E[E[s^{S_N} | N]] = \sum_n P(N = n) E[s^{S_N} | N = n] \\ &= \sum_n P(N = n) E[s^{X_1}]^n = \sum_n P(N = n) (G_X(s))^n \\ &= G_N(G_X(s)). \end{aligned}$$

\square

Example 7.21 Let $N \sim \text{Poisson}(\lambda)$ with X_i be i.i.d. Bernoulli(p). Set $S_N = X_1 + \dots + X_n$. Try to compute the distribution of S_N .

Solution. Note that

$$G_N(s) = e^{\lambda(s-1)}, \quad G_X(s) = ps + 1 - p$$


and

$$G_{S_N}(s) = G_N(G_X(s)) = e^{\lambda(ps+1-p-1)} = e^{\lambda p(s-1)},$$

implying $S_N \sim \text{Poisson}(\lambda p)$. \square

Definition 7.5

The joint generating function for X_1 and X_2 is defined by

$$G_{X_1, X_2}(s_1, s_2) = E(s_1^{X_1} s_2^{X_2}).$$


Theorem 7.3

X_1 and X_2 are independent $\iff G_{X_1, X_2}(s_1, s_2) = G_{X_1}(s_1)G_{X_2}(s_2)$. ♥

Proof. \implies . Recall that if X_1 and X_2 are independent, then $E[f(X_1)g(X_2)] = E[f(X_1)] \cdot E[g(X_2)]$ for any measurable f, g . Take $f(x) = s_1^x$ and $g(x) = s_2^x$, then $G_{X_1, X_2}(s_1, s_2) = G_{X_1}(s_1)G_{X_2}(s_2)$.

\impliedby . Only for discrete cases. Note that

$$G_{X_1, X_2}(s_1, s_2) = \sum_{i, j} P(X_1 = i, X_2 = j) s_1^i s_2^j,$$

$$G_{X_1}(s_1)G_{X_2}(s_2) = \sum_i P(X_1 = i) s_1^i \sum_j P(X_2 = j) s_2^j.$$

Now we compare the coefficient of $s_1^i s_2^j$ and obtain that $P(X_1 = i, X_2 = j) = P(X_1 = i)P(X_2 = j)$. □

Example 7.22 Given independent X and Y obeying $\mathcal{N}(\mu, \sigma^2)$, find the joint distribution of $X + Y$ and $X - Y$.

Solution. Compute the joint generating function:

$$\begin{aligned} E[e^{t(X+Y)} e^{s(X-Y)}] &= E(e^{(t+s)X}) E(e^{(t-s)Y}) \\ &= e^{\mu(t+s) + \frac{1}{2}\sigma^2(t+s)^2} e^{\mu(t-s) + \frac{1}{2}\sigma^2(t-s)^2} \\ &= e^{2\mu t + \sigma^2 t^2} e^{\sigma^2 s^2}. \end{aligned}$$

That is $X + Y \sim \mathcal{N}(2\mu, 2\sigma^2)$ and $X - Y \sim \mathcal{N}(0, 2\sigma^2)$ and they are independent. □

Exercise 7.5 How many methods are there using 1, 2, 5, and 10 to sum up for 30?

Solution. We assign $z, z^2, z^5,$ and z^{10} to represent 1, 2, 5, and 10 respectively, with their powers indicating multiples of those values. This leads us to the generation function:

$$G(z) = (1 + z + z^2 + \dots)(1 + z^2 + z^4 + \dots)(1 + z^5 + z^{10} + \dots)(1 + z^{10} + z^{20} + \dots).$$

The coefficient of z^m in this function precisely reflects the number of ways to form the sum m using combinations of 1, 2, 5, and 10. For example, z^4 can be decomposed as $z^1 \cdot z^1 \cdot z^1 \cdot z^1, z^2 \cdot z^2, z^2 \cdot z^1 \cdot z^1$ and its coefficient is 3 in $G(z)$, that is, there are three ways in total to sum up for 4. In general,

$$G(z) = \frac{1}{1-z} \frac{1}{1-z^2} \frac{1}{1-z^5} \frac{1}{1-z^{10}}$$

and what we want is the coefficient of z^{30} :

$$\frac{1}{30!} \left. \frac{d^{30} G(z)}{dz^{30}} \right|_{z=0}.$$

□

Exercise 7.6 Show that

$$\sum_{\substack{j, k \\ j+k=l}} \binom{m+j-1}{j} \binom{n+k-1}{k} = \binom{m+n+l-1}{l}.$$

Proof. First recall that

$$(1+z)^n = \sum_k \binom{n}{k} z^k$$

and

$$\frac{1}{(1+z)^n} = 1 - nz + \frac{(-n)(-n-1)}{2} z^2 + \frac{(-n)(-n-1)(-n-2)}{3!} z^3 + \dots$$

$$= \sum_k (-1)^k z^k \frac{(n+k-1)!}{(n-1)!k!} = \sum_k (-1)^k \binom{n+k-1}{k} z^k.$$

Note that $(-1)^l \times \text{LHS}$ is exactly the coefficient of z^l in $\frac{1}{(1+z)^m} \frac{1}{(1+z)^n}$ and $(-1)^l \times \text{RHS}$ is exactly the coefficient of z^l in $\frac{1}{(1+z)^{m+n}}$. \square

Exercise 7.7 Let $U : \Omega \rightarrow [0, 1]$ be a random variable and $U(\omega) := (0.\xi_1(\omega_1)\xi_2(\omega_2)\cdots\xi_n(\omega_n)\cdots)_2$ be the binary expression (that is each ξ_i only takes value in 0 or 1). Show that $\xi_i \sim \text{i.i.d. Bernoulli}(1/2) \iff U \sim \text{Unif}[0, 1]$.

Proof. Note that

$$X = (0.\xi_1\xi_2\cdots)_2 = \sum_{i \geq 1} \frac{\xi_i}{2^i}$$

and

$$E(e^{tX}) = E(e^{t \sum_{i \geq 1} \frac{\xi_i}{2^i}}) = \prod_{i \geq 1} E(e^{t \frac{\xi_i}{2^i}}) = \prod_{i \geq 1} \left(\frac{1}{2} + \frac{1}{2} e^{\frac{t}{2^i}} \right)$$

Now we continue with

$$\begin{aligned} \prod_{i=1}^n \left(\frac{1}{2} + \frac{1}{2} e^{\frac{t}{2^i}} \right) &= \frac{1}{2^n} \frac{1}{1 - e^{\frac{t}{2^n}}} \prod_{i=1}^n \left(1 + e^{\frac{t}{2^i}} \right) \left(1 - e^{\frac{t}{2^n}} \right) \\ &= \frac{1}{2^n} \frac{1 - e^t}{1 - e^{t/2^n}} \xrightarrow{n \rightarrow \infty} -\frac{1 - e^t}{t}. \end{aligned}$$

This is exactly the moment generating function for $U \sim \text{Unif}[0, 1]$:

$$E(e^{tU}) = \int_0^1 e^{tu} du = -\frac{1 - e^t}{t}.$$

\square

Exercise 7.8 A coin is tossed repeatedly, heads appearing with probability p on each toss.

- (a) Let X be the number of tosses until the first occasion by which three heads have appeared successively. Write down a difference equation for $f(k) = \mathbb{P}(X = k)$ and solve it. Now write down an equation for $\mathbb{E}(X)$ using conditional expectation. (Try the same thing for the first occurrence of HTH).
- (b) Let N be the number of heads in n tosses of the coin. Write down $G_N(s)$. Hence find the probability that:
- (1) N is divisible by 2,
 - (2) N is divisible by 3.

Solution. (a) The initial sequences T, HT, HHT, HHH induce a partition of the sample space. By conditioning on this initial sequence, we obtain $f(k) = qf(k-1) + pqf(k-2) + p^2qf(k-3)$ for $k > 3$, where $p + q = 1$. Also $f(1) = f(2) = 0, f(3) = p^3$. In principle, this difference equation may be solved in the usual way (see Appendix I). An alternative is to use generating functions. Set $G(s) = \sum_{k=1}^{\infty} s^k f(k)$, multiply throughout the difference equation by s^k and sum, to find that $G(s) = p^3 s^3 / \{1 - qs - pqs^2 - p^2qs^3\}$. To find the coefficient of s^k , factorize the denominator, expand in partial fractions, and use the binomial series.

Another equation for $f(k)$ is obtained by observing that $X = k$ if and only if $X > k-4$ and the last four

tosses were THHH. Hence

$$f(k) = qp^3 \left(1 - \sum_{i=1}^{k-4} f(i) \right), \quad k > 3.$$

Applying the first argument to the mean, we find that $\mu = \mathbb{E}(X)$ satisfies $\mu = q(1 + \mu) + pq(2 + \mu) + p^2q(3 + \mu) + 3p^3$ and hence $\mu = (1 + p + p^2) / p^3$.

As for HTH, consider the event that HTH does not occur in n tosses, and in addition the next three tosses give HTH. The number Y until the first occurrence of HTH satisfies

$$\mathbb{P}(Y > n)p^2q = \mathbb{P}(Y = n + 1)pq + \mathbb{P}(Y = n + 3), \quad n \geq 2.$$

Sum over n to obtain $\mathbb{E}(Y) = (pq + 1) / (p^2q)$.

(b) $G_N(s) = (q + ps)^n$, in the obvious notation.

(1) $\mathbb{P}(2 \text{ divides } N) = \frac{1}{2} \{G_N(1) + G_N(-1)\}$, since only the coefficients of the even powers of s contribute to this probability.

(2) Let ω be a complex cube root of unity. Then the coefficient of $\mathbb{P}(X = k)$ in $\frac{1}{3} \{G_N(1) + G_N(\omega) + G_N(\omega^2)\}$ is

$$\begin{aligned} \frac{1}{3} \{1 + \omega^3 + \omega^6\} &= 1, \text{ if } k = 3r \\ \frac{1}{3} \{1 + \omega + \omega^2\} &= 0, \text{ if } k = 3r + 1 \\ \frac{1}{3} \{1 + \omega^2 + \omega^4\} &= 0, \text{ if } k = 3r + 2 \end{aligned}$$

for integers r . Hence $\frac{1}{3} \{G_N(1) + G_N(\omega) + G_N(\omega^2)\} = \sum_{r=0}^{\lfloor \frac{1}{3}n \rfloor} \mathbb{P}(N = 3r)$, the probability that N is a multiple of 3. Generalize this conclusion. □

Definition 7.6

Let $\{X_n\}_{n \geq 1}$ and Y be random variables, then we say X_n **converges in distribution** to Y , denoted by $X_n \xrightarrow{D} Y$, if $\lim_{n \rightarrow \infty} P(X_n \leq x) = P(Y \leq x) = F_Y(x)$ for all x such that F_Y is continuous at x . ♣

Theorem 7.4

If $M_{X_n}(t) \rightarrow M_Y(t)$ for all $t \in (-\delta, \delta)$, then $X_n \xrightarrow{D} Y$. ♥

Theorem 7.5 (Chernoff bound)

Let X_i be i.i.d. Bernoulli(p) and define the partial sum $X = \sum_{i=1}^N X_i$. Then

$$P[X \geq (1 + \delta)np] \leq \left(\frac{e^\delta}{(1 + \delta)^{1 + \delta}} \right)^{np}.$$
♥

Theorem 7.6 (Hoeffding inequality)

Let X_i be i.i.d. with $X_i \in [a_i, b_i]$ and $\mu = \sum_{i=1}^n E(X_i)$. Then

$$P\left(\left| \sum_{i=1}^n X_i - \mu \right| \geq t \right) \leq C \cdot \exp \left\{ - \frac{2t^2}{\sum_{i=1}^n (b_i - a_i)^2} \right\}.$$
♥

Lemma 7.2

Let X be a random variable with $E(X) = 0$, $X \in [a, b]$. Then

$$E(e^t X) \leq e^{\frac{1}{8}t^2(b-a)^2}.$$



7.3 Galton-Walter process

Suppose that a population evolves in generations, and let Z_n be the number of members of the n th generation. Each member of the n th generation gives birth to a family, possibly empty, of members of the $(n + 1)$ th generation; the size of this family is a random variable. Furthermore, suppose that $Z_0 = 1$ and the number of children for each individual is i.i.d. and each individual has generating function G . Let $G_n(s) = E(s^{Z_n})$.

Lemma 7.3

$$G_n(s) = G_{n-1}(G(s)) = \underbrace{G(G(\cdots G(s)))}_{n \text{ times}}.$$



Proof. Note that $Z_n = X_1 + X_2 + \cdots + X_{Z_{n-1}}$, where X_i is the number of members of the n th generation which stem from the i th member of the $n - 1$ th generation. These variables are independent and are identically distributed with the same distribution. By random sum formula,

$$G_n(s) = G_{n-1}(G_{X_1}(s)) = G_{n-1}(G_1(s)) = G_{n-1}(G(s)).$$

□

In practice, we can evaluate moments of Z_n to moments of Z_1 :

Lemma 7.4

Let $\mu = E(Z_1)$ and $\sigma^2 = \text{var}(Z_1)$. Then $E(Z_n) = \mu^n$ and $\text{var}(Z_n) = \begin{cases} n\sigma^2, & \mu = 1, \\ \frac{\sigma^2(\mu^n - 1)\mu^{n-1}}{\mu - 1}, & \mu \neq 1. \end{cases}$



Proof. Differentiate $G_n(s) = G(G_{n-1}(s))$ and set $s = 1$:

$$G'_n(1) = G'(G_{n-1}(1))G'_{n-1}(1)$$

obtaining $E(Z_n) = \mu E(Z_{n-1})$ and hence $E(Z_n) = \mu^n$.

Differentiate $G_n(s) = G(G_{n-1}(s))$ twice and set $s = 1$:

$$G''_n(1) = G''(1)G'_{n-1}(1)^2 + G'(1)G''_{n-1}(1)$$

Note that

$$\begin{aligned} G''(1) &= E[Z_1(Z_1 - 1)] = E(Z_1^2) - E(Z_1) \\ &= \text{var}(Z_1) + (E(Z_1))^2 - E(Z_1) \\ &= \sigma^2 + \mu^2 - \mu \end{aligned}$$

and thus

$$G''_n(1) = G''(1)\mu^{2(n-1)} + \mu G''_{n-1}(1).$$

Solve the recursion to get the desired.

□

Example 7.23 (Geometric branching) Assume $P(Z_1 = k) = p^k q$ where $q = 1 - p$ and $G(s) = E(s^{Z_1}) = \frac{q}{1 - ps}$. Show that $G_n(s) = \begin{cases} \frac{n - (n-1)s}{n+1 - ns}, & p = q = \frac{1}{2}, \\ \frac{q(p^n - q^n - ps(p^{n-1} - q^{n-1}))}{p^{n+1} - q^{n+1} - ps(p^n - q^n)}, & p \neq q. \end{cases}$

Extinction and non-extinction Note that

$$P(Z_n = 0) = G_n(0) = \begin{cases} \frac{n}{n+1}, & p = q = \frac{1}{2} \\ \frac{q(p^n - q^n)}{p^{n+1} - q^{n+1}}, & p \neq q. \end{cases}$$


As $n \uparrow +\infty$, one has $\{Z_1 = 0\} \uparrow \bigcup_{n=1}^{\infty} \{Z_n = 0\} = \{\text{extinction}\}$. Finally,

$$P(\text{extinction}) = \begin{cases} 1, & p \leq \frac{1}{2}, \\ \frac{q}{p}, & p > \frac{1}{2}. \end{cases}$$

Also note that $E(Z_1) = \frac{p}{q}$ and thus the final conclusion is an extinction happens in probability 1 if $E(Z_1) \leq 1$ whereas there is a positive probability of infinite growth given $E(Z_1) > 1$.


Theorem 7.7 (General case)

- If $\mu = E(Z_1) < 1$, then $P(\text{extinction}) = 1$;
- If $\mu = E(Z_1) > 1$, then $P(\text{extinction}) < 1$;
- If $\mu = E(Z_1) = 1$ and $\sigma^2 = \text{var}(Z_1) > 0$, then $P(\text{extinction}) = 1$.

Moreover, if we set $\eta = P(\text{extinction})$, then η is the smallest non-negative solution to $s = G(s)$. 

Proof. Let $\eta_n = P(Z_n = 0)$, then $\eta_n = G_n(0) = G(G_{n-1}(0)) = G(\eta_{n-1})$. Take $\eta_n \rightarrow \eta$, we have $\eta = G(\eta)$. Now I claim that:

Proposition 7.1

If $\varphi \geq 0$ is a solution to $\varphi = G(\varphi)$, then $\eta \leq \varphi$. 

Proof of the proposition. Indeed, $G(s) = E(s^{Z_1})$ is increasing in $[0, +\infty)$ with

$$\eta_1 = G(0) \leq G(\varphi) = \varphi, \quad \eta_2 = G(\eta_1) \leq G(\varphi) = \varphi, \dots$$

and thus $\eta_n = G(\eta_{n-1}) \leq G(\varphi) = \varphi$. Sending $n \rightarrow \infty$, we get the desired. \square

$G(s)$ is increasing in $[0, +\infty)$ such that $G'(1) = \mu$ and $G''(s) = E(Z_1(Z_1 - 1)s^{Z_1-2}) \geq 0$ since Z_1 takes value in \mathbb{N} (that is $G(s)$ is convex).

A glance at Figure 7.1 (and a more analytical verification) tells us that these intersections are coincident if $\mu = G'(1) < 1$. On the other hand, if $\mu > 1$ then these two intersections are not coincident. In the special case when $\mu = 1$ we need to distinguish between the non-random case in which $\sigma^2 = 0$, $G(s) = s$, and $\eta = 0$, and the random case in which $\sigma^2 > 0$, $G(s) > s$ for $0 \leq s < 1$, and $\eta = 1$. \square




Figure 7.1: $\mu > 1$

7.4 Characteristic functions

Definition 7.7

The **characteristic function** of X is defined by

$$\phi(t) = E(e^{itX}) = \int e^{itx} f(x) dx = E(\cos tX) + iE(\sin tX).$$



Example 7.24 For $X \sim \text{Cauchy}(1)$, its characteristic function is

$$\int_{-\infty}^{+\infty} e^{itx} \frac{1}{\pi(1+x^2)} dx = \int_{-\infty}^{+\infty} \frac{\cos tx}{\pi(1+x^2)} dx$$

To start the computation, we define a contour Γ as a semicircle with diameter $[-R, R]$ on the real axis. Next,

we define the function $f(z) = \frac{e^{itz}}{\pi(z^2 + 1)} = \frac{e^{itz}}{\pi(z+i)(z-i)}$. The pole of $f(z)$ lies exactly at i . At $z = i$:

$$g(z) = (z - i)f(z) = \frac{e^{itz}}{\pi(z + i)}$$

is analytic so the pole is simple and

$$\text{Res}(f, i) = g(i) = \frac{e^{-t}}{2\pi i}.$$

By Residue Theorem,

$$\int_{\Gamma} f(z) dz = 2\pi i \text{Res}(f, i) = e^{-t}.$$

Send $R \rightarrow +\infty$, noting that

$$\int_{\mathbb{C}_{\mathbb{R}^+}} f(z) dz = \left| \int_{\mathbb{C}_{\mathbb{R}^+}} \frac{1}{\pi} \frac{e^{itz}}{1+z^2} dz \right| \leq \frac{1}{\pi} \frac{1}{R^2 - 1} \pi R \rightarrow 0.$$

That is the arc of the semicircle has integration 0 and finally

$$\int_{\mathbb{R}} \frac{e^{itx}}{1+x^2} \frac{1}{\pi} dx = e^{-|t|}, \quad \forall t \in \mathbb{R}.$$




Figure 7.2: The contour

Theorem 7.8

The characteristic ϕ satisfies:

1. $\phi(0) = 1, |\phi(t)| \leq 1$ for any $t \in \mathbb{R}$;
2. ϕ is uniformly continuous;
3. ϕ is positive semi-definite: for any $t_1, \dots, t_n \in \mathbb{R}, z_1, \dots, z_n \in \mathbb{C}$,

$$\sum_{i,j} \phi(t_i - t_j) z_i \bar{z}_j \geq 0.$$



Before proving the theorem, we first introduce some consequences in real analysis:

Lemma 7.5 (Fatou)

If $f_n \geq 0$, then

1. $\int \liminf_{n \rightarrow \infty} f_n d\mu \leq \liminf_{n \rightarrow \infty} \int f_n d\mu$;
2. If $\sup f_n \leq f$, and $\int f d\mu < +\infty$, then

$$\int \limsup_{n \rightarrow \infty} f_n d\mu \geq \limsup_{n \rightarrow \infty} \int f_n d\mu.$$



Remark In a probabilistic way: If $X_n \geq 0$, then

$$E(\liminf X_n) \leq \liminf E(X_n).$$

Proof. Note that $\liminf f_n = \sup_{m \geq 1} \inf_{n \geq m} f_n$, then $g_m := \inf_{n \geq m} f_n \uparrow \liminf f_n$. By M.C.T.,

$$\int \lim g_m d\mu = \int \sup_{m \geq 1} g_m = \lim_{m \geq 1} \int g_m d\mu.$$

On the other hand, $\forall n \geq m$,

$$\int g_m d\mu \leq \int f_n d\mu$$

Taking inf with respect to n on both sides, we obtain

$$\int g_m d\mu \leq \inf_{n \geq m} \int f_n d\mu.$$

Send $m \rightarrow +\infty$, getting

$$\int \liminf f_n d\mu = \lim \int g_m d\mu \leq \liminf_{n \rightarrow \infty} \int f_n d\mu$$

□

Remark

1. The equality may not be attained, i.e., there is $\{f_n\}$ such that $\int \liminf f_n < \liminf \int f_n$: Let $f_n(x) = \mathbb{1}_{[n, n+1]}(x)$, then $\liminf f_n = 0$.
2. If $\{f_n\}$ are not non-negative, Fatou's Lemma may fail: Let $f_n(x) = -\mathbb{1}_{[n, n+1]}(x)$.

Theorem 7.9 (Dominated Convergence Theorem(D.C.T))

Let $(\Omega, \mathcal{F}, \mu)$ be a measure space. $\{f_n\}, f$ are measurable functions such that

1. $f_n \rightarrow f$ a.e.;
2. there is an integration function g such that $|f_n| \leq g, \forall n \geq 1$, and $\int g d\mu < +\infty$,

then f is integrable with $\lim_{n \rightarrow \infty} \int f_n d\mu = \int f d\mu$.



Proof. Since $f_n + g \geq 0$, by Fatou,

$$\int \liminf (f_n + g) \leq \liminf \left[\int f_n + g \right] = \liminf \int f_n + \int g,$$

that is,

$$\int f \leq \liminf \int f_n.$$

Similarly, $g - f_n \geq 0$ and by Fatou,

$$\int \liminf (g - f_n) \leq \liminf \left[\int g - f_n \right] = \int g - \limsup \int f_n$$

that is

$$\limsup \int f_n \leq \int f.$$

In conclusion

$$\limsup \int f_n \leq \int f \leq \liminf \int f_n$$

and this reduces to $\lim \int f_n = \int f$. □

Theorem 7.10 (Bounded Convergence)

For $\mu(\Omega) < +\infty$, if

1. $f_n \rightarrow f$ a.e.;
2. there is $K < +\infty$ such that $|f_n| \leq K$ a.e., then $\int f_n = \int f$.



Proof. Put $g = K$ and apply D.C.T. □

Proof of Theorem 7.8. First prove 3. Note that

$$\begin{aligned} \sum_{i,j} E(e^{i(t_i - t_j)X}) z_i \bar{z}_j &= \sum_{i,j} E[e^{it_i X} e^{-it_j X} z_i \bar{z}_j] \\ &= E \left[\sum_{i,j} e^{it_i X} z_i \cdot \overline{e^{it_j X} z_j} \right] \\ &= E \left(\sum_i \left| e^{it_i X} z_i \right|^2 \right) \geq 0. \end{aligned}$$

Continue with 2. Note that

$$|\phi(t+h) - \phi(t)| = |E(e^{i(t+h)X}) - E(e^{itX})| = |E(e^{itX}(e^{ihX} - 1))| \leq E(|e^{ihX} - 1|)$$

By Bounded Convergence, $\lim_{h \rightarrow 0} E(|e^{ihX} - 1|) = 0$.

As for 1, one has

$$|\phi(t)| = \left| \int_{-\infty}^{\infty} e^{itx} f(x) dx \right| \leq \int_{-\infty}^{\infty} |e^{itx}| f(x) dx = \int_{-\infty}^{\infty} f(x) dx = \phi(0) = 1.$$

□

Theorem 7.11 (Bachner)

If ϕ satisfies statement 1, 2 and 3 in Theorem 7.8, then there is a unique probability measure P such that

$$\phi(t) = \int e^{itx} dP.$$

♥

Theorem 7.12

1. If $\phi^{(k)}(0)$ exists, then

$$\phi^{(k)}(0) = i^k E(X^k),$$

from which we see

$$E(X) = \frac{\phi'(0)}{i}, \quad \text{Var}(X) = -\phi''(0) + (\phi'(0))^2.$$

2. If $\phi^{(k)}(0)$ exists, then $\begin{cases} E|X^k| < \infty, & \text{if } k \text{ is even,} \\ E|X^{k-1}| < \infty, & \text{if } k \text{ is odd.} \end{cases}$

3. If $E(|X|^k) < +\infty$, then

$$\phi(t) = \sum_{j=0}^k \frac{(it)^j}{j!} E(X^j) + o(t^k).$$

♥

Theorem 7.13

If X and Y are independent, then $\phi_{X+Y}(t) = E[e^{it(X+Y)}] = \phi_X(t)\phi_Y(t)$.

♥

Lemma 7.6

If $Y = aX + b$ for some $a, b \in \mathbb{R}$, then $\phi_Y(t) = e^{itb}\phi_X(at)$.

♥

Definition 7.8

The joint characteristic function of X and Y is given by

$$\phi(s, t) = E[e^{isX + itY}].$$

♣

Theorem 7.14

X and Y are independent $\iff \phi(s, t) = \phi_X(s)\phi_Y(t)$.

♥

Theorem 7.15 (Analytic extension of $M(t)$)

Let $M(t) = E(e^{tX})$, $t \in \mathbb{R}$, and $\phi(t) = E(e^{itX})$, $t \in \mathbb{C}$, be the moment generating function and characteristic function, respectively, of a random variable X . For any $a > 0$, the following three statements are equivalent:

1. $|M(t)| < +\infty$ for $|t| < a$;
2. $\phi(z) = E(e^{tX})$ is analytic in the strip $|\text{Re}(z)| < a$;

3. The moments $m_k = E(X^k)$, $k \in \mathbb{N}$ exists and $\lim_{k \rightarrow \infty} \left(\frac{m_k}{k!}\right)^{\frac{1}{k}} \leq \frac{1}{a}$.
 If any of these conditions hold for $a > 0$, the power series expansion for $M(t)$ may be extended analytically to the strip $|\operatorname{Im}(t)| < a$, resulting in a function M with the property that $\phi(t) = M(it)$. ♥

7.5 Common characteristic functions

Example 7.25(Delta Measure) We have $f_X(a) = 1$ and $\phi(t) = E(e^{itX}) = e^{iat}$.

Example 7.26(Bernoulli(p)) $\phi(t) = Ee^{itX} = pe^{it} + q$.

Example 7.27(Binomial(n, p)) Note the decomposition $X = Y_1 + \dots + Y_n$ with $Y_i \sim \text{i.i.d. Bernoulli}(p)$.

Then

$$E(e^{itX}) = (E(e^{itY_1}))^n = (pe^{it} + q)^n.$$

Example 7.28(Poisson(λ)) Recall that $f(k) = \frac{\lambda^k}{k!}e^{-\lambda}$, then

$$\phi(t) = E(e^{itX}) = \sum_k e^{itk} \frac{\lambda^k}{k!} e^{-\lambda} = e^{\lambda e^{it} - \lambda}.$$

Example 7.29(Exponential(λ)) Recall that $f(x) = \lambda e^{-\lambda x} \mathbf{1}_{[0, +\infty)}(x)$. Then

$$\phi(t) = E(e^{itX}) = \int_0^{+\infty} e^{itx} \lambda e^{-\lambda x} dx = \frac{\lambda}{\lambda - it}.$$

Example 7.30(Normal) $\phi(t) = E(e^{itX}) = \int_{\mathbb{R}} e^{itx} e^{-\frac{x^2}{2}} dx$ whereas $M(t) = E(e^{tX}) = e^{\frac{1}{2}t^2}$. By analytic extension, $\phi(t) = M(it) = e^{-\frac{1}{2}t^2}$. If $Y \sim \mathcal{N}(\mu, \sigma^2)$, then $Y = \mu + \sigma X$ for some $X \sim \mathcal{N}(0, 1)$ and

$$\phi_Y(t) = e^{i\mu t} E(e^{i\sigma t X}) = e^{i\mu t - \frac{1}{2}\sigma^2 t^2}.$$



Note Indeed, for $X \sim \mathcal{N}(0, 1)$, note that

$$\phi(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} e^{itx} e^{-\frac{x^2}{2}} dx = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} \sum_{n=0}^{\infty} \frac{(itx)^n}{n!} e^{-\frac{x^2}{2}} dx = \sum_{n=0}^{\infty} \frac{(it)^n}{n!} \left[\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} x^n e^{-\frac{x^2}{2}} dx \right]$$

Recall that $E(X^n) = \begin{cases} 0, & n = 2m - 1, \\ (2m - 1)!! = \frac{(2m)!}{2^m \cdot m!}, & n = 2m. \end{cases}$ Thus

$$\phi(t) = \sum_{m=0}^{\infty} \frac{(it)^{2m}}{(2m)!} \cdot \frac{(2m)!}{2^m \cdot m!} = \sum_{m=0}^{\infty} \left(-\frac{t^2}{2}\right)^m \frac{1}{m!} = e^{-\frac{t^2}{2}}.$$

Example 7.31 $(X_1, \dots, X_n) \sim \mathcal{N}(0, V)$ if and only if $\operatorname{Cov}(X_i, X_j) = V_{ij}$. Their joint density function is

$$f(x_1, \dots, x_n) = \frac{1}{\sqrt{(2\pi)^n \det V}} e^{-\frac{1}{2}x^T V^{-1}x}.$$

For $t_1, \dots, t_n \in \mathbb{R}$, the joint characteristic function is

$$E(e^{i(t_1 X_1 + \dots + t_n X_n)}) = \frac{1}{\sqrt{(2\pi)^n \det V}} \int e^{i(t_1 x_1 + \dots + t_n x_n)} e^{-\frac{1}{2}x^T V^{-1}x} dx.$$

Consider the diagonalization of V^{-1} : there is orthogonal matrix B and diagonal matrix $\Lambda = \operatorname{diag}(\lambda_1, \dots, \lambda_n)$ such that $B^T V^{-1} B = \Lambda$. Let $x = By$, then

$$x^T V^{-1} x = y^T B^T V^{-1} B y = y^T \Lambda y = \sum \lambda_i y_i^2,$$

$$\phi(t) = e^{-\frac{1}{2}t^T V t}.$$

Alternatively, let $Y = (Y_1, \dots, Y_n) = AX$, then $Y \sim \mathcal{N}(0, \lambda)$ with Y_1, \dots, Y_n being independent normal. Thus

$$t_1 X_1 + \dots + t_n X_n = t^T X = t^T A^T Y$$

is a linear transformation of independent normal and is therefore normal. Therefore,

$$E(t_1 X_1 + \dots + t_n X_n) = 0, \quad \text{var}(t_1 X_1 + \dots + t_n X_n) = \sum_{i,j} t_i t_j \text{cov}(X_i, X_j) = \sum_{i,j} t_i V_{ij} t_j = t^T V t.$$

Finally,

$$E[e^{i(t_1 X_1 + \dots + t_n X_n)}] = e^{-\frac{1}{2} \text{var}(t_1 X_1 + \dots + t_n X_n)} = e^{-\frac{1}{2} t^T V t}.$$

Example 7.32(Geometric) The characteristics function of geometric distribution is

$$\begin{aligned} \phi(t) &= E(e^{itX}) = \sum_{x=0}^{\infty} e^{itx} P(X=x) \\ &= \sum_{x=1}^{\infty} e^{itx} q^{x-1} p = pq^{-1} \left(\sum_{x=0}^{\infty} (qe^{it})^x - 1 \right) \\ &= pe^{it} (1 - qe^{it})^{-1} \end{aligned}$$

by noting that $\sum_{x=0}^{\infty} q^x = (1-q)^{-1}$.

Example 7.33(Pascal) The Pascal distributed X has density function

$$f(k) = \binom{k-1}{r-1} p^r (1-p)^{k-r}, \quad k = r, r+1, \dots$$

One can decompose X into r i.i.d. geometric(p) random variables, i.e., $X = X_1 + X_2 + \dots + X_r$. Then

$$\phi_X(t) = \prod_{j=1}^r \phi_{X_j}(t) = \left(\frac{pe^{it}}{1 - qe^{it}} \right)^r.$$

7.6 Inversion and continuity

Recall $\phi(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{itx} f(x) dx$. Then by Fourier inversion theorem, the density function is

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-itx} \phi(t) dt \tag{7.1}$$

at every point x such that f is differentiable.

Example 7.34(Revisit to Cauchy distribution) Knowing $\phi(t) = e^{-|t|}$ and apply (7.1), we get

$$\begin{aligned} f(x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-ixt} \cdot e^{-|t|} dt \\ &= \frac{1}{2\pi} \int_0^{\infty} e^{-(1+ix)t} dt + \frac{1}{2\pi} \int_{-\infty}^0 e^{(1-ix)t} dt \\ &= \frac{1}{2\pi} \left(\frac{1}{1+ix} + \frac{1}{1-ix} \right) = \frac{1}{\pi(1+x^2)}. \end{aligned}$$

Thus X is Cauchy distributed.

Theorem 7.16 (Inversion)

Let X have the distribution F and characteristic function ϕ . Let

$$\bar{F}(x) = \frac{1}{2}(F(x) + \lim_{y \uparrow x} F(y)).$$

Then for any $a < b$,

$$\bar{F}(b) - \bar{F}(a) = \lim_{N \rightarrow +\infty} \int_{-N}^N \frac{e^{-iat} - e^{-ibt}}{2\pi it} \phi(t) dt.$$

**Corollary 7.3**

If $\phi_X(t) = \phi_Y(t)$, then $F_X = F_Y$.



Proof. Apply Inversion Theorem with $a = -\infty$: $\bar{F}_X(b) = \bar{F}_Y(b)$. For any $x \in \mathbb{R}$, we use the right continuity of F_X and F_Y . Take $b_n \downarrow x$ such that F_X, F_Y are continuous at b_n . Then

$$\bar{F}_X(b_n) = F_X(b_n) = F_Y(b_n) = \bar{F}_Y(b_n),$$

implying that $F_X(x) = F_Y(x)$. □

Theorem 7.17 (Uniqueness)

Distribution function is uniquely determined by the characteristic function.



Example 7.35 We now have a more elegant way to demonstrate the sum of two independent Gaussian r.v.s is still Gaussian: suppose that $X \sim \mathcal{N}(\mu_1, \sigma_1^2), Y \sim \mathcal{N}(\mu_2, \sigma_2^2)$ are independent, their characteristic functions are

$$\phi_X(t) = e^{it\mu_1 - \frac{1}{2}\sigma_1^2 t^2}, \quad \phi_Y(t) = e^{it\mu_2 - \frac{1}{2}\sigma_2^2 t^2}.$$

Use independence,

$$\phi_{X+Y}(t) = \phi_X(t) \cdot \phi_Y(t) = e^{it(\mu_1 + \mu_2) - \frac{1}{2}(\sigma_1^2 + \sigma_2^2)t^2}$$

From Uniqueness Theorem, one deduces that $X + Y \sim \mathcal{N}(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$.

Exercise 7.9 Let X be a random variable with density function $f(x)$ and characteristic function $\phi(t)$. Show that $f(x)$ is even if and only if $\phi(t)$ is real and even.

Proof. \Leftarrow . We now have $\phi_X(t) = \phi_X(-t) = \phi_{-X}(t)$, i.e., X and $-X$ have the same characteristic function. Thus they have the same distribution and $f_X(x) = f_{-X}(x) = f_X(-x)$.

\Rightarrow . We now deduce that X and $-X$ have the same distribution (and the same characteristic function) from $f(x) = f(-x)$. Furthermore, note that $\phi_{-X}(t) = \phi_X(-t) = \overline{\phi_X(t)}$, implying that $\phi(t)$ is indeed real and even. □

Definition 7.9

Let $\{X_n\}_{n \geq 1}$ and Y be random variables, then we say X_n **converges in distribution** to Y , denoted by $X_n \xrightarrow{D} Y$, if $\lim_{n \rightarrow \infty} P(X_n \leq x) = P(Y \leq x) = F_Y(x)$ for all x such that F_Y is continuous at x .



Remark The continuity assumption at x is to deal with discrete random variables: Let $X_n = x_n \downarrow x := X$. Then $F_{X_n}(x) = 0$ but $F_X(x) = 1$.

Theorem 7.18 (Continuity)

Suppose a sequence of r.v.s $\{X_n\}_{n \geq 1}$ have corresponding characteristic function $\{\phi_n\}_{n \geq 1}$.

1. If $X_n \xrightarrow{D} X$, then $\phi_n(t) \rightarrow \phi(t)$ for all t ;
2. Conversely, if $\phi_n(t) \rightarrow \phi(t)$ for all t , and $\phi(t)$ is continuous at $t = 0$, then ϕ is the characteristic function of some random variable X such that $X_n \xrightarrow{D} X$.

**Theorem 7.19 (Weak law of large numbers)**

Let X_1, X_2, \dots, X_n be a sequence of independent r.v.s such that $E(X_i) = \mu, i = 1, 2, \dots, n$. Then $S_n = X_1 + \dots + X_n$ satisfies $\frac{1}{n}S_n \xrightarrow{D} \mu$.



Proof. Let ϕ_n be the characteristic function of $\frac{1}{n}S_n$.

$$\phi_n(t) = E(e^{it\frac{1}{n}(X_1+\dots+X_n)}) = [E(e^{i\frac{t}{n}X_1})]^n \stackrel{E(X_1) < \infty}{=} \left(1 + i\frac{t}{n}E(X_1) + O\left(\frac{t}{n}\right)\right)^n$$

and

$$\lim_{n \rightarrow \infty} \phi_n(t) = \lim_{n \rightarrow \infty} \left(1 + i\frac{t}{n}\mu + O\left(\frac{t}{n}\right)\right)^n = e^{it\mu}.$$

By continuity theorem, $\frac{1}{n}S_n \xrightarrow{D} \mu$. □



Note The “independent” restriction indeed can be weakened to “uncorrelated”: Let X_1, X_2, \dots, X_n be a sequence of uncorrelated r.v.s with the same distribution such that $E(X_i X_j) = E(X_i)E(X_j), E(X_i^2) < \infty, E(X_i) = \mu, i, j = 1, 2, \dots, n$. Then $S_n = X_1 + \dots + X_n$ satisfies

$$\overline{X}_n := \frac{1}{n}S_n \xrightarrow{P} \mu.$$

In fact, $\frac{1}{n} \sum_{i=1}^n X_i \xrightarrow{L^2} \mu$:

$$E\left(\left(\frac{1}{n} \sum_{i=1}^n X_i - \mu\right)^2\right) = \frac{1}{n^2} \sum_{i,j=1}^n E[(X_i - \mu)(X_j - \mu)] = \frac{1}{n^2} \sum_{i=1}^n \text{var}(X_i) \leq \frac{c}{n}$$

a.s. as $n \rightarrow \infty$.

Theorem 7.20 (Strong law of large numbers)

Let $X_1, \dots, X_n : \Omega \rightarrow \mathbb{R}$ be i.i.d. with mean μ and $E(|X_1|) < \infty$, then $\frac{1}{n}S_n \rightarrow \mu$ a.s.. ♥

Example 7.36 (Monte Carlo simulations: Numerically stimulate π) We generate $U_i, V_i (i = 1, 2, \dots, r)$ which are independent and uniformly $[-1, 1]$ distributed. We stipulate that

$$X_i = \begin{cases} 1, & U_i^2 + V_i^2 \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Then X_i is Bernoulli and $E(X_i) = P(X_i = 1) = \frac{\pi}{4}$. By law of large numbers,

$$\overline{X}_n := \frac{1}{n} \sum_{i=1}^n X_i \xrightarrow{P} E(X_i) = \frac{\pi}{4}.$$

Definition 7.10

Given r.v.s. $\{X_n\}$, a *confident interval* for the theoretical mean μ with **confident level** $\beta\%$ is an interval of length 2ε such that

$$P[\mu \in (\bar{X}_n - \varepsilon, \bar{X}_n + \varepsilon)] \leq \beta\%.$$



Example 7.37 What is the smallest n such that one obtains a four-digit accuracy of π with confident level 99%?

Solution. Note that the stimulation in Example 7.36 is $\frac{\pi}{4}$ and thus the error ε here is $\frac{1}{4000}$ and the confident level is $\beta = 99\%$. We look for n such that

$$P\left(\left|\bar{X}_n - \frac{\pi}{4}\right| > \frac{1}{4000}\right) \leq 1\% \quad (\clubsuit)$$

Note that X_i is Bernoulli(p) and $\text{Var}(X_i) = p(1-p) \leq \frac{1}{4}$. From Chebychev,

$$P\left(\left|\bar{X}_n - \frac{\pi}{4}\right| > \frac{1}{4000}\right) \leq 4000^2 \cdot \text{Var}(\bar{X}_n) = 4000^2 \frac{1}{n^2} \sum_{i=1}^n \text{Var}(X_i) \leq \frac{4000^2}{n} \cdot \frac{1}{4}.$$

Thus if $n \geq 400 \times 10^6$, (\clubsuit) automatically holds. \square

Example 7.38(Revisit to coupon collector) $\Omega = \{1, 2, \dots, n\}$ and X_n be i.i.d. uniformly $\{1, 2, \dots, n\}$. We use $T_i = \inf\{n : |\{X_1, X_2, \dots, X_n\}| = i\}$, the 1st time to collect i different coupons. We have showed that $\frac{E(T_n)}{n \ln n} \rightarrow 1$. Now I claim that

$$\frac{T_n}{\ln n} \xrightarrow{P} 1.$$

Indeed, note that $T_n = \sum_{i=1}^n (T_i - T_{i-1})$ with $(T_i - T_{i-1}) \sim \text{Geo}\left(1 - \frac{i-1}{n}\right)$. We already showed that $E(T_n) = n \ln n + o(n)$ and

$$\begin{aligned} \text{Var}(T_n) &= \sum_{i=1}^n \text{Var}(T_i - T_{i-1}) = \sum_{i=1}^n \frac{\frac{i-1}{n}}{\left(1 - \frac{i-1}{n}\right)^2} \leq \sum_{i=1}^n \frac{1}{\left(1 - \frac{i-1}{n}\right)^2} \\ &= n^2 \sum_{m=1}^n \frac{1}{m^2} \leq cn^2. \end{aligned}$$

In particular, $\frac{\text{Var}(T_n)}{E(T_n)^2} \rightarrow 0$. By extension of WLLN,

$$\frac{T_n - E(T_n)}{n \ln n} \xrightarrow{P} 0$$

and thus $\frac{T_n}{n \ln n} \xrightarrow{P} 1$.

Note Let $T_n = \sum_{i=1}^n X_i$ for every sequence $\{a_n\}$ such that $\frac{\text{Var}(T_n)}{a_n^2} \rightarrow 0$ as $n \rightarrow \infty$. We have $\frac{T_n - E(T_n)}{a_n} \xrightarrow{P} 0$.

Exercise 7.10 Use characteristic function to prove the Central Limit Theorem:

Theorem 7.21

Let $\{X_i\}$ be i.i.d. r.v.s with $E(X_i) = 0$ and $E(X_i^2) = 1$. Assume $E(X_i^3) < +\infty$, then

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n X_i \xrightarrow{D} \mathcal{N}(0, 1).$$



Proof. Compute the characteristic function:

$$\begin{aligned}\phi_n(t) &= E\left[e^{i\frac{t}{\sqrt{n}}\sum_{i=1}^n X_i}\right] = \left(E\left[e^{i\frac{t}{\sqrt{n}}X_1}\right]\right)^n \\ &= \left[1 + i\frac{t}{\sqrt{n}}E(X_1) - \frac{1}{2}\frac{t^2}{n}E(X_1^2) + O\left(\frac{t^3}{n^{3/2}}\right)\right]^n \\ &\xrightarrow{n\rightarrow\infty} e^{-\frac{t^2}{2}}.\end{aligned}$$

□

Chapter 8 Convergence of random variables

8.1 Different modes of convergence

Definition 8.1

Let $\{X_n\}_{n \geq 1}$ be random variables in $(\Omega, \mathcal{F}, \mu \text{ (or } P))$. Then

1. We say X_n converges to X almost surely (a.s.) if $P(\{\omega : \lim_{n \rightarrow \infty} X_n(\omega) = X(\omega)\}) = 1$;
2. We say X_n converges to X in probability, denoted as $X_n \xrightarrow{P} X$ if for any $\varepsilon > 0$, $P(|X_n - X| > \varepsilon) \rightarrow 0$ as $n \rightarrow \infty$;
3. We say X_n converges to X in the p th moment if $E(|X_n|^p) < +\infty$ for any $n \in \mathbb{N}$ and $E(|X_n - X|^p) \rightarrow 0$ as $n \rightarrow \infty$;
4. We say X_n converges to X almost uniformly if for all $\varepsilon > 0$, there is $E \in \mathcal{F} : \mu(E) < \varepsilon$ such that $f_n \rightrightarrows f$ on E^c ;
5. We say X_n converges in distribution to Y , denoted by $X_n \xrightarrow{D} Y$, if $\lim_{n \rightarrow \infty} P(X_n \leq x) = P(Y \leq x) = F_Y(x)$ for all x such that F_Y is continuous at x .



Remark

1. $X_n \rightarrow X$ in p th moment implies $X_n \rightarrow X$ in probability: for any $\varepsilon > 0$, by Chebychev

$$P(|X_n - X| > \varepsilon) \leq \frac{E(|X_n - X|^p)}{\varepsilon^p} \rightarrow 0.$$

2. $X_n \rightarrow X$ a.e. implies $X_n \rightarrow X$ in probability.
3. $X_n \rightarrow X$ a.s. does not imply $X_n \rightarrow X$ in p th moment: consider $f_n(x) = \mathbb{1}_{[n, n+1)}(x)$ with $f_n \rightarrow 0$ a.s. but f_n is not convergent to 0 in probability or p th moment.
4. $X_n \rightarrow X$ in p th moment does not imply $X_n \rightarrow X$ a.s.

Proposition 8.1

$f_n \rightarrow f$ almost uniformly if and only if for all $\varepsilon > 0$, $\lim_{m \rightarrow \infty} \mu\left(\bigcup_{n \geq m} \{|f_n - f| > \varepsilon\}\right) = 0$.



Proof. \implies . If $f_n \rightarrow f$ a.u., then for all $\delta > 0$, there is E with $\mu(E) < \delta$ such that $f_n \rightrightarrows f$ on E^c . Thus for all $\varepsilon > 0$, there is $m \in \mathbb{N}$ such that for all $n \geq m$, one has $|f_n - f| < \varepsilon$ on E^c . For such m , note that

$\bigcup_{n \geq m} \{|f_n - f| > \varepsilon\} \subseteq E$ and thus

$$\mu\left(\bigcup_{n \geq m} \{|f_n - f| > \varepsilon\}\right) \leq \mu(E) < \delta$$

Send $\delta \rightarrow 0$, we have $\lim_{m \rightarrow \infty} \mu\left(\bigcup_{n \geq m} \{|f_n - f| > \varepsilon\}\right) = 0$.

\impliedby . Suppose that for any $k \geq 1$, we have $\lim_{m \rightarrow \infty} \mu\left(\bigcup_{n \geq m} \left\{|f_n - f| > \frac{1}{k}\right\}\right) = 0$. For any $\delta > 0$, we can

find $m = m(\delta, k)$ such that $\mu\left(\bigcup_{n \geq m} \left\{|f_n - f| > \frac{1}{k}\right\}\right) \leq \frac{\delta}{2^k}$. Define

$$E = \bigcup_{k \geq 1} \bigcup_{m \geq m(\delta, k)} \left(\left\{|f_n - f| > \frac{1}{k}\right\}\right)$$

Then $\mu(E) < \delta$ and $f_n \Rightarrow f$ on E^c . □

Corollary 8.1

1. $f_n \rightarrow f$ a.u. $\implies f_n \rightarrow f$ in measure;
2. If $\mu(\Omega) < +\infty$, then $f_n \rightarrow f$ a.e. $\implies f_n \rightarrow f$ in probability (or in measure).

Theorem 8.1

$X_n \rightarrow X$ in probability implies $X_n \rightarrow X$ in distribution.

Proof. Note that for all $a \in \mathbb{R}, \varepsilon > 0$,

$$\{X_n \leq a\} \subseteq \{X \leq a + \varepsilon\} \cup \{|X_n - X| > \varepsilon\}.$$

Thus

$$P(X_n \leq a) \leq P(X \leq a + \varepsilon) + P(|X_n - X| > \varepsilon)$$

and similarly, $\{X \leq a - \varepsilon\} \subseteq \{X_n \leq a\} \cup \{|X_n - X| > \varepsilon\}$ and

$$P(X \leq a - \varepsilon) \leq P(X_n \leq a) + P(|X_n - X| > \varepsilon)$$

Using $X_n \rightarrow X$ in probability and sending $n \rightarrow \infty$, we have $\begin{cases} \limsup P(X_n \leq a) \leq P(X \leq a + \varepsilon) = F_X(a + \varepsilon), \\ \liminf P(X_n \leq a) \geq P(X \leq a - \varepsilon) = F_X(a - \varepsilon). \end{cases}$

If a is continuity point of F_x , then one can send $\varepsilon \rightarrow 0$ to conclude $\lim P(X_n \leq a) = P(X \leq a)$. □

However, convergence in distribution does not imply convergence in probability:

Example 8.1 Let X have probability density function

$$P(X = -1) = \frac{1}{2}, \quad P(X = 1) = \frac{1}{2}$$

and set $X_n = -X$. Note that X_n and X have the same distribution and hence $X_n \xrightarrow{D} X$. However, for $0 < \varepsilon < 2$, one has

$$P(|X_n - X| \geq \varepsilon) = P(2|X| \geq \varepsilon) = 1 \not\rightarrow 0,$$

that is X_n does not converge to X in probability.

Conclusion

1. $f_n \rightarrow f$ almost uniformly $\implies f_n \rightarrow f$ a.e.
2. If $\mu(\Omega) < +\infty$, then $f_n \rightarrow f$ a.e. $\iff f_n \rightarrow f$ a.u. (Egorov).
3. $f_n \rightarrow f$ in probability(measure) $\implies f_n \rightarrow f$ in distribution.




Figure 8.1: Different convergence

Example 8.2 For $X_n \sim \text{Cauchy}(n)$, i.e., $f_n(x) = \frac{n}{(1+n^2x^2)\pi}$, we have $X_n \xrightarrow{D} 0$.

Proof. In fact, $X_n \xrightarrow{P} 0: \forall \varepsilon > 0$,

$$P(|X_n| > \varepsilon) = 2 \int_{\varepsilon}^{\infty} \frac{n}{(1+n^2x^2)\pi} dx = 2 \left(1 - \frac{2}{\pi} \arctan n\varepsilon \right) \xrightarrow{n \rightarrow \infty} 0$$

□

Proposition 8.2

$X_n \xrightarrow{P} X \iff$ for any subsequence $\{X_{n_l}\}$, there is a further subsequence $\{X_{n_k}\}$ such that $X_{n_k} \rightarrow X$ a.u. (or a.s.).



Proof. \implies . For any subsequence $\{X_{n_l}\}$, we have $X_{n_l} \xrightarrow{P} X$ and thus $\forall k \geq 1, P\left(|X_{n_l} - X| > \frac{1}{k}\right) \rightarrow 0$ as $n_l \rightarrow \infty$. Choose $n_k (> n_{k-1})$ such that $P\left(\bigcup_{k \geq m} |X_{n_k} - X| > \frac{1}{k}\right) \leq \frac{1}{2^{m-1}}$. For all $\varepsilon > 0$,

$$P\left(\bigcup_{k \geq m} |X_{n_k} - X| > \varepsilon\right) \leq P\left(\bigcup_{k \geq m} |X_{n_k} - X| > \frac{1}{k}\right) \leq \frac{1}{2^{m-1}} \xrightarrow{m \rightarrow \infty} 0.$$

Then by Proposition 8.1, we obtain $X_{n_k} \rightarrow X$ a.u..

\impliedby . Prove by contradiction. If X_n does not converge to X in probability, there is $\varepsilon_0 > 0, \sigma_0 > 0$ and a subsequence $\{n_i\}$ such that

$$\lim_{i \rightarrow \infty} P(|X_{n_i} - X| > \varepsilon_0) \geq \sigma_0 \quad (\diamond)$$

But by the condition, there is $\{n_{i_j}\}$ such that $X_{n_{i_j}} \rightarrow X (j \rightarrow \infty)$ a.s., implying $X_{n_{i_j}}$ converges to X in probability, contradicting (\diamond) . □

Theorem 8.2 (Sufficient criterion for a.s. convergence)

1. If for all $\varepsilon > 0, \sum_{n \geq 1} P(|X_n - X| > \varepsilon) < +\infty$, then $X_n \rightarrow X$ a.s.
2. If $\{X_n - X\}$ are pairwise independent for some constant X and there is $\varepsilon_k \downarrow 0$, then $\sum_{n \geq 1} P(|X_n - X| > \varepsilon_k) = +\infty, \forall k \implies X_n \not\rightarrow X$ a.s.



Example 8.3 $\{X_n\} \sim$ i.i.d. unif[0, 1], then $Y_n = \min\{X_1, X_2, \dots, X_n\} \rightarrow 0$ a.s..

Proof. Note that

$$P(Y_n > \varepsilon) = P(X_1 > \varepsilon, X_2 > \varepsilon, \dots, X_n > \varepsilon) = (1 - \varepsilon)^n.$$

That is $\sum_{n \geq 1} P(|Y_n| > \varepsilon) < +\infty$. By the sufficient criterion, $Y_n \rightarrow 0$ a.s.. □

Theorem 8.3 (Littlewood three principles)

1. Measurable set is approximately the finite sum of intervals;
2. Measurable function is approximately continuous functions;
3. Convergent sequence of functions is approximately uniformly convergent.



Now we try to demonstrate $X_n \rightarrow X$: for all $\varepsilon > 0, n \geq 1$, let $B_n(\varepsilon) = \{\omega : |X_n(\omega) - X(\omega)| \leq \varepsilon\}$.

Then

$$\begin{aligned} \{X_n \rightarrow X\} &= \{\omega \in \Omega : \forall \varepsilon > 0, \exists N \text{ such that } \forall n \geq N, |X_n(\omega) - X(\omega)| \leq \varepsilon\} \\ &= \bigcap_{\varepsilon > 0} \bigcup_{N \geq 1} \bigcap_{n \geq N} B_n(\varepsilon) = \bigcap_{k \geq 1} \bigcup_{N \geq 1} \bigcap_{n \geq N} B_n\left(\frac{1}{k}\right) \\ &= \bigcap_{k \geq 1} \liminf_{n \rightarrow \infty} B_n\left(\frac{1}{k}\right) \end{aligned}$$

Lemma 8.1

$$\begin{aligned} X_n \rightarrow X \text{ a.s.} &\iff \lim_{k \rightarrow \infty} P\left(\liminf_{n \rightarrow \infty} B_n\left(\frac{1}{k}\right)\right) = 1 \\ &\iff \lim_{k \rightarrow \infty} P\left(\limsup_{n \rightarrow \infty} B_n^c\left(\frac{1}{k}\right)\right) = 0 \\ &\iff \lim_{\varepsilon \rightarrow 0} P(\liminf_{n \rightarrow \infty} B_n^c(\varepsilon)) = 1 \end{aligned}$$



Lemma 8.2 (Borel Cantelli)

Assume that $\sum_{n \in \mathbb{N}} P(A_n) < \infty$; then

$$P(\limsup_{n \rightarrow \infty} A_n) = P(A_n \text{ i.o.}) = 0.$$

Assume that $A_n, n \in \mathbb{N}$, are independent events. If $\sum_{n \in \mathbb{N}} P(A_n) = \infty$, then

$$P(A_n \text{ i.o.}) = 1.$$



Proof.

$$P(\limsup_{n \rightarrow \infty} A_n) = P\left(\bigcap_{m \geq 1} \bigcup_{n \geq m} A_n\right) = \lim_{m \rightarrow +\infty} P\left(\bigcup_{n \geq m} A_n\right) \leq \lim_{m \rightarrow \infty} \sum_{n \geq m} P(A_n)$$

which goes to 0 since $\sum_{n \geq 1} P(A_n) < +\infty$.

$$\begin{aligned} P[(\limsup_{n \rightarrow \infty} A_n)^c] &= P\left(\left(\bigcap_{m \geq 1} \bigcup_{n \geq m} A_n\right)^c\right) = \lim_{m \rightarrow \infty} P\left(\bigcap_{n \geq m} A_n^c\right) = \lim_{m \rightarrow \infty} \prod_{n \geq m} [1 - P(A_n)] \\ &\leq \lim_{m \rightarrow \infty} \prod_{n \geq m} e^{-P(A_n)} = \lim_{m \rightarrow \infty} e^{-\sum_{n \geq m} P(A_n)} = 1. \end{aligned}$$

Under pairwise independence, let $S_n = \sum_{i=1}^n \mathbf{1}_{A_i}$. The first lemma tells us $P(A_n \text{ i.o.}) = P\left(\sum_{i \geq 1} A_i = +\infty\right)$.

From this, $E(S_n) = \sum_{i=1}^n P(A_i) \rightarrow +\infty$ as $n \rightarrow +\infty$. Note that

$$\begin{aligned} \text{var}(S_n) &= \text{var}\left(\sum_{i=1}^n \mathbf{1}_{A_i}\right) = \sum_{i,j=1}^n \text{cov}(\mathbf{1}_{A_i}, \mathbf{1}_{A_j}) \\ &= \sum_{i=1}^n \text{var}(\mathbf{1}_{A_i}) = \sum_{i=1}^n [P(A_i) - P(A_i)^2] \\ &\leq E(S_n). \end{aligned}$$

Then

$$\begin{aligned} P\left(S_n \leq \frac{1}{2}E(S_n)\right) &\leq P\left(|S_n - E(S_n)| > \frac{1}{2}E(S_n)\right) \leq \frac{4}{(E(S_n))^2} \text{var}(S_n) \\ &\leq \frac{4}{(E(S_n))^2} E(S_n) = \frac{4}{E_n} \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

That is $P\left(\sum_{i=1}^{\infty} \mathbf{1}_{A_i} < \infty\right) = 0$. □

Remark

1. Lemma 1 can be extended to $\{A_i\}$ being pairwise independent;
2. The lemma says certain event of infinite set occurs with probability 0 or 1 (Kolmogorov's 0-1 law).

Example 8.4(Extreme values) Let $\{X_i\}$ be i.i.d. obeying $\exp(1)$, i.e., $f_{X_i}(x) = e^{-x} \mathbf{1}_{[0,+\infty)}(x)$, $i = 1, 2, \dots, n$. Let $M_n = \max_{i=1,2,\dots,n} X_i$, then $\limsup \frac{X_n}{\ln n} = 1$ a.s., $\limsup \frac{M_n}{\ln n} = 1$ a.s.

Proof. Note that

$$P(X_n > \gamma \ln n) = \int_{\gamma \ln n}^{+\infty} e^{-x} dx = e^{-\gamma \ln n} = n^{-\gamma}$$

Sum over n , we have

$$\sum_{n \geq 1} n^{-\gamma} \begin{cases} < +\infty, & \gamma > 1, \\ = +\infty, & \gamma \leq 1. \end{cases}$$

By Borel Cantelli, $P(X_n \geq \gamma \ln n \text{ i.o.}) = \begin{cases} 0, & \gamma > 1, \\ 1, & \gamma \leq 1. \end{cases}$ Put γ to be precisely 1, we have $P(X_n > \ln n \text{ i.o.}) = 1$

and it follows that $\limsup \frac{X_n}{\ln n} \geq 1$ a.s. and $\limsup \frac{M_n}{\ln n} \geq 1$ a.s.


Now we continue to prove $\limsup \frac{M_n}{\ln n} \leq 1$ a.s. Fix $\varepsilon > 0$, since $P(X_n > (1 + \varepsilon) \ln n \text{ i.o.}) = 0$, there is $N(\varepsilon) < +\infty$ such that for any $n \geq N(\varepsilon)$, $X_n < (1 + \varepsilon) \ln n$. Therefore

$$\frac{\max_{n > N(\varepsilon)} X_n}{\ln n} \leq 1 + \varepsilon.$$

Also

$$\frac{\max_{n \leq N(\varepsilon)} X_n}{\ln n} \xrightarrow{n \rightarrow \infty} 0$$

These imply that $\limsup \frac{M_n}{\ln n} \leq 1 + \varepsilon$ a.s. Send $\varepsilon \rightarrow 0$ to conclude. □

 **Note** For all $\varepsilon > 0$, $\liminf \frac{M_n}{\ln n} \geq 1 - \varepsilon$ a.s.. To see this, we first split the interval $[0, n]$ with up to n blocks where most of them except one has length $n^{1-\varepsilon}$. For block i ,

$$\begin{aligned} P_i &:= P(M_i > (1 - \varepsilon) \ln i) = P(X_1 > (1 - \varepsilon) \ln i)^i \\ &= \left(1 - \frac{1}{i^{1-\varepsilon}}\right)^i \leq 2 \left(\frac{1}{e}\right)^{i^\varepsilon} \end{aligned}$$

Since $\sum_{n \geq 1} P_n \leq 2 \sum_{n \geq 1} \left(\frac{1}{e}\right)^{n^\varepsilon} < +\infty$, by Borel Cantelli, $P(M_n > (1 - \varepsilon) \ln n \text{ i.o.}) = 0$ and that is

$\liminf \frac{M_n}{\ln n} \geq 1 - \varepsilon$ with probability 1. Up to now, we have proved that $\lim \frac{M_n}{\ln n} = 1$ a.s..

Example 8.5 Let $\{X_n\}$ be i.i.d. coin flip with $P(H) = P(T) = \frac{1}{2}$. Let L_n denote the longest run of heads at time n , representing the maximum number of consecutive heads observed up to that point. Show that $\frac{L_n}{\log_2 n} \rightarrow 1$ a.s..

Proof. We set $l_j :=$ run of head at time j . Then

$$P(l_j = k) = \frac{1}{2^{k+1}}, \quad L_n = \max_{j \leq n} l_j$$

For any $\varepsilon > 0$,

$$P[l_n > (1 + \varepsilon) \log_2 n] = \sum_{k \geq (1 + \varepsilon) \log_2 n} \frac{1}{2^{k+1}} \leq 2^{-(1 + \varepsilon) \log_2 n} = n^{-(1 + \varepsilon)}.$$

Since $\sum_n P[l_n > (1 + \varepsilon) \log_2 n] < \infty$, one concludes that $P[l_n > (1 + \varepsilon) \log_2 n \text{ i.o.}] = 0$, i.e., there is $N(\varepsilon)$ such that for any $n > N(\varepsilon)$, $l_n \leq (1 + \varepsilon) \log_2 n$. Similarly in Example 8.4, one concludes that $\limsup \frac{L_n}{\log_2 n} \leq 1$ a.s..

Conversely, we show that for all $\varepsilon > 0$, $\liminf \frac{L_n}{\log_2 n} \geq 1 - \varepsilon$ a.s. We still apply the “block” technique: separating the interval $[0, n]$ into many blocks, with most of them have length $(1 - \varepsilon) \log_2 n$ except one. Then

$$\begin{aligned} P(\text{all heads in each block}) &= \frac{1}{2^{(1 - \varepsilon) \log_2 n}} = n^{-(1 - \varepsilon)}, \\ P(L_n < (1 - \varepsilon) \log_2 n) &\leq P(\text{all blocks fail}) = \left(1 - \frac{1}{n^{1 - \varepsilon}}\right)^{\frac{n}{(1 - \varepsilon) \log_2 n}} \\ &= \left(1 - \frac{1}{n^{1 - \varepsilon}}\right)^{n^{1 - \varepsilon} \frac{n^\varepsilon}{(1 - \varepsilon) \log_2 n}} \sim e^{\frac{-n^\varepsilon}{(1 - \varepsilon) \log_2 n}} \end{aligned}$$

Hence we get $\sum_n P(L_n < (1 - \varepsilon) \log_2 n) < \infty$ and by B.C, $P(L_n < (1 - \varepsilon) \log_2 n \text{ i.o.}) = 0$, i.e., there is $N(\varepsilon)$ such that for all $n > N(\varepsilon)$, $\frac{L_n}{\log_2 n} \geq 1 - \varepsilon$. \square

 **Exercise 8.1 (Revisit to matching problem)** n people are to pick n hats. We set

$$X_i = \begin{cases} 1, & \text{if the } i\text{th person takes his own hat,} \\ 0, & \text{otherwise,} \end{cases}$$

and $X = X_1 + X_2 + \cdots + X_n$.

1. Try to find $E(X)$ and $\text{Var}(X)$.

2. Conclude that $\frac{X - E(X)}{n} \xrightarrow{P} 0$.

Solution. Note that every X_i have the same distribution but they are not independent:

$$P(X_i = 1) = \frac{1}{n}, \quad P(X_i = 0) = 1 - \frac{1}{n}, \quad i = 1, 2, \dots, n.$$

Thus

$$E(X_i) = \frac{1}{n}, \quad \text{Var}(X_i) = \frac{1}{n} \left(1 - \frac{1}{n}\right), \quad i = 1, 2, \dots, n.$$

Therefore,

$$E(X) = \sum_{i=1}^n E(X_i) = 1$$

and

$$\text{Var}(X) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{i=1}^n \sum_{j=i+1}^n \text{Cov}(X_i, X_j).$$

We now try to compute $\text{Cov}(X_i, X_j)$. By the condition, $X_i X_j$ only takes value in 0, 1 and

$$P(X_i X_j = 1) = P(X_i = 1, X_j = 1) = \frac{1}{n} \cdot \frac{1}{n-1}$$

implying

$$E(X_i X_j) = 0 \cdot P(X_i X_j = 0) + 1 \cdot P(X_i X_j = 1) = \frac{1}{n(n-1)}.$$

Hence

$$\text{Cov}(X_i, X_j) = E(X_i X_j) - E(X_i)E(X_j) = \frac{1}{n(n-1)} - \left(\frac{1}{n}\right)^2 = \frac{1}{n^2(n-1)},$$

$$\text{Var}(X) = \frac{n-1}{n} + 2 \binom{n}{2} \frac{1}{n^2(n-1)} = 1.$$

Now for any $\varepsilon > 0$, by Chebychev,

$$P\left(\frac{X - E(X)}{n} \geq \varepsilon\right) \leq \frac{1}{n^2 \varepsilon^2} \rightarrow 0$$

as $n \rightarrow \infty$. □

8.2 Tail events

Definition 8.2

Let $\{X_n\}_{n \in \mathbb{N}}$ be a sequence of random variables. Define

$$\mathcal{T}_n := \sigma(X_{n+1}, X_{n+2}, \dots), \text{ and } \mathcal{T} := \bigcap_{n \in \mathbb{N}} \mathcal{T}_n.$$

Then \mathcal{T} is called the **tail σ -algebra** of the sequence $\{X_n\}_{n \in \mathbb{N}}$. We can think of it as containing the events describing the limiting behaviour of the sequence. ♣

Theorem 8.4 (Kolmogorov's 0 – 1 law)

Let $\{X_n\}_{n \in \mathbb{N}}$ be a sequence of independent random variables. Then the tail σ -algebra \mathcal{T} of $\{X_n\}_{n \in \mathbb{N}}$ contains only events of probability 0 or 1. ♥

Example 8.6 (Coin flips for infinitely many times) Let $A = \{\text{patterns } \underbrace{HHH \cdots H}_{1000 \text{ times}} \text{ occurs infinitely many times}\}$.

Then $A \in \mathcal{T}$ and by Kolmogorov 0-1, $P(A) = 0$ or 1.

Example 8.7 (Percolation) $P(\text{infinite open clusters}) = 0$ or 1.

References

- [1] R. Durrett and R. Durrett. *Probability: Theory and Examples*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 2019. ISBN: 9781108473682. URL: <https://books.google.com.hk/books?id=b22MDwAAQBAJ>.
- [2] G. Grimmett, D.R.P.E.M.S.G. Grimmett, and D. Stirzaker. *Probability and Random Processes: Fourth Edition*. Oxford University Press, 2020. ISBN: 9780198847601. URL: <https://books.google.com.hk/books?id=1-vqDwAAQBAJ>.
- [3] 茆诗松等. 概率论基础 (第三版). 北京: 中国统计出版社, 2008.
- [4] Pierre Tarrès. *Lecture notes on Probability Limit Theorems*. 2023.
- [5] Wei Wu. *Lecture notes on Honors Theorey of Probability*. 2024.
- [6] 实变函数论. 21世纪数学规划教材, 数学基础课系列. 北京大学出版社, 2016. ISBN: 9787301276471. URL: <https://books.google.com.hk/books?id=m4MixQEACAAJ>.